

## Kinematic Analysis of Relativistic Hypernuclei Decays \*

Olga Majlingova<sup>1</sup>, Vit Sopko<sup>2</sup>,

<sup>1</sup> Faculty of Mechanical Engineering, Department of Technical Mathematics,  
Czech Technical University, Karlovo nam.13, 120 00 Prague, Czech Republic

<sup>2</sup> Faculty of Mechanical Engineering, Department of Applied Physics,  
Czech Technical University, Technicka 4, 16607 Prague, Czech Republic

Results in kinematic analysis of decay relativistic hypernuclei, produced on nuclei beams in Nuclotron experiments in the Joint Institute for Nuclear research, Dubna, are presented. Methods and ways to determine binding energy for hyperon,  $B_\Lambda$ , in hypernucleus  ${}^6_\Lambda\text{H}$  [1], [2] are discussed. Mathematical model developed for identification exclusive channels of the non mesonic weak decays based on four particle decay of hypernuclei  ${}^{10}_\Lambda\text{Be} \rightarrow \alpha + \alpha + n + n$ , [3] is described.

\* This work is supported by grant No 684 077 0029 VZ MSM

[1] J. Lukstins, Nucl. Phys. **A 691**, 491c (2001).

[2] C. Granja *et al.*, this Conference.

[3] L. Majling *et al.*, Nucl. Phys. **A 754**, 154c (2005).