

A SEARCH FOR NEUTRAL HEAVY LEPTONS [☆]

D.I. MEYER ¹, H.K. NGUYEN ², G.S. ABRAMS, M.S. ALAM,
 A. BARBARO-GALTIERI, A.M. BOYARSKI,
 J.M. DORFAN, R. ELY, G.J. FELDMAN, J. FELLER,
 A. FONG, B. GOBBI, G. HANSON, A.D. JOHNSON,
 J.A. JAROS, J.A. KADYK, B.P. KWAN, P. LECOMTE, D. LÜKE ³,
 R.J. MADARAS, J. MARTIN, D.H. MILLER, S.I. PARKER, J.M. PATERSON,
 M.L. PERL, I. PERUZZI ⁴, M. PICCOLO ⁴, T.P. PUN, P. RAPIDIS, M.T. RONAN, R. ROSS,
 W. TANENBAUM, G.H. TRILLING, T. TRIPPE, V. VUILLEMIN ⁵, J.E. WISS ⁶, D. YOUNT
Stanford Linear Accelerator Center, Stanford University, Stanford, California 94305, USA
Lawrence Berkeley Laboratory, University of California, Berkeley, California 94720, USA
Northwestern University, Evanston, Illinois 60201, USA
and University of Hawaii, Honolulu, Hawaii 96822, USA

Received 8 August 1977

In e^+e^- annihilations at SPEAR we have searched for narrow neutral resonances in the $e\pi$ and $\mu\pi$ channels for masses between $0.5 \text{ GeV}/c^2$ and $1.5 \text{ GeV}/c^2$. No evidence for such states has been found. Under standard theoretical assumptions for production cross sections and decay branching ratios of heavy leptons a lower limit of $1.2 \text{ GeV}/c^2$ can be set on the mass.

There has been considerable speculation about the existence of a massive neutral lepton, N, which would couple to the electron in a right-handed doublet as the neutrino does in the left-handed doublet. Since the decay $K^\pm \rightarrow Ne^\pm$ has not been observed, the mass of such a lepton if it exists must be greater than $0.5 \text{ GeV}/c^2$.

One of the main decay modes of such a lepton would be expected to be $e^\pm\pi^\mp$ with a predicted branching ratio [1] which falls from 80% at $M_N = 0.5 \text{ GeV}/c^2$ to 10% at $M_N = 2 \text{ GeV}/c^2$. (There could of course be a similar lepton decaying to $\mu^\pm\pi^\mp$ and in addition μ -e mixing. Therefore wherever ℓ is written in this paper μ or e is implied.)

A possible source of such leptons would be

$e^+e^- \rightarrow \tau^+\tau^-$, τ representing a charged heavy lepton [2] with $m_\tau = 1.9 \text{ GeV}/c^2$. The subsequent decay $\tau = \nu_\tau eN$ could then occur. Since the primary decay of the second τ is expected to be into modes with a single charged particle, the signature for N would be a 4-prong event with at least 2 leptons and a clustering of events at a unique $\ell\pi$ mass.

The data which was used in the search was taken on the SLAC-LBL Magnetic Detector at SPEAR which has been described elsewhere [3]. An electron is identified by a shower in Pb-scintillation counters, the μ by penetration through 20 cm of steel. Electrons and μ 's below $0.6 \text{ GeV}/c$ momentum could not be identified and were not used in the analysis. There is approximately a 15% chance of a hadron being misidentified as either an electron or a muon. The electron detectors cover 0.65 of the total solid angle, the muon detectors 0.45. The resolution for $\ell\pi$ pairs is 40 MeV FWHM at 1.0 GeV and is roughly proportional to the mass.

In the search 170 000 hadron events recorded at total center-of-mass energy $>5 \text{ GeV}$ were used corresponding to an integrated luminosity of $2.3 \times 10^{37} \text{ cm}^{-2}$. Selection was made on events with 3 or 4 observed prongs of which at least 2 had been identified

[☆] Work supported by the Energy Research and Development Administration.

¹ Permanent address: University of Michigan, Ann Arbor, Michigan.

² Permanent address: LPNHE, Université Paris VI, Paris, France.

³ Fellow of Deutsche Forschungsgemeinschaft.

⁴ Permanent address: Laboratori Nazionali di Frascati dell' INFN, Rome, Italy.

⁵ Swiss National Science Foundation Fellow.

⁶ University of Illinois, Urbana, Illinois.




Fig. 1. Number of events detected as a function of $l\pi$ mass.

as lepton candidates; 85% of the events had 4 prongs. The mass was calculated for all lepton-hadron mass combinations assuming the hadron to be a π . Corrections were made for the solid angle acceptance of the detector and the momentum acceptance for leptons. The data are shown in fig. 1. Fig. 2 shows the 90% confidence upper limit for the cross section as a function of the N mass at the average center-of-mass energy of 6.8 GeV.

The experimental upper limit can be compared with that expected if the N exists. It is assumed that τ pair production occurs at a level of one unit of R and hadron production at a level of 4 units of R in agreement with the best experimental knowledge. In addition the branching ratios $B_\tau \equiv \Gamma(\tau \rightarrow \nu_\tau l N) / \Gamma_\tau$ and $B_N \equiv \Gamma(N \rightarrow l\pi) / \Gamma_N$ are needed. These have been estimated theoretically [1, 4]. The estimate of B_τ as a function of mass is essentially a calculation of the reduction in the available phase space for $\tau \rightarrow \nu_\tau e N$ as compared with the 5 competing channels $\tau \rightarrow \nu_\tau e \bar{\nu}$, $\nu_\tau \mu \bar{\nu}_\mu$ and $\nu_\tau u \bar{d}$ with 3 color degrees of freedom. That is $B_\tau = f / (5 + f)$ where f is a factor less than 1 due to kinematics. Similarly in the pertinent region of M_N around 1.3 GeV/ c^2 , $\Gamma_N = \Gamma(e^- e^+ \nu_e) + \Gamma(e^- \mu^+ \nu_\mu) + \Gamma(e^- u \bar{d})$. The $e\pi$ and $e\rho$ channels compete approximately equally for the 60% of the decay modes involving hadrons. Using $B_\tau B_N$ above the theoretically predicted rate is shown in fig. 2.

Since the e 's and μ 's were identified separately, it was also possible to look at the four individual channels $\tau \rightarrow \nu_\tau N e$ with $N \rightarrow e\pi$ or $\mu\pi$ and $\tau \rightarrow \nu_\tau N \mu$ with




Fig. 2. The solid curve gives the 90% confidence upper limit for an $l\pi$ mass peak. The dashed curve is the theoretical estimate based on expected τ and N branching ratios and τ pair production of one unit of R .

$N \rightarrow e\pi$ or $\mu\pi$. The background in the individual channels is of course lower than the combined background but so are the theoretical branching ratio curves. The mass limit which can be set is therefore approximately the same for the individual channels and the combined sample. No evidence is found in any of the reaction channels for an N .

It would be desirable to carry the search for a neutral heavy lepton to higher mass. Probably the most feasible reactions for such a search are $N\bar{N}$ and $N\bar{\nu}$ production in e^+e^- collisions. At PEP and PETRA energies these final states should have sizeable production cross sections [5] through weak coupling if the N exists.

References

- [1] J.L. Rosner, to be published.
- [2] M. Perl et al., Phys. Lett. 63B (1976) 117.
- [3] J.E.-Augustin et al., Phys. Rev. Lett. 34 (1975) 233; F. Vannucci et al., SLAC-PUB-1862 (Phys. Rev. D, to be published).
- [4] H.B. Thacker and J.J. Sakurai, Phys. Lett. 36B (1971) 103; Yung-Su Tsai, Phys. Rev. 4D (1971) 2821.
- [5] B. Humpert and P.N. Scharbach, Phys. Lett. 68B (1977) 71 and SLAC-PUB-1941.