

Seventh International Colloquium on Multiparticle Production, Tutzing, Germany, 1976, edited by J. Benecke *et al.* (Max-Planck-Institut für Physik und Astrophysik, München, Germany, 1976).

²R. Gustafson *et al.*, private communication.

³We thank R. Gustafson *et al.* (the University of Michigan group) for the use of their calorimeter monitor.

⁴To obtain the differential charged-particle multiplicity we assumed that the inelastic cross section for n -A collisions is the same as for p -A collisions, namely,

$46A^{0.69}$ mb. See Ref. 1 for details.

⁵See, for example, J. Koplik and A. H. Mueller, Phys. Rev. D **12**, 3638 (1975); A. Capella and A. Krzywicki, Orsay Report No. LPTPE 77/16, 1976 (unpublished). In addition, see the review of N. N. Nikolaev, to be published.

⁶Similar effects have been reported previously in measurements at different values of y_{lab} . See, for example, J. W. Cronin *et al.*, Phys. Rev. D **11**, 3105 (1975); D. Garbutt *et al.*, Phys. Lett. **67B**, 355 (1977).

Observation of D^0 -Meson Decay into $K^-\pi^+\pi^0$

D. L. Scharre, A. Barbaro-Galtieri, J. M. Dorfan, R. Ely, G. J. Feldman, J. M. Feller, A. Fong, B. Gobbi, G. Hanson, J. A. Jaros, B. P. Kwan, P. Lecomte, A. M. Litke, D. Lüke, R. J. Madaras, J. F. Martin, D. H. Miller, S. I. Parker, M. L. Perl, I. Peruzzi, M. Piccolo, T. P. Pun, P. A. Rapidis, M. T. Ronan, R. R. Ross, T. G. Trippe, V. Vuillemin, and D. E. Yount

Stanford Linear Accelerator Center and Department of Physics, Stanford University, Stanford, California 94305, and Lawrence Berkeley Laboratory and Department of Physics, University of California, Berkeley, California 94720, and Department of Physics and Astronomy, Northwestern University, Evanston, Illinois 60201, and Department of Physics and Astronomy, University of Hawaii, Honolulu, Hawaii 96822

(Received 6 October 1977)

In a sample of multihadron events with a π^0 from e^+e^- annihilation data at 3.77-GeV center-of-mass energy, we observe the decay $D^0 \rightarrow K^-\pi^+\pi^0$ with direct observation of the π^0 . The observed branching fraction is $(12 \pm 6)\%$.

In e^+e^- annihilation data taken at the $\psi(3772)$,¹ we have observed the $K^-\pi^+\pi^0$ decay mode of the D^0 ($\bar{D}^0$). This is the first reported observation of a D decay mode containing a π^0 .

The data were collected with Stanford Linear Accelerator Center—Lawrence Berkeley Laboratory magnetic detector at SPEAR,² augmented by a system of lead-glass counters which replaced one octant of the magnet return yoke³ for improved γ and electron detection. This system (referred to as the LGW) consists of a 2×26 array of lead-glass blocks (used as active converters) 3.3 radiation lengths (X_0) thick, a 14×19 array of lead-glass blocks 10.5 X_0 thick, and three planes of magnetostrictive spark chambers. The fiducial volume of the LGW covers a solid angle of $0.053 \times 4\pi$ sr. This slightly smaller than the solid angle covered by the actual dimensions of the lead-glass counters in order to insure containment of the entire shower resulting from a γ entering the system.

γ 's are identified by energy deposited in active converters which are cleanly separated spatially from the calculated intersection points of charged tracks (identified in the inner detector) with the

surface of the active converter plane. Correlated deposits in the 10.5 X_0 back blocks and spark chambers are used to give complete information on the γ energy and angles. γ 's which convert in the 1 X_0 aluminum magnet coil are tagged by the spark chambers between the coil and active converters, and the appropriate energy-loss correction is made (approximately 55 MeV). If only γ 's with energy greater than 100 MeV are considered, there is essentially no background. Thus, in the remainder of the analysis, this energy cutoff is used. The energy resolution for γ 's of energy less than 1 GeV is approximately described by $\sigma/E = 0.09/E^{1/2}$ (E in GeV) and the angular resolution is $\Delta\theta \approx 0.3^\circ$.

π^0 's are identified by pairs of γ 's in the LGW which reconstruct to have a mass consistent with the π^0 mass. Figure 1(a) shows the $\gamma\gamma$ invariant mass ($M_{\gamma\gamma}$) for a sample of multihadronic events containing two or more γ 's in the LGW. Clear evidence for π^0 production is observed. A cleaner signal which more clearly shows the magnitude and width of the π^0 signal is obtained if the γ -energy cutoff is increased, as can be seen by the shaded region in the histogram in which an ener-




FIG. 1. (a) $\gamma\text{-}\gamma$ invariant-mass distribution for events with two or more γ 's detected in the LGW. A γ -energy cutoff of 150 MeV is required for events in the shaded region. (b) π^0 acceptance in the LGW as a function of π^0 momentum for isotropically produced π^0 's.

gy cutoff of 150 MeV is used.

The π^0 acceptance is calculated by means of a Monte Carlo program in which the effects of the finite size of the lead-glass counters and the effects of shower spreading are included. Figure 1(b) shows the π^0 acceptance as a function of π^0 momentum. The decrease in acceptance at high momentum is due to the shower overlap and small opening angle at high energy so that it is sometimes not possible to separate the two γ 's.

The analysis is based on a sample of approximately 25 000 multihadronic events near the peak of the $\psi(3772)$, in the center-of-mass energy ($E_{c.m.}$) range from 3.76 to 3.79 GeV. The analysis techniques are similar to those described previously as far as the charged particles are concerned.⁴⁻⁶ Charged-particle identification is based on time-of-flight measurements. We have examined invariant-mass combinations which include a kaon and one or more pions in order to search for D -meson decay modes. We consider all particle combinations which include a π^0 observed in the LGW (defined by $0.08 \leq M_{\gamma\gamma} < 0.16$ GeV/ c^2), a kaon (identified by weight greater than




FIG. 2. $K^+\pi^+\pi^0$ invariant mass vs kinematic-fit χ^2 for constrained events. Dashed line indicates χ^2 cutoff.

0.10 as defined in Ref. 4 for charged kaons and by mass cut for K_s^0), and possibly other charged pions.

In order to improve the resolution of the invariant-mass combinations, we use the property that all D mesons produced at the $\psi(3772)$ are produced in the reaction $e^+e^- \rightarrow D\bar{D}$.⁶ Thus it is possible to fit the kinematic variables with the constraint

$$E_D = \frac{1}{2}E_{c.m.},$$

where E_D is the sum of the energies of the decay products. An additional constraint requiring the $\gamma\text{-}\gamma$ invariant mass to be equal to the π^0 mass is also imposed. We find one invariant-mass combination with evidence for D^0 ($\bar{D}^0$) production, $K^+\pi^+\pi^0$. A plot of invariant mass versus kinematic-fit χ^2 for these $K^+\pi^+\pi^0$ combinations is shown in Fig. 2. A clustering of events is observed near the mass of the D^0 with low χ^2 . A cut is made at $\chi^2 = 5$ and these events are shown in Fig. 3 as a function of $K^+\pi^+\pi^0$ mass. There are 9




FIG. 3. $K^+\pi^+\pi^0$ invariant-mass distribution for constrained events with a $\chi^2 = 5$ cutoff.

events near the mass of the D^0 with mean value of $1.861 \text{ GeV}/c^2$. (This is consistent with the measured value⁶ of $1.8633 \pm 0.0009 \text{ GeV}/c^2$.) The observed rms width of about $4 \text{ MeV}/c^2$ is consistent with the expected experimental resolution. The background estimate is 1.7 events in a $20\text{-MeV}/c^2$ mass interval centered on the D^0 mass based on Monte Carlo studies and the event distribution in the region $1.70\text{--}1.85 \text{ GeV}/c^2$. (The probability of observing 9 events as a result of this background is $\sim 7 \times 10^{-5}$.) This leaves an excess of 7.3 ± 3.0 events above background.

Using the $K^-\pi^+\pi^0$ acceptance calculated by Monte Carlo and the measured luminosity of the data sample (1.3 pb^{-1}), the product of the cross section and branching fraction is calculated to be $\sigma B = 1.4 \pm 0.6 \text{ nb}$. As in Ref. 6, we can calculate the absolute branching fraction of this decay mode by assuming that the $\psi(3.772)$ is a state of definite isospin (either 0 or 1) and that its only substantial decay mode is $D\bar{D}$. The absolute branching fraction is $(12 \pm 6)\%$, where a $\pm 20\%$ systematic error is included.

It is also possible to set an upper limit on the $D^0 \rightarrow \bar{K}^0\pi^0$ decay since no events are observed near the D^0 mass and the acceptance is fairly good. A 90%-confidence-level upper limit of 6% is obtained for the absolute branching fraction. Branching-fraction upper limits (90% confidence

level.) for other Cabibbo-favored decay modes involving π^0 's are all greater than 25% because of either poor acceptance or background. The ratio of the $K^-\pi^+\pi^0$ branching fraction to other previously measured branching fractions⁶ is larger than would be expected from the statistical model.⁷

This work was supported primarily by the U. S. Energy Research and Development Administration. Support for individuals came from the listed institutions plus the Deutsche Forschungsgemeinschaft, the Laboratori Nazionali de Frascati dell'Istituto Nazionale di Fisica Nucleare, and the Swiss National Science Foundation.

¹P. A. Rapidis *et al.*, Phys. Rev. Lett. **39**, 526 (1977).

²J.-E. Augustin *et al.*, Phys. Rev. Lett. **34**, 233 (1975).

³A. Barbaro-Galtieri *et al.*, Phys. Rev. Lett. **39**, 1058 (1977).

⁴G. Goldhaber *et al.*, Phys. Rev. Lett. **37**, 255 (1976); I. Peruzzi *et al.*, Phys. Rev. Lett. **37**, 569 (1976).

⁵G. J. Feldman *et al.*, Phys. Rev. Lett. **38**, 1313 (1977).

⁶I. Peruzzi *et al.*, Phys. Rev. Lett. **39**, 1301 (1977).

⁷C. Quigg and J. L. Rosner, Fermilab Report No. Fermilab-Pub-77/60-THY, 1977 (unpublished).

Newly Found Resonance $\Upsilon(9.5)$ and the Charge of the Heavy Quark

T. Hagiwara

The Rockefeller University, New York, New York 10021

and

Yoichi Kazama^(a)

Fermi National Accelerator Laboratory, Batavia, Illinois 60510

and

E. Takasugi

*Fermi National Accelerator Laboratory, Batavia, Illinois 60510, and Department of Physics,
The Ohio State University, Columbus, Ohio 43210*

(Received 8 August 1977)

The newly discovered resonance $\Upsilon(9.5)$ is studied as a bound state of a heavy quark and its antiquark. From the estimate of the production cross section, it is argued that the charge of the constituent quark is likely to be $-\frac{1}{3}$.

A striking enhancement named Υ with the mass of around 9.5 GeV was discovered¹ in the $\mu^+\mu^-$ production in proton-nucleus collisions at Fermi National Accelerator Laboratory. This enhancement seems to consist of at least two resonances. With a two-Gaussian fit the masses are determined to be $9.44 \pm 0.3 \text{ GeV}$ and $10.17 \pm 0.05 \text{ GeV}$, and their decay widths are consistent with zero. The reported $\mu^+\mu^-$