

Measurement of the WW Production Cross Section in $p\bar{p}$ Collisions at $\sqrt{s}=1.96$ TeV

V. M. Abazov,³³ B. Abbott,⁷⁰ M. Abolins,⁶¹ B. S. Acharya,²⁷ M. Adams,⁴⁸ T. Adams,⁴⁶ M. Agelou,¹⁷ J.-L. Agram,¹⁸ S. H. Ahn,²⁹ M. Ahsan,⁵⁵ G. D. Alexeev,³³ G. Alkhalaf,³⁷ A. Alton,⁶⁰ G. Alverson,⁵⁹ G. A. Alves,² M. Anastasoiaie,³² S. Anderson,⁴² B. Andrieu,¹⁶ Y. Arnoud,¹³ A. Askew,⁷⁴ B. Åsman,³⁸ O. Atramentov,⁵³ C. Autermann,²⁰ C. Avila,⁷ F. Badaud,¹² A. Baden,⁵⁷ B. Baldin,⁴⁷ P. W. Balm,³¹ S. Banerjee,²⁷ E. Barberis,⁵⁹ P. Bargassa,⁷⁴ P. Baringer,⁵⁴ C. Barnes,⁴⁰ J. Barreto,² J. F. Bartlett,⁴⁷ U. Bassler,¹⁶ D. Bauer,⁵¹ A. Bean,⁵⁴ S. Beauceron,¹⁶ M. Begel,⁶⁶ A. Bellavance,⁶³ S. B. Beri,²⁶ G. Bernardi,¹⁶ R. Bernhard,^{47,*} I. Bertram,³⁹ M. Bessaçon,¹⁷ R. Beuselinck,⁴⁰ V. A. Bezzubov,³⁶ P. C. Bhat,⁴⁷ V. Bhatnagar,²⁶ M. Binder,²⁴ K. M. Black,⁵⁸ I. Blackler,⁴⁰ G. Blazey,⁴⁹ F. Blekman,³¹ S. Blessing,⁴⁶ D. Bloch,¹⁸ U. Blumenschein,²² A. Boehnlein,⁴⁷ O. Boeriu,⁵² T. A. Bolton,⁵⁵ F. Borchering,⁴⁷ G. Borissov,³⁹ K. Bos,³¹ T. Bose,⁶⁵ A. Brandt,⁷² R. Brock,⁶¹ G. Brooijmans,⁶⁵ A. Bross,⁴⁷ N. J. Buchanan,⁴⁶ D. Buchholz,⁵⁰ M. Buehler,⁴⁸ V. Buescher,²² S. Burdin,⁴⁷ T. H. Burnett,⁷⁶ E. Busato,¹⁶ J. M. Butler,⁵⁸ J. Bystricky,¹⁷ W. Carvalho,³ B. C. K. Casey,⁷¹ N. M. Cason,⁵² H. Castilla-Valdez,³⁰ S. Chakrabarti,²⁷ D. Chakraborty,⁴⁹ K. M. Chan,⁶⁶ A. Chandra,²⁷ D. Chapin,⁷¹ F. Charles,¹⁸ E. Cheu,⁴² L. Chevalier,¹⁷ D. K. Cho,⁶⁶ S. Choi,⁴⁵ T. Christiansen,²⁴ L. Christofek,⁵⁴ D. Claes,⁶³ B. Clément,¹⁸ C. Clément,³⁸ Y. Coadou,⁵ M. Cooke,⁷⁴ W. E. Cooper,⁴⁷ D. Coppage,⁵⁴ M. Corcoran,⁷⁴ J. Coss,¹⁹ A. Cothenet,¹⁴ M.-C. Cousinou,¹⁴ S. Crépe-Renaudin,¹³ M. Cristetiu,⁴⁵ M. A. C. Cummings,⁴⁹ D. Cutts,⁷¹ H. da Motta,² B. Davies,³⁹ G. Davies,⁴⁰ G. A. Davis,⁵⁰ K. De,⁷² P. de Jong,³¹ S. J. de Jong,³² E. De La Cruz-Burelo,³⁰ C. De Oliveria Martins,³ S. Dean,⁴¹ F. Déliot,¹⁷ P. A. Delsart,¹⁹ M. Demarteau,⁴⁷ R. Demina,⁶⁶ P. Demine,¹⁷ D. Denisov,⁴⁷ S. P. Denisov,³⁶ S. Desai,⁶⁷ H. T. Diehl,⁴⁷ M. Diesburg,⁴⁷ M. Doidge,³⁹ H. Dong,⁶⁷ S. Doulas,⁵⁹ L. Duflot,¹⁵ S. R. Dugad,²⁷ A. Duperrin,¹⁴ J. Dyer,⁶¹ A. Dyshkant,⁴⁹ M. Eads,⁴⁹ D. Edmunds,⁶¹ T. Edwards,⁴¹ J. Ellison,⁴⁵ J. Elmsheuser,²⁴ J. T. Eltzroth,⁷² V. D. Elvira,⁴⁷ S. Eno,⁵⁷ P. Ermolov,³⁵ O. V. Eroshin,³⁶ J. Estrada,⁴⁷ D. Evans,⁴⁰ H. Evans,⁶⁵ A. Evdokimov,³⁴ V. N. Evdokimov,³⁶ J. Fast,⁴⁷ S. N. Fatakia,⁵⁸ L. Felgioni,⁵⁸ T. Ferbel,⁶⁶ F. Fiedler,²⁴ F. Filthaut,³² W. Fisher,⁶⁴ H. E. Fisk,⁴⁷ M. Fortner,⁴⁹ H. Fox,²² W. Freeman,⁴⁷ S. Fu,⁴⁷ S. Fuess,⁴⁷ T. Gadfort,⁷⁶ C. F. Galea,³² E. Gallas,⁴⁷ E. Galyaev,⁵² C. Garcia,⁶⁶ A. Garcia-Bellido,⁷⁶ J. Gardner,⁵⁴ V. Gavrilov,³⁴ P. Gay,¹² D. Gelé,¹⁸ R. Gelhaus,⁴⁵ K. Genser,⁴⁷ C. E. Gerber,⁴⁸ Y. Gershtein,⁷¹ G. Ginther,⁶⁶ T. Golling,²¹ B. Gómez,⁷ K. Gounder,⁴⁷ A. Goussiou,⁵² P. D. Grannis,⁶⁷ S. Greder,¹⁸ H. Greenlee,⁴⁷ Z. D. Greenwood,⁵⁶ E. M. Gregores,⁴ Ph. Gris,¹² J.-F. Grivaz,¹⁵ L. Groer,⁶⁵ S. Grünendahl,⁴⁷ M. W. Grünewald,²⁸ S. N. Gurzhiev,³⁶ G. Gutierrez,⁴⁷ P. Gutierrez,⁷⁰ A. Haas,⁶⁵ N. J. Hadley,⁵⁷ S. Hagopian,⁴⁶ I. Hall,⁷⁰ R. E. Hall,⁴⁴ C. Han,⁶⁰ L. Han,⁴¹ K. Hanagaki,⁴⁷ K. Harder,⁵⁵ R. Harrington,⁵⁹ J. M. Hauptman,⁵³ R. Hauser,⁶¹ J. Hays,⁵⁰ T. Hebbeker,²⁰ D. Hedin,⁴⁹ J. M. Heinmiller,⁴⁸ A. P. Heinson,⁴⁵ U. Heintz,⁵⁸ C. Hensel,⁵⁴ G. Hesketh,⁵⁹ M. D. Hildreth,⁵² R. Hirosky,⁷⁵ J. D. Hobbs,⁶⁷ B. Hoeneisen,¹¹ M. Hohlfeld,²³ S. J. Hong,²⁹ R. Hooper,⁷¹ P. Houben,³¹ Y. Hu,⁶⁷ J. Huang,⁵¹ I. Iashvili,⁴⁵ R. Illingworth,⁴⁷ A. S. Ito,⁴⁷ S. Jabeen,⁵⁴ M. Jaffré,¹⁵ S. Jain,⁷⁰ V. Jain,⁶⁸ K. Jakobs,²² A. Jenkins,⁴⁰ R. Jesik,⁴⁰ K. Johns,⁴² M. Johnson,⁴⁷ A. Jonckheere,⁴⁷ P. Jonsson,⁴⁰ H. Jöstlein,⁴⁷ A. Juste,⁴⁷ M. M. Kado,⁴³ D. Käfer,²⁰ W. Kahl,⁵⁵ S. Kahn,⁶⁸ E. Kajfasz,¹⁴ A. M. Kalinin,³³ J. Kalk,⁶¹ D. Karmanov,³⁵ J. Kasper,⁵⁸ D. Kau,⁴⁶ R. Kehoe,⁷³ S. Kermiche,¹⁴ S. Kesisoglou,⁷¹ A. Khanov,⁶⁶ A. Kharchilava,⁵² Y. M. Kharzheev,³³ K. H. Kim,²⁹ B. Klima,⁴⁷ M. Klute,²¹ J. M. Kohli,²⁶ M. Kopal,⁷⁰ V. M. Korablev,³⁶ J. Kotcher,⁶⁸ B. Kothari,⁶⁵ A. Koubarovsky,³⁵ A. V. Kozelov,³⁶ J. Kozminski,⁶¹ S. Krzywdzinski,⁴⁷ S. Kuleshov,³⁴ Y. Kulik,⁴⁷ S. Kunori,⁵⁷ A. Kupco,¹⁷ T. Kurča,¹⁹ S. Lager,³⁸ N. Lahrichi,¹⁷ G. Landsberg,⁷¹ J. Lazoflores,⁴⁶ A.-C. Le Bihan,¹⁸ P. Lebrun,¹⁹ S. W. Lee,²⁹ W. M. Lee,⁴⁶ A. Leflat,³⁵ F. Lehner,^{47,*} C. Leonidopoulos,⁶⁵ P. Lewis,⁴⁰ J. Li,⁷² Q. Z. Li,⁴⁷ J. G. R. Lima,⁴⁹ D. Lincoln,⁴⁷ S. L. Linn,⁴⁶ J. Linnemann,⁶¹ V. V. Lipaev,³⁶ R. Lipton,⁴⁷ L. Lobo,⁴⁰ A. Lobodenko,³⁷ M. Lokajicek,¹⁰ A. Lounis,¹⁸ H. J. Lubatti,⁷⁶ L. Lueking,⁴⁷ M. Lynker,⁵² A. L. Lyon,⁴⁷ A. K. A. Maciel,⁴⁹ R. J. Madaras,⁴³ P. Mättig,²⁵ A. Magerkurth,⁶⁰ A.-M. Magnan,¹³ N. Makovec,¹⁵ P. K. Mal,²⁷ S. Malik,⁵⁶ V. L. Malyshev,³³ H. S. Mao,⁶ Y. Maravin,⁴⁷ M. Martens,⁴⁷ S. E. K. Mattingly,⁷¹ A. A. Mayorov,³⁶ R. McCarthy,⁶⁷ R. McCroskey,⁴² D. Meder,²³ H. L. Melanson,⁴⁷ A. Melnitchouk,⁶² M. Merkin,³⁵ K. W. Merritt,⁴⁷ A. Meyer,²⁰ H. Miettinen,⁷⁴ D. Mihalcea,⁴⁹ J. Mitrevski,⁶⁵ N. Mokhov,⁴⁷ J. Molina,³ N. K. Mondal,²⁷ H. E. Montgomery,⁴⁷ R. W. Moore,⁵ G. S. Muanza,¹⁹ M. Mulders,⁴⁷ Y. D. Mutaf,⁶⁷ E. Nagy,¹⁴ M. Narain,⁵⁸ N. A. Naumann,³² H. A. Neal,⁶⁰ J. P. Negret,⁷ S. Nelson,⁴⁶ P. Neustroev,³⁷ C. Noeding,²² A. Nomerotski,⁴⁷ S. F. Novaes,⁴ T. Nunnemann,²⁴ E. Nurse,⁴¹ V. O'Dell,⁴⁷ D. C. O'Neil,⁵ V. Oguri,³ N. Oliveira,³ N. Oshima,⁴⁷ G. J. Otero y Garzón,⁴⁸ P. Padley,⁷⁴ N. Parashar,⁵⁶ J. Park,²⁹ S. K. Park,²⁹ J. Parsons,⁶⁵ R. Partridge,⁷¹ N. Parua,⁶⁷ A. Patwa,⁶⁸ P. M. Perea,⁴⁵ E. Perez,¹⁷ O. Peters,³¹ P. Pétrouff,¹⁵ M. Petteini,⁴⁰ L. Phaf,³¹ R. Piegaia,¹ P. L. M. Podesta-Lerma,³⁰ V. M. Podstavkov,⁴⁷ Y. Pogorelov,⁵² B. G. Pope,⁶¹ W. L. Prado da Silva,³ H. B. Prosper,⁴⁶ S. Protopopescu,⁶⁸ M. B. Przybycien,^{50,†} J. Qian,⁶⁰ A. Quadt,²¹ B. Quinn,⁶² K. J. Rani,²⁷ P. A. Rapidis,⁴⁷ P. N. Ratoff,³⁹ N. W. Reay,⁵⁵ S. Reucroft,⁵⁹ M. Rijssenbeek,⁶⁷

I. Ripp-Baudot,¹⁸ F. Rizatdinova,⁵⁵ C. Royon,¹⁷ P. Rubinov,⁴⁷ R. Ruchti,⁵² G. Sajot,¹³ A. Sánchez-Hernández,³⁰ M. P. Sanders,⁴¹ A. Santoro,³ G. Savage,⁴⁷ L. Sawyer,⁵⁶ T. Scanlon,⁴⁰ R. D. Schamberger,⁶⁷ H. Schellman,⁵⁰ P. Schieferdecker,²⁴ C. Schmitt,²⁵ A. A. Schukin,³⁶ A. Schwartzman,⁶⁴ R. Schwienhorst,⁶¹ S. Sengupta,⁴⁶ H. Severini,⁷⁰ E. Shabalina,⁴⁸ M. Shamim,⁵⁵ V. Shary,¹⁷ W. D. Shephard,⁵² D. Shpakov,⁵⁹ R. A. Sidwell,⁵⁵ V. Simak,⁹ V. Sirotenko,⁴⁷ P. Skubic,⁷⁰ P. Slattery,⁶⁶ R. P. Smith,⁴⁷ K. Smolek,⁹ G. R. Snow,⁶³ J. Snow,⁶⁹ S. Snyder,⁶⁸ S. Söldner-Rembold,⁴¹ X. Song,⁴⁹ Y. Song,⁷² L. Sonnenschein,⁵⁸ A. Sopczak,³⁹ M. Sosebee,⁷² K. Soustruznik,⁸ M. Souza,² B. Spurlock,⁷² N. R. Stanton,⁵⁵ J. Stark,¹³ J. Steele,⁵⁶ G. Steinbrück,⁶⁵ K. Stevenson,⁵¹ V. Stolin,³⁴ A. Stone,⁴⁸ D. A. Stoyanova,³⁶ J. Strandberg,³⁸ M. A. Strang,⁷² M. Strauss,⁷⁰ R. Ströhmer,²⁴ M. Strovink,⁴³ L. Stutte,⁴⁷ S. Sumowidagdo,⁴⁶ A. Sznajder,³ M. Talby,¹⁴ P. Tamburello,⁴² W. Taylor,⁵ P. Telford,⁴¹ J. Temple,⁴² S. Tentindo-Repond,⁴⁶ E. Thomas,¹⁴ B. Thooris,¹⁷ M. Tomoto,⁴⁷ T. Toole,⁵⁷ J. Torborg,⁵² S. Towers,⁶⁷ T. Trefzger,²³ S. Trincaz-Duvoid,¹⁶ B. Tuchming,¹⁷ C. Tully,⁶⁴ A. S. Turcot,⁶⁸ P. M. Tuts,⁶⁵ L. Uvarov,³⁷ S. Uvarov,³⁷ S. Uzunyan,⁴⁹ B. Vachon,⁵ R. Van Kooten,⁵¹ W. M. van Leeuwen,³¹ N. Varelas,⁴⁸ E. W. Varnes,⁴² I. A. Vasilyev,³⁶ M. Vaupel,²⁵ P. Verdier,¹⁵ L. S. Vertogradov,³³ M. Verzocchi,⁵⁷ F. Villeneuve-Seguirer,⁴⁰ J.-R. Vlimant,¹⁶ E. Von Toerne,⁵⁵ M. Vreeswijk,³¹ T. Vu Anh,¹⁵ H. D. Wahl,⁴⁶ R. Walker,⁴⁰ L. Wang,⁵⁷ Z.-M. Wang,⁶⁷ J. Warchol,⁵² M. Warsinsky,²¹ G. Watts,⁷⁶ M. Wayne,⁵² M. Weber,⁴⁷ H. Weerts,⁶¹ M. Wegner,²⁰ N. Wermes,²¹ A. White,⁷² V. White,⁴⁷ D. Whiteson,⁴³ D. Wicke,⁴⁷ D. A. Wijngaarden,³² G. W. Wilson,⁵⁴ S. J. Wimpenny,⁴⁵ J. Wittlin,⁵⁸ M. Wobisch,⁴⁷ J. Womersley,⁴⁷ D. R. Wood,⁵⁹ T. R. Wyatt,⁴¹ Q. Xu,⁶⁰ N. Xuan,⁵² R. Yamada,⁴⁷ M. Yan,⁵⁷ T. Yasuda,⁴⁷ Y. A. Yatsunenko,³³ Y. Yen,²⁵ K. Yip,⁶⁸ S. W. Youn,⁵⁰ J. Yu,⁷² A. Yurkewicz,⁶¹ A. Zabi,¹⁵ A. Zatserklyaniy,⁴⁹ M. Zdrzil,⁶⁷ C. Zeitnitz,²³ D. Zhang,⁴⁷ X. Zhang,⁷⁰ T. Zhao,⁷⁶ Z. Zhao,⁶⁰ B. Zhou,⁶⁰ J. Zhu,⁵⁷ M. Zielinski,⁶⁶ D. Zieminska,⁵¹ A. Zieminski,⁵¹ R. Zitoun,⁶⁷ V. Zutshi,⁴⁹ E. G. Zverev,³⁵ and A. Zylberstein¹⁷

(D0 Collaboration)

¹Universidad de Buenos Aires, Buenos Aires, Argentina²LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil³Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil⁴Instituto de Física Teórica, Universidade Estadual Paulista, São Paulo, Brazil⁵Simon Fraser University, Burnaby, Canada; University of Alberta, Edmonton, Canada; McGill University, Montreal, Canada; and York University, Toronto, Canada⁶Institute of High Energy Physics, Beijing, People's Republic of China⁷Universidad de los Andes, Bogotá, Colombia⁸Charles University, Center for Particle Physics, Prague, Czech Republic⁹Czech Technical University, Prague, Czech Republic¹⁰Institute of Physics, Academy of Sciences, Center for Particle Physics, Prague, Czech Republic¹¹Universidad San Francisco de Quito, Quito, Ecuador¹²Laboratoire de Physique Corpusculaire, IN2P3-CNRS, Université Blaise Pascal, Clermont-Ferrand, France¹³Laboratoire de Physique Subatomique et de Cosmologie, IN2P3-CNRS, Université de Grenoble I, Grenoble, France¹⁴CPPM, IN2P3-CNRS, Université de la Méditerranée, Marseille, France¹⁵Laboratoire de l'Accélérateur Linéaire, IN2P3-CNRS, Orsay, France¹⁶LPNHE, Universités Paris VI and VII, IN2P3-CNRS, Paris, France¹⁷DAPNIA/Service de Physique des Particules, CEA, Saclay, France¹⁸IReS, IN2P3-CNRS, Université Louis Pasteur, Strasbourg, France, and Université de Haute Alsace, Mulhouse, France¹⁹Institut de Physique Nucléaire de Lyon, IN2P3-CNRS, Université Claude Bernard, Villeurbanne, France²⁰RWTH Aachen, III. Physikalisches Institut A, Aachen, Germany²¹Universität Bonn, Physikalisches Institut, Bonn, Germany²²Universität Freiburg, Physikalisches Institut, Freiburg, Germany²³Universität Mainz, Institut für Physik, Mainz, Germany²⁴Ludwig-Maximilians-Universität München, München, Germany²⁵Fachbereich Physik, University of Wuppertal, Wuppertal, Germany²⁶Panjab University, Chandigarh, India²⁷Tata Institute of Fundamental Research, Mumbai, India²⁸University College Dublin, Dublin, Ireland²⁹Korea Detector Laboratory, Korea University, Seoul, Korea³⁰CINVESTAV, Mexico City, Mexico³¹FOM-Institute NIKHEF and University of Amsterdam/NIKHEF, Amsterdam, The Netherlands³²University of Nijmegen/NIKHEF, Nijmegen, The Netherlands³³Joint Institute for Nuclear Research, Dubna, Russia³⁴Institute for Theoretical and Experimental Physics, Moscow, Russia

- ³⁵Moscow State University, Moscow, Russia
³⁶Institute for High Energy Physics, Protvino, Russia
³⁷Petersburg Nuclear Physics Institute, St. Petersburg, Russia
³⁸Lund University, Lund, Sweden; Royal Institute of Technology and Stockholm University, Stockholm, Sweden;
and Uppsala University, Uppsala, Sweden
³⁹Lancaster University, Lancaster, United Kingdom
⁴⁰Imperial College, London, United Kingdom
⁴¹University of Manchester, Manchester, United Kingdom
⁴²University of Arizona, Tucson, Arizona 85721, USA
⁴³Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA
⁴⁴California State University, Fresno, California 93740, USA
⁴⁵University of California, Riverside, California 92521, USA
⁴⁶Florida State University, Tallahassee, Florida 32306, USA
⁴⁷Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA
⁴⁸University of Illinois at Chicago, Chicago, Illinois 60607, USA
⁴⁹Northern Illinois University, DeKalb, Illinois 60115, USA
⁵⁰Northwestern University, Evanston, Illinois 60208, USA
⁵¹Indiana University, Bloomington, Indiana 47405, USA
⁵²University of Notre Dame, Notre Dame, Indiana 46556, USA
⁵³Iowa State University, Ames, Iowa 50011, USA
⁵⁴University of Kansas, Lawrence, Kansas 66045, USA
⁵⁵Kansas State University, Manhattan, Kansas 66506, USA
⁵⁶Louisiana Tech University, Ruston, Louisiana 71272, USA
⁵⁷University of Maryland, College Park, Maryland 20742, USA
⁵⁸Boston University, Boston, Massachusetts 02215, USA
⁵⁹Northeastern University, Boston, Massachusetts 02115, USA
⁶⁰University of Michigan, Ann Arbor, Michigan 48109, USA
⁶¹Michigan State University, East Lansing, Michigan 48824, USA
⁶²University of Mississippi, University, Mississippi 38677, USA
⁶³University of Nebraska, Lincoln, Nebraska 68588, USA
⁶⁴Princeton University, Princeton, New Jersey 08544, USA
⁶⁵Columbia University, New York, New York 10027, USA
⁶⁶University of Rochester, Rochester, New York 14627, USA
⁶⁷State University of New York, Stony Brook, New York 11794, USA
⁶⁸Brookhaven National Laboratory, Upton, New York 11973, USA
⁶⁹Langston University, Langston, Oklahoma 73050, USA
⁷⁰University of Oklahoma, Norman, Oklahoma 73019, USA
⁷¹Brown University, Providence, Rhode Island 02912, USA
⁷²University of Texas, Arlington, Texas 76019, USA
⁷³Southern Methodist University, Dallas, Texas 75275, USA
⁷⁴Rice University, Houston, Texas 77005, USA
⁷⁵University of Virginia, Charlottesville, Virginia 22901, USA
⁷⁶University of Washington, Seattle, Washington 98195, USA
- (Received 26 October 2004; published 20 April 2005)

We present a measurement of the W boson pair-production cross section in $p\bar{p}$ collisions at a center-of-mass energy of $\sqrt{s} = 1.96$ TeV. The data, collected with the Run II D0 detector at Fermilab, correspond to an integrated luminosity of 224–252 pb^{-1} depending on the final state (ee , $e\mu$, or $\mu\mu$). We observe 25 candidates with a background expectation of $8.1 \pm 0.6(\text{stat}) \pm 0.6(\text{syst}) \pm 0.5(\text{lum})$ events. The probability for an upward fluctuation of the background to produce the observed signal is 2.3×10^{-7} , equivalent to 5.2 standard deviations. The measurement yields a cross section of $13.8_{-3.8}^{+4.3}(\text{stat})_{-0.9}^{+1.2}(\text{syst}) \pm 0.9(\text{lum})$ pb, in agreement with predictions from the standard model.

DOI: 10.1103/PhysRevLett.94.151801

PACS numbers: 13.38.Be, 13.85.Qk, 14.70.Fm

The measurement of the W boson pair-production cross section $\sigma_{p\bar{p} \rightarrow W^+W^-}$ offers a good opportunity to test the non-Abelian structure of the standard model (SM). Furthermore, W pair production could be enhanced by new phenomena, such as anomalous trilinear couplings

[1], or the production and decay of new particles, such as the Higgs boson [2]. The next-to-leading order (NLO) calculations for $\sigma_{p\bar{p} \rightarrow W^+W^-}$ [3] predict a cross section of 12.0–13.5 pb at $\sqrt{s} = 1.96$ TeV. The CDF Collaboration reported evidence for W boson pair production, based on

108 pb⁻¹ of data collected in Run I of the Fermilab Tevatron Collider at $\sqrt{s} = 1.8$ TeV, with a cross section $\sigma_{p\bar{p} \rightarrow W^+W^-} = 10.2_{-5.1}^{+6.3}(\text{stat}) \pm 1.6(\text{syst})$ pb [4]. The four experiments at the CERN e^+e^- Collider (LEP) have observed W boson pair production in e^+e^- collisions [5]. The probed mass range of the W boson pairs at the Tevatron Collider is much higher than at LEP because of the much higher accessible energies.

In this Letter we present a measurement of the W^+W^- production cross section in leptonic final states $p\bar{p} \rightarrow W^+W^- \rightarrow \ell^+\nu\ell^-\bar{\nu}(\ell = e, \mu)$. We use data collected between April 2002 and March 2004 in $p\bar{p}$ of Run II of the Tevatron Collider. The integrated luminosities are 252 ± 16 pb⁻¹, 235 ± 15 pb⁻¹, and 224 ± 15 pb⁻¹ for the e^+e^- , $e^\pm\mu^\mp$, and $\mu^+\mu^-$ channels, respectively. The differences in the integrated luminosities for various channels are primarily due to different triggering conditions.

We briefly describe the main components of the D0 Run II detector [6] important to this analysis. The D0 detector has a magnetic central-tracking system, consisting of a silicon microstrip tracker and a central fiber tracker, both located within a 2 T superconducting solenoidal magnet [6]. A liquid-argon and uranium calorimeter has a central section (CC) covering pseudorapidities $|\eta|$ up to ≈ 1.1 [$\eta = -\ln(\tan\frac{\theta}{2})$ with polar angle θ], and two end calorimeters extending coverage to $|\eta| \approx 4.2$, all three housed in separate cryostats [7]. A muon system resides beyond the calorimetry, and consists of a layer of tracking detectors and scintillation trigger counters before 1.8 T toroids, followed by two more similar layers after the toroids.

The $W^+W^- \rightarrow \ell^+\nu\ell^-\bar{\nu}$ candidates are selected by triggering on single or di-lepton events using a three level trigger system. The first trigger level uses hardware to select electron candidates based on energy deposition in the electromagnetic part of the calorimeter and selects muon candidates formed by hits in two layers of the muon scintillator system. Digital signal processors in the second trigger level form muon track candidate segments defined by hits in the muon drift chambers and scintillators. At the third level, software algorithms running on a computing farm and exploiting the full event information are used to make the final selection of events which are recorded for off-line analysis.

In further off-line analysis electrons are identified by electromagnetic showers in the calorimeter. These showers are chosen by comparing the longitudinal and transverse shower profiles to those of simulated electrons. The showers must be isolated, deposit most of their energy in the electromagnetic part of the calorimeter, and pass a likelihood criterion that includes a spatial track match and, in the CC region, an E/p requirement, where E is the energy of the calorimeter cluster and p is the momentum of the track. All electrons are required to be in the pseudorapidity range $|\eta| < 3.0$. The transverse momentum measurement

of the electrons is based on calorimeter cell energy information.

To select isolated muons, the scalar sum of the transverse momentum of all tracks other than that of the muon in a cone of $\mathcal{R} = 0.5$ around the muon track must be less than 4 GeV, where $\mathcal{R} = \sqrt{(\Delta\phi)^2 + (\Delta\eta)^2}$ and ϕ is the azimuthal angle. Muon detection is restricted to the coverage of the muon system $|\eta| < 2.0$. Muons from cosmic rays are rejected by requiring a timing criterion on the hits in the scintillator layers as well as applying restrictions on the position of the muon track with respect to the primary vertex.

The decay of two W bosons into electrons or muons results in three different final states $e^+e^- + X$ (ee channel), $e^\pm\mu^\mp + X$ ($e\mu$ channel), and $\mu^+\mu^- + X$ ($\mu\mu$ channel), each of which consists of two oppositely charged isolated high transverse momentum, p_T , leptons and large missing transverse energy, \cancel{E}_T , due to the escaping neutrinos. The selection criteria for each channel were chosen to maximize the expected signal significance, while keeping high efficiency for WW production.

In all three channels, two leptons originating from the same vertex are required to be of opposite charge, and must have $p_T > 20$ GeV for the leading lepton and $p_T > 15$ GeV for the trailing one. Figure 1 shows the good agreement between data and Monte Carlo (MC) calculations in \cancel{E}_T distributions for the ee channel (a), the $\mu\mu$ channel (c), and the $e\mu$ channel (e) after applying the lepton transverse momentum cuts. In all cases, the background is largely dominated by Z/γ^* production which is suppressed by requiring the \cancel{E}_T to be greater than 30, 40, and 20 GeV in the ee , $\mu\mu$, and $e\mu$ channels, respectively. The different cut values among the three channels are due to the different momentum resolution of electrons and muons.

In the ee channel, additional cuts are applied to further reduce the Z/γ^* background and other backgrounds. The minimal transverse mass $m_T^{\min} = \min(m_T^{e1}, m_T^{e2})$ must exceed 60 GeV, where $m_T = \sqrt{2\cancel{E}_T p_T^e [1 - \cos\Delta\phi(p_T^e, \cancel{E}_T)]}$. Events are removed if the invariant di-electron mass is between 76 and 106 GeV. Events are also removed if the \cancel{E}_T has a large contribution from the mismeasurement of jet energy, using the following procedure. The fluctuation in the measurement of jet energy in the transverse plane can be approximated by $\Delta E^{\text{jet}} \sin\theta^{\text{jet}}$ where ΔE^{jet} is proportional to $\sqrt{E^{\text{jet}}}$. The opening angle $\Delta\phi(\text{jet}, \cancel{E}_T)$ between each jet and the missing transverse energy in the transverse plane provides a measure of the contribution of the jet to the missing transverse energy. The scaled missing transverse energy defined as

$$\cancel{E}_T^{\text{Sc}} = \frac{\cancel{E}_T}{\sqrt{\sum_{\text{jets}} (\Delta E^{\text{jet}} \sin\theta^{\text{jet}} \cos\Delta\phi(\text{jet}, \cancel{E}_T))^2}} \quad (1)$$

is required to be greater than 15. Finally, to suppress the




FIG. 1. Distribution of the missing transverse energy \cancel{E}_T after applying the initial transverse momentum cuts in the (a) ee , (c) $\mu\mu$, and (e) $e\mu$ channel. Figures (b), (d), and (f) show the \cancel{E}_T distributions after the final selection except for the \cancel{E}_T criterion for the ee , $\mu\mu$, and $e\mu$ channels, respectively. The arrows indicate the cut values. The QCD contribution is negligible in (c) and (d).

background from $t\bar{t}$ production, the scalar sum of the transverse energies of all jets with $E_T^{\text{jet}} > 20$ GeV and $|\eta| < 2.5$, H_T , is required to be less than 50 GeV. Figure 1(b) shows the \cancel{E}_T distribution after the final selection without applying the \cancel{E}_T criterion for the ee channel, and Fig. 2(a) shows the distribution of the minimal transverse mass after applying all selection criteria except the cut on the minimal transverse mass. Six events remain in the ee data sample after all of these cuts are applied.

For the $\mu\mu$ channel, to further reduce the Z/γ^* background, only events with an invariant di-muon mass between 20 and 80 GeV are retained. Since the momentum resolution is worsening for high p_T tracks, an additional constrained fit is performed to reject events compatible with Z boson production. The opening angle between the two muons in the transverse plane is required to be $\Delta\phi_{\mu\mu} < 2.4$. Finally, requiring $H_T < 100$ GeV removes the remaining background from $t\bar{t}$ events. Figure 1(d) shows the \cancel{E}_T distribution after the final selection without applying the \cancel{E}_T criterion for the $\mu\mu$ channel. Four events are observed in the $\mu\mu$ data sample after application of all selection criteria.

In the $e\mu$ channel, to suppress the WZ and ZZ backgrounds, events are rejected if a third lepton is found and the invariant mass of two leptons of the same flavor and

opposite charge is in the range from 61 to 121 GeV. To remove the background from multijet production and $Z/\gamma^* \rightarrow \tau\tau$ events, the minimal transverse mass $m_T^{\text{min}} = \min(m_T^e, m_T^\mu)$ must exceed 20 GeV. Remaining $Z/\gamma^* \rightarrow \tau\tau$ events, where large missing transverse energy is most likely introduced by mismeasured jets, are suppressed by removing events with $\cancel{E}_T^{\text{sc}} < 15$. Requiring $H_T < 50$ GeV rejects most of the $t\bar{t}$ events. To remove $W + \gamma$ events in which photons convert to electron-positron pairs, at least three hits in the silicon tracker are required for the electron track if the transverse mass determined from the muon and \cancel{E}_T is consistent with the W boson transverse mass. Figure 1(f) shows the \cancel{E}_T distribution after the final selection without applying the \cancel{E}_T criterion for the $e\mu$ channel, whereas Fig. 2(b) shows the distribution of the minimal transverse mass after applying all selection criteria except the cut on the minimal transverse mass. Fifteen events survive the final selection criteria in the $e\mu$ data sample.

The efficiency for WW signal events to pass the acceptance and kinematic criteria is determined using the PYTHIA 6.2 [8] event generator followed by a detailed GEANT-based [9] simulation of the D0 detector. All trigger and reconstruction efficiencies are derived from the data. For the ee channel, the overall detection efficiency is $(8.76 \pm 0.13)\%$. The overall efficiencies for the $\mu\mu$ and




FIG. 2. Distribution of the minimal transverse mass m_T^{\min} after applying all selection criteria except the cut on m_T^{\min} for (a) the ee and (b) the $e\mu$ channels. The arrows indicate the cut values.

$e\mu$ channels are $(6.22 \pm 0.15)\%$ and $(15.40 \pm 0.20)\%$, respectively. Using a NLO cross section of 13.5 pb [3] and branching fractions B of 0.1072 ± 0.0016 for $W \rightarrow e\nu$ and 0.1057 ± 0.0022 for $W \rightarrow \mu\nu$ [10], the expected number of events for the pair production of W bosons combined for all three channels is $16.6 \pm 0.1(\text{stat}) \pm 0.6(\text{syst}) \pm 1.1(\text{lum})$ events, where the statistical error is given by the statistics of the MC sample. The signal breakdown for the three channels is given by the first line of Table I.

Background contributions from Z/γ^* , $W + \text{jet}/\gamma$, $t\bar{t}$, WZ , and ZZ events are estimated using the PYTHIA event generator. In addition, $W + \text{jet}/\gamma$ contributions are verified using ALPGEN [11] and are cross checked with an estimation from the data using a matrix method that takes into account electron efficiencies and jet fake rates. All events are processed through the full detector simulation. The background due to multijet production, when a jet is misidentified as an electron, is determined from the data using a sample of like-sign di-lepton events with inverted lepton quality cuts (called QCD background in Figs. 1 and 2).

For the normalization of Z/γ^* and $W + \text{jet}/\gamma$ events, the next-to-next-to-leading order cross sections from

TABLE I. Number of signal and background events expected and number of events observed after all selections are applied for the three channels. Only statistical uncertainties are given.

Process	ee	$e\mu$	$\mu\mu$
WW signal	3.42 ± 0.05	11.10 ± 0.10	2.10 ± 0.05
$Z/\gamma^* \rightarrow ee$	0.20 ± 0.06
$Z/\gamma^* \rightarrow \mu\mu$...	0.28 ± 0.09	1.60 ± 0.40
$Z/\gamma^* \rightarrow \tau\tau$	<0.01	0.0 ± 0.1	<0.01
$t\bar{t}$	0.18 ± 0.02	0.34 ± 0.03	0.09 ± 0.01
WZ	0.33 ± 0.17	0.38 ± 0.02	0.15 ± 0.08
ZZ	0.19 ± 0.06	0.02 ± 0.02	0.10 ± 0.04
W + jet	0.97 ± 0.06	2.41 ± 0.06	0.01 ± 0.01
W + γ	0.43 ± 0.04	0.31 ± 0.04	...
Multijet	<0.05	0.07 ± 0.07	<0.05
Background sum	2.30 ± 0.21	3.81 ± 0.17	1.95 ± 0.41
Data	6	15	4

Ref. [12] are used. The cross section times branching ratio of Z/γ^* production in the invariant mass region $60 < m_{\ell\ell} < 130$ GeV is $\sigma \times B = 254$ pb. For inclusive W boson production with decays into a single lepton flavor state, this value is $\sigma \times B = 2717$ pb. The NLO WZ and ZZ production cross section values are taken from Ref. [4] with $\sigma \times B = 0.014$ pb for WZ and $\sigma \times B = 0.002$ pb for ZZ production with decay into a single lepton flavor state. The calculations of Ref. [13] are used for $t\bar{t}$ production with $\sigma \times B = 0.076$ pb with single flavor lepton decays of both W bosons. A summary of the background contributions together with signal expectations and events observed in the data after the final selection for the individual channels is shown in Table I. The total background sum is $8.1 \pm 0.6(\text{stat}) \pm 0.6(\text{syst}) \pm 0.5(\text{lum})$ events. The $e\mu$ channel has both the highest signal efficiency and the best signal-to-background ratio. There is good agreement between the number of events observed in the data and the sum of the expectations from WW production and the various backgrounds in all three channels.

Systematic uncertainties that affect the WW production cross section measurement are listed in Table II. In these estimates, parameters are varied within $\pm 1\sigma$ of the respective theoretical or experimental errors. Sources such as the trigger efficiency, electron and muon identification (ID) efficiencies, jet energy scale (JES), electron and muon momentum resolution, branching fraction $B(W \rightarrow \ell\nu)$, cross section calculation of Z/γ^* and $t\bar{t}$ events, and the determination of the $W + \text{jet}/\gamma$ background contribute to the systematic uncertainty. The PYTHIA Monte Carlo calculation tends to underestimate jet multiplicities, since a parton-shower approach is used for initial and final state radiation instead of the full matrix element. To compensate for this underestimation, events are reweighted in the MC to reproduce the jet multiplicities seen in the data. The systematic uncertainty for this approach is determined from a measurement of the WW production cross section with and without the reweighting. The total systematic uncertainties are given in Table II. The uncertainty on the luminosity measurement is 6.5%.

TABLE II. Systematic uncertainties for the ee , $e\mu$, and $\mu\mu$ channels.

Source	Change in the WW cross section (%)					
	ee	$e\mu$	$\mu\mu$	ee	$e\mu$	$\mu\mu$
Trigger, ID	+4.7	-4.6	+3.9	-3.8	+6.2	-5.8
JES	+3.2	-3.2	+1.6	-1.2	+7.2	-4.8
μ resolution	+4.7	-2.2	+10.0	-4.1
e resolution	+4.6	-2.9	+1.3	-1.1
$B(W \rightarrow \ell\nu)$	+4.4	-3.9	+5.3	-4.6	+4.3	-4.1
$\sigma(Z/\gamma^*, t\bar{t})$	+0.9	-0.7	+0.4	-0.4	+3.2	-3.2
W + jet/ γ	+4.0	-4.0	+3.0	-3.0
Reweighting	+4.3	-4.4	+1.5	-1.5
Total	+10.3	-9.5	+8.9	-7.3	+14.9	-10.1

The cross section for W boson pair production is estimated using a likelihood method [14,15] with Poisson statistics. The cross section for each channel $\sigma_{p\bar{p}\rightarrow W^+W^-}$ is given by

$$\sigma_{p\bar{p}\rightarrow W^+W^-} = \frac{N_{\text{obs}} - N_{\text{bg}}}{\int \mathcal{L} dt B \epsilon}, \quad (2)$$

where N_{obs} is the number of observed events, N_{bg} is the expected background, $\int \mathcal{L} dt$ is the integrated luminosity, B is the branching fraction for $W \rightarrow \ell \nu$, and ϵ is the efficiency for the signal. The likelihood for N_{obs} events in the data is given by

$$L(\sigma_{p\bar{p}\rightarrow W^+W^-}, N_{\text{obs}}, N_{\text{bg}}, \int \mathcal{L} dt, B, \epsilon) = \frac{N_{\text{obs}}^{N_{\text{obs}}}}{N_{\text{obs}}!} e^{-N}, \quad (3)$$

where N is the number of signal and background events:

$$N = \sigma_{p\bar{p}\rightarrow W^+W^-} B \int \mathcal{L} dt \epsilon + N_{\text{bg}}. \quad (4)$$

The cross section $\sigma_{p\bar{p}\rightarrow W^+W^-}$ is estimated by minimizing $-2 \ln L(\sigma_{p\bar{p}\rightarrow W^+W^-}, N_{\text{obs}}, N_{\text{bg}}, \int \mathcal{L} dt, B, \epsilon)$. To combine the channels, the individual likelihood functions are multiplied. As a final result, the combined cross section for WW production at a center-of-mass energy of $\sqrt{s} = 1.96$ TeV is

$$\sigma_{p\bar{p}\rightarrow W^+W^-} = 13.8_{-3.8}^{+4.3}(\text{stat})_{-0.9}^{+1.2}(\text{syst}) \pm 0.9(\text{lum}) \text{ pb}. \quad (5)$$

This value is in good agreement with the NLO calculation prediction of 12.0–13.5 pb at $\sqrt{s} = 1.96$ TeV [3].

The significance for the signal observation can be estimated using the likelihood ratio method [16]. The confidence levels for a background only hypothesis, CL_B , is obtained using the background expectation and the number of events observed as input. The signal significance is extracted from $1 - CL_B$. The confidence level for an upward fluctuation of the background to the observed number of events or higher in the absence of signal is 2.3×10^{-7} , which corresponds to 5.2 standard deviations for a Gaussian probability distribution.

To conclude, we have measured the W boson pair-production cross section in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV. We observe 25 events in the data, corresponding to integrated luminosities of 224–252 pb^{-1} depending on the final state, with a background expectation from non- WW processes of $8.1 \pm 0.6(\text{stat}) \pm 0.6(\text{syst}) \pm 0.5(\text{lum})$ events. The expectation for SM pair production of W bosons in our data sample is $16.6 \pm 0.1(\text{stat}) \pm 0.6(\text{syst}) \pm 1.1(\text{lum})$ events. We obtain a production cross section of $\sigma_{p\bar{p}\rightarrow W^+W^-} = 13.8_{-3.8}^{+4.3}(\text{stat})_{-0.9}^{+1.2}(\text{syst}) \pm 0.9(\text{lum})$ pb, consistent with the NLO prediction. The probability that the observed events are caused by a fluctuation of the background is 2.3×10^{-7} .

We thank the staffs at Fermilab and collaborating institutions, and acknowledge support from the DOE and NSF (USA), CEA and CNRS/IN2P3 (France), Ministry of Education and Science, Agency for Atomic Energy and RF President Grants Program (Russia), CAPES, CNPq, FAPERJ, FAPESP, and FUNDUNESP (Brazil), DAE and DST (India), Colciencias (Colombia), CONACyT (Mexico), KRF (Korea), CONICET and UBACyT (Argentina), FOM (The Netherlands), PPARC (United Kingdom), Ministry of Education (Czech Republic), Natural Sciences and Engineering Research Council and WestGrid Project (Canada), BMBF and DFG (Germany), A.P. Sloan Foundation, Research Corporation, Texas Advanced Research Program, and the Alexander von Humboldt Foundation.

*Visitor from University of Zurich, Zurich, Switzerland.

†Visitor from Institute of Nuclear Physics, Krakow, Poland.

- [1] D0 Collaboration, B. Abbott *et al.*, Phys. Rev. D **60**, 072002 (1999); K. Hagiwara, J. Woodside, and D. Zeppenfeld, Phys. Rev. D **41**, 2113 (1990).
- [2] E. Eichten, I. Hinchliffe, K. Lane, and C. Quigg, Rev. Mod. Phys. **56**, 579 (1984); **58**, 1065 (1986).
- [3] J. Ohnemus, Phys. Rev. D **44**, 1403 (1991); **50**, 1931 (1994); J.M. Campbell and R.K. Ellis, *ibid.* **60**, 113006 (1999).
- [4] CDF Collaboration, F. Abe *et al.*, Phys. Rev. Lett. **78**, 4536 (1997).
- [5] R. Strohmer, Int. J. Mod. Phys. A **18**, 5127 (2003).
- [6] D0 Collaboration, V. Abazov *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A (to be published); T. LeCompte and H.T. Diehl, Annu. Rev. Nucl. Part. Sci. **50**, 71 (2000).
- [7] D0 Collaboration, S. Abachi *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **338**, 185 (1994).
- [8] T. Sjöstrand *et al.*, Comput. Phys. Commun. **135**, 238 (2001).
- [9] R. Brun and F. Carminati, CERN Program Library Long Writeup, 1993 (unpublished), W5013.
- [10] Particle Data Group, S. Eidelman *et al.*, Phys. Lett. B **592**, 1 (2004).
- [11] M.L. Mangano *et al.*, J. High Energy Phys. 07 (2003) 001.
- [12] R. Hamberg, W.L. van Neerven, and T. Matsuura, Nucl. Phys. **B359**, 343 (1991); **B644**, 403(E) (2002).
- [13] N. Kidonakis and R. Vogt, Phys. Rev. D **68**, 114014 (2003).
- [14] G.J. Feldman and R.D. Cousins, Phys. Rev. D **57**, 3873 (1998).
- [15] B.P. Roe and M.B. Woodroffe, Phys. Rev. D **60**, 053009 (1999).
- [16] T. Junk, Nucl. Instrum. Methods Phys. Res., Sect. A **434**, 435 (1999).

Erratum: Measurement of the WW Production Cross Section in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV [Phys. Rev. Lett. 94, 151801 (2005)]

V. M. Abazov, B. Abbott, M. Abolins, B. S. Acharya, M. Adams, T. Adams, M. Agelou, J.-L. Agram, S. H. Ahn, M. Ahsan, G. D. Alexeev, G. Alkhazov, A. Alton, G. Alverson, G. A. Alves, M. Anastasoae, S. Anderson, B. Andrieu, Y. Arnaud, A. Askew, B. Åsman, O. Atramentov, C. Autermann, C. Avila, F. Badaud, A. Baden, B. Baldin, P. W. Balm, S. Banerjee, E. Barberis, P. Bargassa, P. Baringer, C. Barnes, J. Barreto, J. F. Bartlett, U. Bassler, D. Bauer, A. Bean, S. Beauceron, M. Begel, A. Bellavance, S. B. Beri, G. Bernardi, R. Bernhard, I. Bertram, M. Beşançon, R. Beuselinck, V. A. Bezzubov, P. C. Bhat, V. Bhatnagar, M. Binder, K. M. Black, I. Blackler, G. Blazey, F. Blekman, S. Blessing, D. Bloch, U. Blumenschein, A. Boehnlein, O. Boeriu, T. A. Bolton, F. Borchering, G. Borissov, K. Bos, T. Bose, A. Brandt, R. Brock, G. Brooijmans, A. Bross, N. J. Buchanan, D. Buchholz, M. Buehler, V. Buescher, S. Burdin, T. H. Burnett, E. Busato, J. M. Butler, J. Bystricky, W. Carvalho, B. C. K. Casey, N. M. Cason, H. Castilla-Valdez, S. Chakrabarti, D. Chakraborty, K. M. Chan, A. Chandra, D. Chapin, F. Charles, E. Cheu, L. Chevalier, D. K. Cho, S. Choi, T. Christiansen, L. Christofek, D. Claes, B. Clément, C. Clément, Y. Coadou, M. Cooke, W. E. Cooper, D. Coppage, M. Corcoran, J. Coss, A. Cothenet, M.-C. Cousinou, S. Crépe- Renaudin, M. Cristetiu, M. A. C. Cummings, D. Cutts, H. da Motta, B. Davies, G. Davies, G. A. Davis, K. De, P. de Jong, S. J. de Jong, E. De La Cruz-Burelo, C. De Oliveira Martins, S. Dean, F. Déliot, P. A. Delsart, M. Demarteau, R. Demina, P. Demine, D. Denisov, S. P. Denisov, S. Desai, H. T. Diehl, M. Diesburg, M. Doidge, H. Dong, S. Doulas, L. Dufлот, S. R. Dugad, A. Duperrin, J. Dyer, A. Dyshkant, M. Eads, D. Edmunds, T. Edwards, J. Ellison, J. Elmsheuser, J. T. Eltzroth, V. D. Elvira, S. Eno, P. Ermolov, O. V. Eroshin, J. Estrada, D. Evans, H. Evans, A. Evdokimov, V. N. Evdokimov, J. Fast, S. N. Fatakia, L. Feligioni, T. Ferbel, F. Fiedler, F. Filthaut, W. Fisher, H. E. Fisk, M. Fortner, H. Fox, W. Freeman, S. Fu, S. Fuess, T. Gadfort, C. F. Galea, E. Gallas, E. Galyaev, C. Garcia, A. Garcia-Bellido, J. Gardner, V. Gavrilov, P. Gay, D. Gelé, R. Gelhaus, K. Genser, C. E. Gerber, Y. Gershtein, G. Ginther, T. Golling, B. Gómez, K. Gounder, A. Goussiou, P. D. Grannis, S. Greder, H. Greenlee, Z. D. Greenwood, E. M. Gregores, Ph. Gris, J.-F. Grivaz, L. Groer, S. Grünendahl, M. W. Grünewald, S. N. Gurzhiev, G. Gutierrez, P. Gutierrez, A. Haas, N. J. Hadley, S. Hagopian, I. Hall, R. E. Hall, C. Han, L. Han, K. Hanagaki, K. Harder, R. Harrington, J. M. Hauptman, R. Hauser, J. Hays, T. Hebbeker, D. Hedin, J. M. Heinmiller, A. P. Heinson, U. Heintz, C. Hensel, G. Hesketh, M. D. Hildreth, R. Hirosky, J. D. Hobbs, B. Hoeneisen, M. Hohlfeld, S. J. Hong, R. Hooper, P. Houben, Y. Hu, J. Huang, I. Iashvili, R. Illingworth, A. S. Ito, S. Jabeen, M. Jaffré, S. Jain, V. Jain, K. Jakobs, A. Jenkins, R. Jesik, K. Johns, M. Johnson, A. Jonckheere, P. Jonsson, H. Jöstlein, A. Juste, M. M. Kado, D. Käfer, W. Kahl, S. Kahn, E. Kajfasz, A. M. Kalinin, J. Kalk, D. Karmanov, J. Kasper, D. Kau, R. Kehoe, S. Kermiche, S. Kesisoglou, A. Khanov, A. Kharchilava, Y. M. Kharzhev, K. H. Kim, B. Klima, M. Klute, J. M. Kohli, M. Kopal, V. M. Korablev, J. Kotcher, B. Kothari, A. Koubarovsky, A. V. Kozelov, J. Kozminski, S. Krzywdzinski, S. Kuleshov, Y. Kulik, S. Kunori, A. Kupco, T. Kurča, S. Lager, N. Lahrichi, G. Landsberg, J. Lazoflores, A.-C. Le Bihan, P. Lebrun, S. W. Lee, W. M. Lee, A. Leflat, F. Lehner, C. Leonidopoulos, P. Lewis, J. Li, Q. Z. Li, J. G. R. Lima, D. Lincoln, S. L. Linn, J. Linnemann, V. V. Lipaev, R. Lipton, L. Lobo, A. Lobodenko, M. Lokajicek, A. Lounis, H. J. Lubatti, L. Lueking, M. Lynker, A. L. Lyon, A. K. A. Maciel, R. J. Madaras, P. Mättig, A. Magerkurth, A.-M. Magnan, N. Makovec, P. K. Mal, S. Malik, V. L. Malyshev, H. S. Mao, Y. Maravin, M. Martens, S. E. K. Mattingly, A. A. Mayorov, R. McCarthy, R. McCroskey, D. Meder, H. L. Melanson, A. Melnitchouk, M. Merkin, K. W. Merritt, A. Meyer, H. Miettinen, D. Mihalcea, J. Mitrevski, N. Mokhov, J. Molina, N. K. Mondal, H. E. Montgomery, R. W. Moore, G. S. Muanza, M. Mulders, Y. D. Mutaf, E. Nagy, M. Narain, N. A. Naumann, H. A. Neal, J. P. Negret, S. Nelson, P. Neustroev, C. Noeding, A. Nomerotski, S. F. Novaes, T. Nunnemann, E. Nurse, V. O'Dell, D. C. O'Neil, V. Oguri, N. Oliveira, N. Oshima, G. J. Otero y Garzón, P. Padley, N. Parashar, J. Park, S. K. Park, J. Parsons, R. Partridge, N. Parua, A. Patwa, P. M. Perea, E. Perez, O. Peters, P. Pétróff, M. Petteni, L. Phaf, R. Piegaiia, P. L. M. Podesta-Lerma, V. M. Podstavkov, Y. Pogorelov, B. G. Pope, W. L. Prado da Silva, H. B. Prosper, S. Protopopescu, M. B. Przybycien, J. Qian, A. Quadt, B. Quinn, K. J. Rani, P. A. Rapidis, P. N. Ratoff, N. W. Reay, S. Reucroft, M. Rijssenbeek, I. Ripp-Baudot, F. Rizatdinova, C. Royon, P. Rubinov, R. Ruchti, G. Sajot, A. Sánchez-Hernández, M. P. Sanders, A. Santoro, G. Savage, L. Sawyer, T. Scanlon, R. D. Schamberger, H. Schellman, P. Schieferdecker, C. Schmitt, A. A. Schukin, A. Schwartzman, R. Schwienhorst, S. Sengupta, H. Severini, E. Shabalina, M. Shamim, V. Shary, W. D. Shephard, D. Shpakov, R. A. Sidwell, V. Simak, V. Sirotenko, P. Skubic, P. Slattery, R. P. Smith, K. Smolek, G. R. Snow, J. Snow, S. Snyder, S. Söldner-Rembold, X. Song, Y. Song, L. Sonnenschein, A. Sopczak, M. Sosebee, K. Soustruznik, M. Souza, B. Spurlock, N. R. Stanton, J. Stark, J. Steele, G. Steinbrück,

K. Stevenson, V. Stolin, A. Stone, D. A. Stoyanova, J. Strandberg, M. A. Strang, M. Strauss, R. Ströhmer, M. Strovink, L. Stutte, S. Sumowidagdo, A. Sznajder, M. Talby, P. Tamburello, W. Taylor, P. Telford, J. Temple, S. Tentindo-Repond, E. Thomas, B. Thooris, M. Tomoto, T. Toole, J. Torborg, S. Towers, T. Trefzger, S. Trincaz-Duvoid, B. Tuchming, C. Tully, A. S. Turcot, P. M. Tuts, L. Uvarov, S. Uvarov, S. Uzunyan, B. Vachon, R. Van Kooten, W. M. van Leeuwen, N. Varelas, E. W. Varnes, I. A. Vasilyev, M. Vaupel, P. Verdier, L. S. Vertogradov, M. Verzocchi, F. Villeneuve-Seguiet, J.-R. Vlimant, E. Von Toerne, M. Vreeswijk, T. Vu Anh, H. D. Wahl, R. Walker, L. Wang, Z.-M. Wang, J. Warchol, M. Warsinsky, G. Watts, M. Wayne, M. Weber, H. Weerts, M. Wegner, N. Wermes, A. White, V. White, D. Whiteson, D. Wicke, D. A. Wijngaarden, G. W. Wilson, S. J. Wimpenny, J. Wittlin, M. Wobisch, J. Womersley, D. R. Wood, T. R. Wyatt, Q. Xu, N. Xuan, R. Yamada, M. Yan, T. Yasuda, Y. A. Yatsunenko, Y. Yen, K. Yip, S. W. Youn, J. Yu, A. Yurkewicz, A. Zabi, A. Zatserklyaniy, M. Zdrzil, C. Zeitnitz, D. Zhang, X. Zhang, T. Zhao, Z. Zhao, B. Zhou, J. Zhu, M. Zielinski, D. Zieminska, A. Zieminski, R. Zitoun, V. Zutshi, E. G. Zverev, and A. Zylberstejn

(D0 Collaboration)

(Received 23 January 2008; published 3 April 2008)

DOI: [10.1103/PhysRevLett.100.139901](https://doi.org/10.1103/PhysRevLett.100.139901)

PACS numbers: 13.38.Be, 13.85.Qk, 14.70.Fm, 99.10.Cd

The measurement of the cross section $\sigma(p\bar{p} \rightarrow W^+W^-)$ published in 2005 [1] requires a correction for an increase in the reported integrated luminosity. The instantaneous luminosity at D0 is measured by counting the number of inelastic collisions that produce charged particles within the acceptance of the luminosity monitor [2]. The determination of the luminosity has recently been improved through studies of the multiplicities observed in the luminosity monitor [3]. These studies indicated that the fraction of observable inelastic collisions was overestimated in our previous analysis [4]. For this analysis the estimated integrated luminosities were 252 pb^{-1} , 235 pb^{-1} , and 224 pb^{-1} for the e^+e^- , $e^\pm\mu^\pm$, and $\mu^+\mu^-$ channels, respectively. The improved estimates are 294 pb^{-1} , 274 pb^{-1} , and 261 pb^{-1} . The luminosity uncertainty decreased from 6.5% to 6.1%. The corrected value for the cross section is $\sigma(p\bar{p} \rightarrow W^+W^-) = 11.8_{-3.2}^{+3.7}(\text{stat})_{-0.8}^{+1.0}(\text{syst}) \pm 0.7(\text{lum}) \text{ pb}$. The new value is in good agreement with the NLO standard model predictions of 12.0–13.5 pb [5].

- [1] V. M. Abazov *et al.* (D0 Collaboration), *Phys. Rev. Lett.* **94**, 151801 (2005).
- [2] V. M. Abazov *et al.* (D0 Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **565**, 463 (2006).
- [3] T. Andeen *et al.*, Fermilab Report No. FERMILAB-TM-2365, 2007.
- [4] T. Edwards *et al.*, Fermilab Report No. FERMILAB-TM-2278-E, 2004.
- [5] J. Ohnemus, *Phys. Rev. D* **44**, 1403 (1991); **50**, 1931 (1994); J. M. Campbell and R. K. Ellis, *ibid.* **60**, 113006 (1999).