



KM3NeT

Opens a new window on our universe

arXiv:1601.07459v1 [astro-ph.IM] 27 Jan 2016

KM3NeT 2.0

Letter of Intent for ARCA and ORCA

– Astroparticle & Oscillation Research with Cosmics in the Abyss –

27th January 2016

Contact: spokesperson@km3net.de





The KM3NeT optical module. The KM3NeT infrastructure comprises several thousand identical optical modules arranged in three-dimensional spatial arrays located in the deep waters of the Mediterranean Sea. The spacing between the optical modules is different for the ARCA and ORCA detectors to optimally detect neutrinos with the targeted energies. Each optical module consists of a glass sphere with a diameter of 42 cm, housing 31 photo-sensors (yellowish disks). The glass sphere can withstand the pressure of the water and is transparent to the faint light that must be detected to see neutrinos.

The KM3NeT Collaboration:

S. Adrián-Martínez³⁹, M. Ageron², F. Aharonian³³, S. Aiello¹⁸, A. Albert⁴⁰, F. Ameli²², E. Anassontzis³¹, M. Andre³⁸, G. Androulakis²⁹, M. Anghinolfi¹⁹, G. Anton⁷, M. Ardid³⁹, T. Avgitas³, G. Barbarino^{20,48}, E. Barbarito¹⁶, B. Baret³, J. Barrios-Martí¹¹, B. Belhorma²⁷, A. Belias²⁹, E. Berbee⁶, A. van den Berg²⁵, V. Bertin², S. Beurthey², V. van Beveren⁶, N. Beverini^{36,21}, S. Biagi¹⁴, A. Biagioni²², M. Billault², R. Bormuth^{6,26}, B. Bouhadef²¹, G. Bourlis¹⁰, S. Bourret³, C. Boutonnet³, M. Bouwhuis⁶, C. Bozza⁴⁹, R. Bruijn⁴³, J. Brunner², E. Buis³⁴, J. Busto², G. Cacapardo¹⁴, L. Caillat², M. Calamai²¹, D. Calvo¹¹, A. Capone^{37,22}, L. Caramete²⁴, S. Cecchini¹⁷, S. Celli^{37,22,9}, C. Champion³, R. Cherkaoui El Moursli⁴², S. Cherubini^{14,45}, T. Chiarusi¹⁷, M. Circella¹⁶, L. Classen⁷, R. Cocimano¹⁴, J. A. B. Coelho³, A. Coleiro³, S. Colonges³, R. Coniglione¹⁴, M. Cordelli¹⁵, A. Cosquer², P. Coyle², A. Creusot³, G. Cuttone¹⁴, A. D'Amico⁶, G. De Bonis²², G. De Rosa^{20,48}, C. De Sio⁴⁹, F. Di Capua²⁰, I. Di Palma^{37,22}, A. F. Díaz García³⁵, C. Distefano¹⁴, C. Donzaud³, D. Dornic², Q. Dorosti-Hasankiadeh²⁵, E. Drakopoulou²⁹, D. Drouhin⁴⁰, L. Drury³³, M. Durocher^{14,9}, T. Eberl⁷, S. Eichie^{7,8}, D. van Eijk⁶, I. El Bojaddaini⁴¹, N. El Khayati⁴², D. Elsaesser⁵¹, A. Enzenhöfer², F. Fassi⁴², P. Favali²³, P. Fermani²², G. Ferrara^{14,45}, C. Filippidis²⁹, L. A. Fusco^{17,44}, T. Gal⁷, S. Galatà³, F. Garufi^{20,48}, P. Gay^{3,13}, M. Gebyehu⁶, V. Giordano¹⁸, N. Gizani¹⁰, R. Gracia³, K. Graf⁷, T. Grégoire³, G. Grella⁴⁹, R. Habel¹⁵, S. Hallmann⁷, H. van Haren³⁰, S. Harissopulos²⁹, T. Heid⁷, A. Heijboer⁶, E. Heine⁶, S. Henry², J. J. Hernández-Rey¹¹, M. Hevinga²⁵, J. Hofestädt⁷, C. M. F. Hugon¹⁹, G. Illuminati¹¹, C. W. James⁷, P. Jansweijer⁶, M. Jongen⁶, M. de Jong⁶, M. Kadler⁵¹, O. Kalekin⁷, A. Kappes⁷, U. F. Katz⁷, P. Keller², G. Kieft⁶, D. Kießling⁷, E. N. Koffeman⁶, P. Kooijman^{43,52}, A. Kouchner³, V. Kulikovskiy¹⁴, R. Lahmann⁷, P. Lamare², E. Leonora¹⁸, M. Lindsey Clark³, A. Liolios⁴, C. D. Llorens Alvarez³⁹, D. Lo Presti¹⁸, H. Löhner²⁵, A. Lonardo²², M. Lotze¹¹, S. Loucatos³, E. Maccioni^{36,21}, K. Mannheim⁵¹, A. Margiotta^{17,44}, A. Marinelli^{36,21}, O. Mariş²⁴, C. Markou²⁹, J. A. Martínez-Mora³⁹, A. Martini¹⁵, R. Mele^{20,48}, K. W. Melis⁶, T. Michael⁶, P. Migliozi²⁰, E. Migneco¹⁴, P. Mijakowski²⁸, A. Miraglia¹⁴, C. M. Mollo²⁰, M. Mongelli¹⁶, M. Morganti^{21,1}, A. Moussa⁴¹, P. Musico¹⁹, M. Musumeci¹⁴, S. Navas³⁵, C. A. Nicolau²², I. Olcina¹¹, C. Olivetto³, A. Orlando¹⁴, A. Papaikonomou¹⁰, R. Papaleo¹⁴, G. E. Păvălaş²⁴, H. Peek⁶, C. Pellegrino^{17,44}, C. Perrina^{37,22}, M. Pfitzner⁶, P. Piattelli¹⁴, K. Pikounis²⁹, G. E. Poma^{14,45}, V. Popa²⁴, T. Pradier¹², F. Pratolongo¹⁹, G. Pühlhofer⁵, S. Pulvirenti¹⁴, L. Quinn², C. Racca⁴⁰, F. Raffaelli²¹, N. Randazzo¹⁸, P. Rapidis²⁹, P. Razis⁴⁶, D. Real¹¹, L. Resvanis³¹, J. Reubelt⁷, G. Riccobene¹⁴, C. Rossi¹⁹, A. Rovelli¹⁴, M. Saldaña³⁹, I. Salvadori², D. F. E. Samtleben^{6,26}, A. Sánchez García¹¹, A. Sánchez Losa¹⁶, M. Sanguineti¹⁹, A. Santangelo⁵, D. Santonocito¹⁴, P. Sapienza¹⁴, F. Schimmel⁶, J. Schmelling⁶, V. Sciacca¹⁴, M. Sedita¹⁴, T. Seitz⁷, I. Sgura¹⁶, F. Simeone²², I. Siotis²⁹, V. Sipala¹⁸, B. Spisso²⁰, M. Spurio^{17,44}, G. Stavropoulos²⁹, J. Steijger⁶, S. M. Stellacci⁴⁹, D. Stransky⁷, M. Taiuti^{19,47}, Y. Tayalati^{41,42}, D. Tézier², S. Theraube², L. Thompson⁵⁰, P. Timmer⁶, C. Tönnis¹¹, L. Trasatti¹⁵, A. Trovato¹⁴, A. Tsirigotis¹⁰, S. Tzamarias¹⁰, E. Tzamariudaki²⁹, B. Vallage³, V. Van Elewyck³, J. Vermeulen⁶, P. Vicini²², S. Viola¹⁴, D. Vivolo^{20,48}, M. Volkert⁷, G. Voulgaris³¹, L. Wiggers⁶, J. Wilms⁸, E. de Wolf^{6,43}, K. Zachariadou³², J. D. Zornoza¹¹, J. Zúñiga¹¹

Additional Supporters:

D. Allard^a, A. Esmaili^{b,i}, G. Fogli^{f,m}, O. Mena^e, K. Murase^c, S. Palomares Ruiz^e, E. Parizot^a, S. T. Petcov^{h,k}, S. Razzaqueⁿ, D. Semikoz^a, A. Yu. Smirnov^{g,j}, F.L. Villante^{i,o}, F. Vissani^{d,i}, R. Zukanovich Funchal^l

Affiliations of KM3NeT members:

- ¹ Accademia Navale di Livorno, Viale Italia 72, Livorno, 57100 Italy
- ² Aix-Marseille Université, CNRS/IN2P3, CPPM UMR 7346, 13288, Marseille, France
- ³ APC, Université Paris Diderot, CNRS/IN2P3, CEA/IRFU, Observatoire de Paris, Sorbonne Paris Cité, 75205 Paris, France
- ⁴ Aristotle University Thessaloniki, University Campus, Thessaloniki, 54124 Greece
- ⁵ Eberhard Karls Universität Tübingen, Institut für Astronomie und Astrophysik, Sand 1, 72076 Tübingen, Germany
- ⁶ FOM, Nikhef, PO Box 41882, Amsterdam, 1098 DB Netherlands
- ⁷ Friedrich-Alexander-Universität Erlangen-Nürnberg, Erlangen Centre for Astroparticle Physics, Erwin-Rommel-Straße 1, 91058 Erlangen, Germany
- ⁸ Friedrich-Alexander-Universität Erlangen-Nürnberg, Remeis Sternwarte, Sternwartstraße 7, 96049 Bamberg, Germany
- ⁹ Gran Sasso Science Institute, GSSI, Viale Francesco Crispi 7, L'Aquila, 67100 Italy
- ¹⁰ Hellenic Open University, School of Science / Technology, Natural Sciences, Sahtouri St. / Ag. Andreou St. 16, Patra, 26222 Greece
- ¹¹ IFIC - Instituto de Física Corpuscular (CSIC - Universitat de València), c/Catedrático José Beltrán, 2, 46980 Paterna, Valencia, Spain
- ¹² IN2P3, IPHC, 23 rue du Loess, Strasbourg, 67037 France
- ¹³ IN2P3, LPC, Campus des Cézeaux 24, avenue des Landais BP 80026, Aubière Cedex, 63171 France
- ¹⁴ INFN, Laboratori Nazionali del Sud, Via S. Sofia 62, Catania, 95123 Italy
- ¹⁵ INFN, LNF, Via Enrico Fermi , 40, Frascati, 00044 Italy
- ¹⁶ INFN, Sezione di Bari, Via Amendola 173, Bari, 70126 Italy
- ¹⁷ INFN, Sezione di Bologna, v.le C. Berti-Pichat, 6/2, Bologna, 40127 Italy
- ¹⁸ INFN, Sezione di Catania, Via Santa Sofia 64, Catania, 95123 Italy
- ¹⁹ INFN, Sezione di Genova, Via Dodecaneso 33, Genova, 16146 Italy
- ²⁰ INFN, Sezione di Napoli, Complesso Universitario di Monte S. Angelo, Via Cintia ed. G, Napoli, 80126 Italy
- ²¹ INFN, Sezione di Pisa, Largo Bruno Pontecorvo 3, Pisa, 56127 Italy
- ²² INFN, Sezione di Roma, Piazzale Aldo Moro 2, Roma, 00185 Italy
- ²³ INGV, Via di Vigna Murata, 605, Rome, 00143 Italy
- ²⁴ ISS, 242, Vacaresti, Bucharest, 40061 Romania
- ²⁵ KVI-CART University of Groningen, Groningen, The Netherlands
- ²⁶ Leiden University, Leiden Institute of Physics, PO Box 9504, Leiden, 2300 RA Netherlands
- ²⁷ National Center for Energy Sciences and Nuclear Techniques, B.P. 1382, R.P. 10001 Rabat, Morocco
- ²⁸ National Centre for Nuclear Research, 00-681 Warsaw, Poland
- ²⁹ NCSR Demokritos, Institute of Nuclear and Particle Physics, Ag. Paraskevi Attikis, Athens, 15310 Greece
- ³⁰ NIOZ, PO Box 59, Den Burg, Texel, 1790 AB Netherlands
- ³¹ Physics Department, N. and K. University of Athens, Athens, Greece
- ³² Technological Education Institute of Pireaus, Thivon and P. Ralli Str. 250, Egaleo - Athens, 12244 Greece
- ³³ The Dublin Institute for Advanced Studies, 10 Burlington Road, Dublin, 4 Ireland
- ³⁴ TNO, Technical Sciences, PO Box 155, Delft, 2600 AD Netherlands
- ³⁵ Universidad de Granada & C.A.F.P.E, Av. del Hospicio s/n, 18071 Granada, Spain
- ³⁶ Università di Pisa, Dipartimento di Fisica, Largo Bruno Pontecorvo 3, Pisa, 56127 Italy
- ³⁷ Università La Sapienza, Dipartimento di Fisica, Piazzale Aldo Moro 2, Roma, 00185 Italy
- ³⁸ Universitat Politècnica de Catalunya, Laboratori d'Aplicacions Bioacústiques, Centre Tecnològic de Vilanova i la Geltrú, Avda. Rambla Exposició, s/n, Vilanova i la Geltrú, 08800 Spain
- ³⁹ Universitat Politècnica de València, Instituto de Investigación para la Gestión Integrada de las Zonas Costeras, C/ Paranímf, 1, Gandia, 46730 Spain
- ⁴⁰ Université de Strasbourg, Université de Haute Alsace, GRPHE, 34, Rue du Grillenbreit, Colmar, 68008 France
- ⁴¹ University Mohammed I, Faculty of Sciences, BV Mohammed VI, B.P. 717, R.P. 60000 Oujda, Morocco

- ⁴² University Mohammed V in Rabat, Faculty of Sciences, 4 av. Ibn Battouta, B.P. 1014, R.P. 10000 Rabat, Morocco
- ⁴³ University of Amsterdam, Institute of Physics/IHEF, PO Box 94216, Amsterdam, 1090 GE Netherlands
- ⁴⁴ University of Bologna, Dipartimento di Fisica e Astronomia, v.le C. Berti-Pichat, 6/2, Bologna, 40127 Italy
- ⁴⁵ University of Catania, Dipartimento di Fisica ed Astronomia di Catania, Via Santa Sofia 64, Catania, 95123 Italy
- ⁴⁶ University of Cyprus, Physics, Kallipoleos 75, Nicosia, 1678 Cyprus
- ⁴⁷ University of Genova, Via Dodecaneso 33, Genova, 16146 Italy
- ⁴⁸ University of Napoli, Dip. Scienze fisiche, Complesso Universitario di Monte S. Angelo, Via Cintia ed. G, Napoli, 80126 Italy
- ⁴⁹ University of Salerno, Department of Physics, Via Giovanni Paolo II 132, Fisciano, 84084 Italy
- ⁵⁰ University of Sheffield, Department of Physics and Astronomy, Hounsfield Road, Sheffield, S3 7RH United Kingdom
- ⁵¹ University Würzburg, Emil-Fischer-Straße 31, 97074 Würzburg, Germany
- ⁵² Utrecht University, Department of Physics and Astronomy, PO Box 80000, Utrecht, 3508 TA Netherlands

Affiliations of additional supporters:

- ^a APC, Université Paris Diderot, CNRS/IN2P3, CEA/IRFU, Observatoire de Paris, Sorbonne Paris Cité, 75205 Paris, France
- ^b Departamento de Física, Pontificia Universidade Católica do Rio de Janeiro, C.P. 38071, 22452-970, Rio de Janeiro, Brazil
- ^c Department of Physics; Department of Astronomy & Astrophysics, The Pennsylvania State University, University Park, Pennsylvania 16802, USA
- ^d Gran Sasso Science Institute, GSSI, Viale Francesco Crispi 7, L'Aquila, 67100 Italy
- ^e IFIC - Instituto de Física Corpuscular (CSIC - Universitat de València), c/Catedrático José Beltrán, 2, 46980 Paterna, Valencia, Spain
- ^f INFN, Sezione di Bari, Via Amendola 173, 70126 Bari, Italy
- ^g International Centre for Theoretical Physics, Strada Costiera 11, I-34100 Trieste, Italy
- ^h Kavli IPMU (WPI), University of Tokyo, 5-1-5 Kashiwanoha, 277-8583 Kashiwa, Japan
- ⁱ Laboratori Nazionali del Gran Sasso, INFN, Assergi (AQ), Italy
- ^j Max-Planck-Institut für Kernphysik, Saupfercheckweg 1, 69117 Heidelberg, Germany
- ^k SISSA/INFN, Via Bonomea 265, 34136 Trieste, Italy
- ^l Universidade de São Paulo, Instituto de Física, Departamento de Física Matemática, C.P.66318, 05315-970 São Paulo, Brazil
- ^m University of Bari, Dipartimento di Fisica, Via Amendola 173, 70126 Bari, Italy
- ⁿ University of Johannesburg, Department of Physics, PO Box 524, Auckland Park 2006, South Africa
- ^o University of L'Aquila, Dip. Scienze Fisiche e Chimiche, via Vetoio, 67100 L'Aquila, Italy

Contents

Executive Summary	1
1 Detector Design and Technology	3
1.1 KM3NeT/ARCA: deep sea and onshore infrastructures	3
1.2 KM3NeT/ORCA: deep sea and onshore infrastructures	4
1.3 Detection string	5
1.4 Digital optical module	6
1.5 Fibre-optic data transmission system	9
1.6 Data acquisition	9
1.6.1 Event trigger	10
1.6.2 Performance	10
2 Astroparticle Research with Cosmics in the Abyss (ARCA)	13
2.1 Introduction	13
2.1.1 Cosmic neutrinos	13
2.1.2 Assumptions	14
2.2 Simulations	16
2.2.1 Event generation	16
2.2.2 Detector response	17
2.2.3 Further improvements	19
2.2.4 Event reconstruction	20
2.2.5 Cascade reconstruction	21
2.2.6 Prospects for improved reconstruction	23
2.2.7 Background suppression	25
2.3 Sensitivity studies	26
2.3.1 Isotropic diffuse neutrino flux	27
2.3.2 Diffuse neutrino flux from the Galactic plane	34
2.3.3 Point-like neutrino sources	35
2.3.4 Further physics opportunities	41
2.4 Investigation of systematic effects	43
2.4.1 Optics: water properties and DOM response	44
2.4.2 Detector calibration and alignment	45
2.4.3 Ageing effects	45
2.5 Detector geometry studies	45
3 Oscillation Research with Cosmics in the Abyss (ORCA)	47
3.1 Introduction	47
3.2 Simulations	54
3.2.1 Benchmark detector	54
3.2.2 Event generation and characterization	54
3.2.3 Implementation of different vertical spacings	59
3.2.4 Triggering	59
3.3 Muon neutrino studies	60
3.3.1 Muon direction reconstruction	60
3.3.2 Neutrino energy estimate	63
3.3.3 Performance	67
3.4 Electron neutrino studies	68
3.4.1 Phenomenology of shower events	68
3.4.2 Shower reconstruction algorithm	72
3.4.3 Event selection	76
3.4.4 Reconstruction performance	77
3.4.5 Performance for different vertical spacings	84

3.4.6	Effect of variation in water/PMT properties and noise level on reconstruction performance	85
3.5	Flavour identification and muon rejection	88
3.5.1	Methodology	88
3.5.2	Event preselection	88
3.5.3	Classification input	89
3.5.4	Classification performance	90
3.6	Sensitivity studies for the neutrino mass hierarchy	92
3.6.1	Global fit	92
3.6.2	Spacing studies	97
3.6.3	Measurement of ΔM^2 and θ_{23}	99
3.6.4	Systematic uncertainties	101
3.7	Outlook	101
4	Organisation	103
5	Data policy	105
6	Cost and Time Schedule	107
	References	109

Executive Summary

The main objectives of the KM3NeT¹ Collaboration are *i)* the discovery and subsequent observation of high-energy neutrino sources in the Universe and *ii)* the determination of the mass hierarchy of neutrinos. These objectives are strongly motivated by two recent important discoveries, namely: *1)* The high-energy astrophysical neutrino signal reported by IceCube and *2)* the sizable contribution of electron neutrinos to the third neutrino mass eigenstate as reported by Daya Bay, Reno and others. To meet these objectives, the KM3NeT Collaboration plans to build a new Research Infrastructure consisting of a network of deep-sea neutrino telescopes in the Mediterranean Sea. A phased and distributed implementation is pursued which maximises the access to regional funds, the availability of human resources and the synergetic opportunities for the earth and sea sciences community. Three suitable deep-sea sites are identified, namely off-shore Toulon (France), Capo Passero (Italy) and Pylos (Greece).

The infrastructure will consist of three so-called building blocks. A building block comprises 115 strings, each string comprises 18 optical modules and each optical module comprises 31 photo-multiplier tubes (PMTs). Each building block thus constitutes a 3-dimensional array of photo sensors that can be used to detect the Cherenkov light produced by relativistic particles emerging from neutrino interactions. Two building blocks will be configured to fully explore the IceCube signal with different methodology, improved resolution and complementary field of view, including the Galactic plane. Collectively, these building blocks are referred to as ARCA: Astroparticle Research with Cosmics in the Abyss. One building block will be configured to precisely measure atmospheric neutrino oscillations. This building block is referred to as ORCA: Oscillation Research with Cosmics in the Abyss. ARCA will be realised at the Capo Passero site and ORCA at the Toulon site. Due to KM3NeT's flexible design, the technical implementation of ARCA and ORCA is almost identical. The deep-sea sites are linked to shore with a network of cables for electrical power and high-bandwidth data communication. On site, shore stations are equipped to provide power, computing and a high-bandwidth internet connection to the data repositories. The readout of the detectors is based on the "All-data-to-shore" concept, pioneered in ANTARES. In this, all analogue signals from the photo-multiplier tubes are digitised and all digital data are sent to shore where they are processed in real-time by a farm of commodity PCs. The overall design allows for a flexible and cost-effective implementation of the Research Infrastructure. The costs to realise ARCA and ORCA amount to 95 M€ in addition to Phase-1 (see below), whereas the operational costs are estimated at about 2 M€ per year.

The whole project is organised in a single Collaboration with a central management and common data analysis and repository centres. A Memorandum of Understanding for the first phase (Phase-1), covering the currently available budget of about 31 M€, has been signed by the representatives of the corresponding funding agencies. During Phase-1, data analysis tools have been developed; assembly sites for the production of optical modules and strings have been setup; and deployment and connection of strings in the deep sea are being optimised for speed and reliability. During the next phase (Phase-2.0), the Collaboration will complete the construction of ARCA and ORCA by 2020. The ultimate goal is to fully develop the KM3NeT Research Infrastructure to comprise a distributed installation at the three foreseen sites (Phase-3). The phased implementation of the KM3NeT Research Infrastructure is summarised in Table 1. The Collaboration aspires to establish a European Research Infrastructure Consortium (ERIC) hosted in the Netherlands.

The first part of this document focuses on the technical design of the infrastructure. As a preview to the science objectives presented later in this document, Fig. 1 shows the significance as a function of time for the detection of a diffuse, flavour-symmetric neutrino flux corresponding to the result reported by IceCube. Thanks to the purity of the event sample, a high-significance detection of this neutrino flux will be possible for both track-like and cascade-like events within one year of operation. The excellent angular and energy resolutions, combined with the large effective mass, provide for a significant discovery potential to find neutrino sources in the Universe. Fig. 2 shows the significance as a function of observation time for the determination of the neutrino mass hierarchy. A determination of the neutrino mass hierarchy with at least 3-sigma significance can be made after three years of operation, i.e. as early as 2023. This precedes results of other experiments and provides timely input for experiments aiming at a measurement of the CP-violation phase with high sensitivity. In addition, ORCA will provide improved measurements of some of the neutrino oscillation parameters.

¹<http://www.km3net.org>

Phase	Total costs (cumulative) M€	Primary deliverables
1	31	Proof of feasibility and first science results;
2.0	125	Study of the neutrino signal reported by IceCube; Determination of the neutrino mass hierarchy;
3	220–250	Neutrino astronomy including Galactic sources;

Table 1: Phased implementation of the KM3NeT Research Infrastructure (see text). The quoted costs for each phase include the costs of the previous phase(s). The funds for Phase-1 are secured.




Figure 1: Significance as a function of time for the detection of a diffuse flux of neutrinos corresponding to the signal reported by IceCube, for cascade-like events (red line) and track-like events (black line). The black and red bands represent the uncertainties due to the conventional and prompt component of the atmospheric neutrino flux, respectively. The blue line indicates the result of the combined analysis.




Figure 2: Median significance as a function of time for the determination of the neutrino mass hierarchy. The different lines denote expectations for different combinations of hierarchy and atmospheric mixing angle θ_{23} . Note that the CP-violating phase δ_{CP} has been assumed to be zero.

1 Detector Design and Technology

The successful deployment and operation of the ANTARES neutrino telescope [1] has demonstrated the feasibility of performing neutrino studies with large volume detectors in the deep sea. The detection of neutrinos is based on the detection of Cherenkov light produced by relativistic particles emerging from a neutrino interaction. The same technology can be used for studying neutrinos from GeV (for KM3NeT/ORCA) to PeV energies and above (for KM3NeT/ARCA). The KM3NeT design builds upon the ANTARES experience and improves the cost effectiveness of its design by about a factor four.

The goal of the KM3NeT technology is to instrument, at minimal cost and maximal reliability, the largest possible volume of seawater with a three dimensional spatial grid of ultra-sensitive photo-sensors, while remaining sensitive to neutrino interactions in the target energy range. The system should provide nanosecond precision on the arrival time of single photons, while the position and orientation of the photo-sensors must be known to a few centimetres and few degrees, respectively. The photo-sensors and the readout electronics are hosted within pressure-resistant glass spheres, so called digital optical modules (DOMs). The DOMs are distributed in space along flexible strings, one end of which is fixed to the sea floor and the other end is held close to vertical by a submerged buoy. The concept of strings is modular by design. The construction and operation of the research infrastructure thus allows for a phased and distributed implementation.

A collection of strings forms a single KM3NeT building block. The modular design allows building blocks with different spacings between lines/DOMs to be constructed, in order to target different neutrino energies. The full KM3NeT telescope comprises seven building blocks distributed on three sites. For Phase-2.0, three building blocks are planned: two KM3NeT/ARCA blocks, with a large spacing to target astrophysical neutrinos at TeV energies and above; and one KM3NeT/ORCA block, to target atmospheric neutrinos in the few-GeV range.

1.1 KM3NeT/ARCA: deep sea and onshore infrastructures




Figure 3: Map of the Mediterranean Sea close to Sicily, Italy. The cable and the location of the KM3NeT-Italy installation are indicated (left). Layout of an ARCA building block (right).

The KM3NeT-Italy infrastructure is located at $36^{\circ} 16' N 16^{\circ} 06' E$ at a depth of 3500 m, about 100 km offshore from Porto Palo di Capo Passero, Sicily, Italy (Fig. 3, left). The site is the former NEMO site and is shared with the EMSO facility for Earth and Sea science research.

The ARCA installation comprises two KM3NeT building blocks. Fig. 3 right illustrates the layout of a single 115 string building block. The power/data are transferred to/from the infrastructure via a single main electro-optic cable comprising 24 optical fibres and a single power conductor (the return is via the sea). The main electro-optic cable is connected in a ring to four primary junction boxes (Fig. 4, left). Each primary junction box is connected to three secondary junction boxes (Fig. 4, right). Each secondary junction box allows the connection of up to 12 KM3NeT detection strings. The underwater connection of the strings to




Figure 4: Photograph of the primary junction box after deployment on the seabed (left). Photograph of two secondary junction boxes on the boat prior to deployment (right).

the junction boxes is via interlink cables running along the seabed. For the ARCA configuration, the average horizontal spacing between detection strings is about 95 m.

The main electro-optic cable is connected to a power supply (50 kVA) located in the shore station at Porto Palo di Capo Passero. Power is transferred at 10 kVDC and is converted to 375 VDC at the primary junction boxes for transmission, via the secondary junction boxes, along the interlink cables to the strings.

The shore station also hosts the data acquisition electronics and a commodity PC farm used for data filtering. In December, 2008, the main electro-optic cable was deployed. A primary junction box and two secondary junction boxes were successfully connected in summer 2015.

1.2 KM3NeT/ORCA: deep sea and onshore infrastructures




Figure 5: Map of the Mediterranean Sea south of Toulon, France. The location of the KM3NeT-France and ANTARES installations are indicated (left). Layout of the full ORCA array (right).

The KM3NeT-France infrastructure is located at $42^{\circ} 48' N 06^{\circ} 02' E$ at a depth of 2450 m, about 40 km offshore from Toulon, France (see Fig. 5, left). The site is outside of the French territorial waters and about 10 km west of the site of the existing ANTARES telescope.

Fig. 5 right illustrates the layout of the full ORCA array; a single KM3NeT building block of 115 strings. The power/data are transferred to/from the infrastructure via two main electro-optic cables each comprising 36/48 optical fibres and a single power conductor (the return is via the sea).

The strings are connected to six junction boxes (Fig. 6, left), located on the periphery of the array (three on each main electro-optic cable). Each junction box has eight connectors, each of which can power four




Figure 6: Photograph of a KM3NeT-France junction box (left). Schematic of the connections to the junction box (right).

strings daisy chained in series (Fig. 6, right). Some daisy chains include calibration units, which incorporate laser beacons and/or hydrophone acoustic emitters. In the baseline design, five connectors on the junction box are dedicated for the neutrino array and one is dedicated for Earth and Sea science sensors and two are spares. The underwater connection of the strings to the junction box is via interlink cables running along the seabed. For the ORCA configuration, the average horizontal spacing between detection strings is about 20 m.

Due to the shorter transmission distance involved in the ORCA configuration, it is feasible to transfer power in Alternating Current. The power station, dimensioned for a single building block (92 KVA) is located at the shore end of the main cable near the 'Les Sablettes' beach. Power is transferred at 3500 VAC. The offshore junction boxes use a AC transformer to convert this to 400 VAC for transmission along the interlink cables to the strings. The control room is located at the Institute Michel Pacha, La Seyne-sur-Mer, and hosts the data acquisition electronics and a commodity PC farm used for data filtering.

In December, 2014, the first main electro-optic cable was successfully deployed by Orange Marine. Once ANTARES is decommissioned, its main electro-optic cable will be reused for ORCA. The first junction box was connected in spring 2015.

1.3 Detection string

The detection strings [2] (Fig. 7) each host 18 DOMs. For KM3NeT/ARCA, each is about 700 m in height, with DOMs spaced 36 m apart in the vertical direction, starting about 80 m from the sea floor. For KM3NeT/ORCA, each string is 200 m in height with DOMs spaced 9 m apart in the vertical direction, starting about 40 m from the sea floor. Each string comprises two thin (4 mm diameter) parallel Dyneema[®] ropes to which the DOMs are attached via a titanium collar. Additional spacers are added in between the DOMs to maintain the ropes parallel. Attached to the ropes is the vertical electro-optical cable, a pressure balanced, oil-filled, plastic tube that contains two copper wires for the power transmission (400 VDC) and 18 optical fibres for the data transmission. At each storey two power conductors and a single fibre are branched out via the breakout box. The breakout box also contains a DC/DC converter (400 V to 12 V). The power conductors and optical fibre enter the glass sphere via a penetrator.

Even though the string design minimises drag and itself is buoyant, additional buoyancy is introduced at the top of the string to reduce the horizontal displacement of the top relative to the base for the case of large sea currents.

For deployment and storage, the string is coiled around a large spherical frame, the so-called launcher vehicle, in which the DOMs slot into dedicated cavities (see Fig. 8). The anchor at the bottom of the string is the interface with the seabed infrastructure. It is external to the launcher vehicle and is sufficiently heavy to keep the string fixed on the seabed. The anchor houses an interlink cable, equipped with a wet-mateable connectors, and the base container. The base container incorporates dedicated optical components and an acoustic receiver used for positioning of the detector elements.

A surface vessel (Fig. 9 (left), Fig. 10 (left)), with dynamic positioning capability, is used at each site to deploy the launcher vehicle at its designated position on the seabed with an accuracy of 1 m. A remotely



Figure 7: The detection string (left) and the breakout box and the fixation of the DOM on the two parallel Dyneema® ropes (right).

operated vehicle (Fig. 9, Fig. 10, right) is used to deploy and connect the interlink cables from the base of a string to the junction box. Once the connection to the string has been verified onshore, an acoustic signal from the boat triggers the unfurling of the string. During this process, the launcher vehicle starts to rise to the surface while slowly rotating and releasing the DOMs. The empty launcher vehicle floats to the surface and is recovered by the surface vessel. The use of compact strings allows for transportation of many units onboard and thus multiple deployments during a single cruise. This method reduces costs and also has advantages in terms of risk reduction for ship personnel and material during the deployment. It also improves tolerance to rough sea conditions.

In May 2014, a prototype string comprising three active DOMs was successfully deployed and connected to the KM3NeT-Italy site and operated for more than one year [3]. This test deployment validated many aspects of the deployment scheme. The first ORCA-style string will be connected to KM3NeT-France infrastructure spring 2016.

1.4 Digital optical module

The Digital Optical Module [4] (Fig. 11 left) is a transparent 17 inch diameter glass sphere comprising two separate hemispheres, housing 31 photo-multiplier tubes (PMT) and their associated readout electronics. The design of the DOM has several advantages over traditional optical modules using single large PMTs, as it houses three to four times the photo-cathode area in a single sphere and has an almost uniform angular coverage. As the photo-cathode is segmented, the identification of more than one photon arriving at the DOM can be done with high efficiency and purity. In addition, the directional information provides improved rejection of optical background.

The PMTs are arranged in 5 rings of 6 PMTs plus a single PMT at the bottom pointing vertically downwards. The PMTs are spaced at 60° in azimuth and successive rings are staggered by 30° . There are 19 PMTs in the lower hemisphere and 12 PMTs in the upper hemisphere. The PMTs are held in place by a 3D printed support. The photon collection efficiency is increased by 20–40% by a reflector ring around the




Figure 8: Photo of a launch vehicle deployment (left). Principle of the launch vehicle unfurling (right, picture courtesy Marijn van der Meer/Quest).




Figure 9: Photograph of the Ambrosius Tide boat, used for the KM3NeT/ARCA string deployment (left). Photograph of the remote operated vehicle, used for the string connection (right).

face of each PMT. In order to assure optical contact, an optical gel fills the cavity between the support and the glass. The support and the gel are sufficiently flexible to allow for the deformation of the glass sphere under the hydrostatic pressure.

Each PMT has an individual low-power high-voltage base with integrated amplification and tuneable discrimination. The arrival time and the time-over-threshold (ToT) of each PMT, are recorded by an individual time-to-digital converter implemented in a FPGA. The threshold is set at the level of 0.3 of the mean single photon pulse height and the high voltage is set to provide an amplification of 3×10^6 . The FPGA is mounted on the central logic board, which transfers the data to shore via an Ethernet network of optical fibres. Each DOM in a string has a dedicated wavelength to be later multiplexed with other DOM wavelengths for transfer via a single optical fibre to the shore. The broadcast of the onshore clock signal, needed for time stamping in each DOM, is embedded in the Gb Ethernet protocol. The white rabbit protocol has been modified to implement the broadcast of the clock signal. The power consumption of a single DOM is about 7W.

The specification for the PMTs are summarised in Tab. 2. Prototype PMTs from Hamamatsu and ETEL




Figure 10: Photo of the Castor boat, used for the KM3NeT/ORCA string deployment (left). Photo of the Comex Apache ROV used for the KM3NeT/ORCA string connection (right).




Figure 11: A photograph of a completed Digital Optical Module. The central white spot is the acoustic piezo sensor (left). Exploded view of the inside of a DOM (right).

have been developed and satisfy the requirements (see Fig. 12). The PMTs have a photo-cathode diameter of at least 72 mm and a length of less than 122 mm. The reflector effectively increases the diameter to about 85 mm. The photo-multiplier tube has a ten stage dynode structure with a minimum gain of 10^6 . The front face of the photo-multiplier tube is convex with a radius smaller than the inner radius of the glass sphere. Due to the small size of the PMT, the influence of the Earth's magnetic field is negligible and a mu-metal shield is not required.



Figure 12: Photographs of the PMTs. ETEL D792KFL (left) and Hamamatsu R12199-02 (right).

The optical module also contains three calibration sensors: 1) The LED nano-beacon, which illuminates

Radiant blue sensitivity at 404 nm	130 mA/W
Quantum efficiency (QE)	20% @ 470 nm and 28% @ 404 nm
Inhomogeneity of cathode response	10%
Supply voltage for a gain of 3×10^6	900–1300 V
Dark count at 15°C and 0.3 p.e. threshold	1.5 kHz
Transit time spread (TTS)	4.5 ns (FWHM)
Peak to valley ratio	2.5

Table 2: Specification of the PMTs.

the optical module(s) vertically above; 2) A compass and tilt-meter for orientation calibration; 3) An acoustic piezo sensor glued to the inner surface of the glass sphere for position calibration.

In May 2013, a prototype DOM was successfully installed on an ANTARES detection line and operated in-situ for over a year [5]. Starting in spring 2014, three prototype DOMs were operated for over a year at the KM3NeT-Italy site [3]. In December 2015, a first production string of 18 DOMs was successfully operated at the KM3NeT-Italy site.

1.5 Fibre-optic data transmission system

The KM3NeT fibre-optic data transmission system performs the following functions:

- Transfers all the data to shore: The bandwidth per DOM is 1 Gb/s. The observed singles rate, dominated by ^{40}K , is typically 8 kHz per PMT [5] or 250 kHz per DOM which amounts to 12 Mb/s per DOM. Additional contributions from bioluminescence can be accommodated up to levels of a factor of 10 compared to ^{40}K .
- Provide timing synchronisation: Relative time offsets between any pair of DOMs are stable within 1 ns;
- Provide individual control for each DOM: Setting the HV of a PMT, turn off/on a single PMT, turn on/off nano-beacon, update soft- and firmware;
- Provide individual control for each base of a string: Turn string power on/off, control optical amplifiers, monitor AC/DC converter;
- Provide slow control for the junction boxes.

The slow-control system is implemented via a broadcast mechanism (same as that of the clock), in which control information for all strings is sent on a single common wavelength. If it is a message for just a single string or DOM it is ignored by all the others. The communication from offshore exploits a Dense Wavelength Division Multiplexing (DWDM) technique. The return signals for the slow control are transmitted on 34 wavelengths via the slow-control fibre(s). The data return path is based on a 50 GHz spacing system with a 72 wavelengths uplink. Each DOM of 4 strings produces a unique wavelength that is combined on one fibre. EDFA optical amplifiers are introduced onshore and at the base of a string to maintain the optical margins above 10 dB.

1.6 Data acquisition

The readout [6] of the KM3NeT detector is based on the “All-data-to-shore” concept in which all analogue signals from the PMTs that pass a preset threshold (typically 0.3 p.e.) are digitised and all digital data are sent to shore where they are processed in real time. The physics events are filtered from the background using designated software. To maintain all available information for the offline analyses, each event will contain a snapshot of all the data in the detector during the event. Different filters can be applied to the data simultaneously.

The optical data contain the time of the leading edge and the time over threshold of every analogue pulse, commonly referred to as a hit. Each hit corresponds to 6 Bytes of data (1 B for PMT address, 4 B for time and 1 B for time over threshold). The least significant bit of the time information corresponds to 1 ns. The total data rate for a single building block amounts to about 25 Gb/s. A reduction of the data

rate by a factor of about 10^5 is thus required to store the filtered data on disk. In addition to physics data, summary data containing the singles rates of all PMTs in the detector are stored with a sampling frequency of 10 Hz. This information is used in the simulations and the reconstruction to take into account the actual status and optical background conditions of the detector.

In parallel to the optical data, the data from the acoustics positioning system are processed and represents a data volume of about one third of that of the optical data.

1.6.1 Event trigger

For the detection of muons and showers, the time-position correlations that are used to filter the data follow from causality. In the following, the level-zero filter (L0) refers to the threshold for the analogue pulses which is applied off shore. All other filtering is applied on shore. The level-one filter (L1) refers to a coincidence of two (or more) L0 hits from different PMTs in the same optical module within a fixed time window. The scattering of light in deep-sea water is such that the time window can be very small. A typical value is $\Delta T = 10$ ns. The estimated L1 rate per optical module is then about 1000 Hz, of which about 600 Hz is due to genuine coincidences from ^{40}K decays. The remaining part arises from random coincidences which can be reduced by a factor of two by making use of the known orientations of the PMTs. This is referred to as the level-two filter (L2). Separate trigger algorithms operate in parallel on this data, each optimised for a different event topology.

A general solution to trigger on a muon track event consists of a scan of the sky combined with a directional filter [7]. In the directional filter, the direction of the muon is assumed. For each direction, an intersection of a cylinder with the 3D array of optical modules can be considered. The diameter of this cylinder (i.e. road width) corresponds to the maximal distance traveled by the light. It can safely be set to a few times the absorption length without a significant loss of the signal. The number of PMTs to be considered is then reduced by a factor of 100 or more, depending on the assumed direction. Furthermore, the time window that follows from causality is reduced by a similar factor². This improves the signal-to-noise ratio (S/N) of an L1 hit by a factor of (at least) 10^4 compared to the general causality relation. With a requirement of five (or more) L1 hits, this filter shows a very small contribution of random coincidences.

The field of view of the directional filter is about 10 degrees. So, a set of 200 directions is sufficient to cover the full sky. By design, this trigger can be applied to any detector configuration. Furthermore, the minimum number of L1 hits to trigger an event can be lowered for a limited number of directions. A set of astrophysical sources can thus be tracked continuously with higher detection efficiency for each source.

For shower events, triggering is simpler, since the maximal 3D-distance between PMTs can be applied without consideration of the direction of the shower.

A maximum distance travelled by the light can be assumed, limiting the maximum distance D between hit PMTs. This reduces the number of PMTs to be considered and the time window that follows from causality. Hence, an improvement of the S/N ratio compared to the general causality relation can be obtained.

Alternative signals with different time-position correlations, such as slow magnetic monopoles, can be searched for in parallel. It is obvious but worth noting that the number of computers and the speed of the algorithms determine the performance of the system and hence the physics output of KM3NeT.

1.6.2 Performance

The performance of the online data filter can be summarised by the effective volume, the event purity and the time needed to process a time slice of raw data. The effective volume is the volume in which a neutrino interaction would trigger the event to be written to disk and the event purity is the fraction of events that contain a neutrino interaction or atmospheric muon bundle. The effective volume of the ARCA and ORCA detectors are presented in Sec. 2.2 and Sec. 3.2.2, respectively.

To process the data, the concept of time slicing is applied. In this, the data from each optical module are stored in a frame corresponding to a preset time period. All data frames corresponding to the same time period are sent to a single CPU core based on IP level 2 switching. A complete set of data frames is referred to as a time slice. Data corresponding to subsequent time periods are sent to different CPU cores

²Only the transverse distance between the PMTs should be taken into account because the propagation time of the muon can be corrected for.




Figure 13: Trigger rate due to random coincidences (left) and required number of CPU cores (right) as a function of the singles rate.

until the number of available CPU cores is exhausted. The first CPU core should then be ready to receive and process the data from the next time period.

In the following, the performance of the online data filter is presented for a one ARCA and one ORCA building block. In this, different triggers are operated in parallel. The typical trigger settings correspond to a L1 time window of $\Delta T = 10$ ns, a maximum space angle between the PMT axes of 90 degrees (L2), and a minimum number of L1 hits of 4 or 5. The detection threshold thus corresponds to 8 or 10 photons. The trigger rate due to random coincidences and the number of CPU cores are shown in Fig. 13 as a function of the singles rate.

The typical singles rate due to radioactive decays and bioluminescence in the sea water, including the dark count of the PMT, is about 6–8 kHz. To limit the average singles rate, short bursts of bioluminescence can be filtered in real-time. The number of CPU cores needed to process the data in real time is less than 50 up to singles rates of 20 kHz. It should be noted that the number of CPU cores may be larger than one for a modern PC.

2 Astroparticle Research with Cosmics in the Abyss (ARCA)

2.1 Introduction

The main science objective of KM3NeT/ARCA is the detection of high-energy neutrinos of cosmic origin. Since neutrinos propagate directly from their sources to the Earth, even modest numbers of detected neutrinos can be of utmost scientific relevance, by indicating the astrophysical objects in which cosmic rays are accelerated, or pointing to places where dark matter particles annihilate or decay. The prospect of such fundamental physics discoveries have led the astroparticle and astrophysics communities to include KM3NeT as a high priority in their respective European road maps (APPEC/ASPERA, AstroNet) and the European Strategy Forum on Research Infrastructures (ESFRI) to include it in their list of priority projects. The KM3NeT Research Infrastructure will also provide user ports for real-time, long-term Earth and Sea science measurements in the deep-sea environment.

One priority goal of KM3NeT/ARCA is indisputably to find neutrinos from the cosmic ray accelerators in our Galaxy. The search strategy is to identify upward-moving muons, which unambiguously indicate neutrino reactions since only neutrinos can traverse the Earth without being absorbed. A neutrino telescope in the Mediterranean Sea is ideal for this purpose, since most of the potential Galactic sources are in the Southern sky; in contrast, the IceCube detector at the South Pole is much less sensitive to these individual sources, at least in the energy range where the signal is expected (a few TeV to a few tens of 10 TeV – see Sec. 2.3.3). The KM3NeT/ARCA design has been carefully optimised to maximise the sensitivity to these Galactic sources. One of the findings in this process is that the overall sensitivity is not reduced if the neutrino telescope is split into separate building blocks, provided they are large enough, at least 0.5 cubic kilometres each [8]. It has thus been decided to make a distributed infrastructure, thereby maximising the influx of regional funding and human resources. Furthermore, the concept of independent building blocks complies with the technical specifications for the construction and operation of the Research Infrastructure.

Currently, the KM3NeT Collaboration is proceeding with the first construction phase (Phase-1). Until 2017, 31 strings equipped with 558 optical modules (see Sec. 1) will be assembled and deployed. Of these, 24 strings will be configured for ARCA and deployed at the Italian site. The resulting array will provide the equivalent of 10–20% of the size of the IceCube detector. The recent experience from a combined analysis of ANTARES and IceCube data [9], increasing the sensitivity to point-like neutrino sources by up to a factor of two with respect to the individual analyses, indicates that Phase-1 will already have a decent discovery potential and provide significant new data.

2.1.1 Cosmic neutrinos

A new situation has emerged since IceCube has presented evidence for a neutrino signal of cosmic origin. This signal includes upward- and downward-going events with neutrino energies from a few 10 TeV to above 1 PeV. Even though the signal is statistically very significant, its astrophysical implications are not yet clear. This signal is the first high-energy extra-terrestrial neutrino signal ever observed and thus marks a major turning point in the history of neutrino astronomy. Detailed studies have been and are being conducted to estimate the sensitivity of KM3NeT/ARCA to a neutrino flux with the reported properties, to investigate the consequences of a re-optimisation of the detector for such a signal (in particular in terms of geometry parameters and building block size) and to evaluate the prospects of Phase-2.0. Results of these studies are presented in the following.

IceCube's high-energy starting event (HESE) analysis [10] has now observed 54 events with a reconstructed energy above 30 TeV, 39 of which are identified as cascades and 14 as track events [11]³. Most of the observed events originate from the Southern hemisphere, corresponding to down-going neutrinos in the IceCube detector. Due to the different topologies of the events, the angular resolution is roughly 10–15° for cascades and 1° for muons. The expected background due to atmospheric muons and neutrinos is about 12 and 9 events respectively, resulting in a significance of over 5 σ for the observation to be incompatible with the background. This significance has been obtained by applying designated event selection cuts using the outer layers of the detector as veto against incoming charged particles. The best constraints on an (assumed diffuse) astrophysical spectrum come from a maximum likelihood analysis using both HESE and other event samples [12], finding a neutrino flux proportional to $E^{-2.5}$, disfavouring at 2.1 σ an E^{-2} spectrum with a

³One of them has been identified as a coincident air-shower event.




Figure 14: Footprint of one building block of the ARCA detector.

cutoff at a few PeV. The distribution of the neutrino directions combined with the angular resolution does not (yet) allow for the identification of one or more point sources. Deviations from flavour-uniformity are only weakly constrained [13], and tau neutrino events have not yet been identified [14, 15].

The prime physics case for KM3NeT Phase-2.0 ARCA is to measure and investigate the signal of neutrinos observed by IceCube with different methodology, improved resolution and a complementary field of view.

2.1.2 Assumptions

The basic assumption in the following studies is that the ARCA detector will comprise two KM3NeT building blocks, providing an instrumented volume of about one cubic kilometre, i.e. of similar size as the IceCube detector. All analyses reported in this document are performed for a horizontal distance between strings of 90 m and vertical distance between adjacent optical modules of 36 m. The footprint of one block is shown in Fig. 14. To estimate the dependence of the sensitivity on the geometrical detector configuration, an alternative layout with 120 m distance between strings but unchanged vertical distances is being investigated; this configuration corresponds to an increase of the instrumented volume to 1.7 km³. In both cases a water depth of 3.5 km and a latitude of 36° 16' N were assumed, corresponding to the Italian KM3NeT site (KM3NeT-It, see Sec. 1.1).

The following sensitivity studies are discussed in the following:

- **Cascade events from a diffuse flux, including high-energy starting muon tracks**

This analysis includes all neutrino flavours. Owing to an efficient suppression of the atmospheric muon and neutrino backgrounds (see below), a 4π angular coverage has been achieved.

- **Upgoing, diffuse flux of muon (anti-)neutrinos**

This analysis is usually referred to as the “conventional” diffuse flux analysis. Traditionally, it does not include the upper hemisphere, with the exception of a small zenith region above the horizon.

- **Muon (anti-)neutrinos from a diffuse Galactic plane flux**




Figure 15: Atmospheric neutrino fluxes as a function of the neutrino energy. The bands represent the uncertainties in the conventional (red and black bands) and in the prompt (green and blue bands) components assumed in this work (see text).

Upgoing muon track events are used for an analysis covering an extended region of the Galactic plane near the Galactic centre in the Southern sky.

- **Upgoing flux of muon (anti-)neutrinos from point sources**

In order to quantify the sensitivity of KM3NeT Phase-2.0 to extragalactic and Galactic point sources of neutrinos, both a generic E^{-2} spectrum from point sources and spectra with energy cut-off for specific Galactic sources with non-zero radial extension have been considered.

- **Cascade events from point sources**

KM3NeT/ARCA's resolution in the cascade channel will allow us to use these events in point-source searches. The sensitivity of such an analysis is evaluated against generic E^{-2} point-sources.

The background of atmospheric neutrinos assumed in these analyses corresponds to the so-called Honda flux [16] with a prompt component as calculated by Enberg [17]. A correction taking into account the "knee" of the cosmic ray spectrum has been applied to both conventional and prompt atmospheric neutrino fluxes according to the prescription in [18] and references therein. The Honda parameterisation includes an anisotropy caused by the Earth's magnetic field, while the prompt component is assumed to be isotropic in the full solid angle. Moreover, in the sensitivity studies the effect of the uncertainties on the atmospheric neutrino flux has been estimated. An uncertainty of $\pm 25\%$ was assumed for the intensity of the conventional Honda flux. For the prompt component, the uncertainty band estimated in [17] has been used. Recently a new calculation of the prompt component has been reported in [19]. In this calculation constraints set by the charm measurement from the LHCb experiments at 7 TeV have been taken into account. In Fig. 15 the different components of the atmospheric neutrino flux is reported for ν_e and ν_μ ; see Sec. 2.2 for details on the background from atmospheric muons.

It should be noted that the results reported in the following are preliminary and some analysis details are not yet fully completed and optimised. Also, the analyses reported above do not reflect the full physics potential of ARCA; the event resolutions shown in Sec. 2.2.4 can be used to characterise ARCA's ability to probe any assumed extraterrestrial neutrino fluxes.

2.2 Simulations

Monte Carlo simulations have been used to simulate the detector response to particles incident on the detector, their interaction with the medium surrounding the detector and subsequent Cherenkov light production, and the detector response in terms of the PMT data sent to shore. The software packages described in this section have mostly been developed in the ANTARES Collaboration and then adapted to KM3NeT. The simulation is based on the nominal detector geometry described in Sec. 2.1.2 and Fig. 14 – see Sec. 1.1 for further details. Each of the two ARCA blocks are treated identically and independently – simulations are performed for a single block, and the effective lifetime (event rate) is multiplied by two. The effects of position and orientation calibration uncertainties are estimated using dedicated simulations, as described in Sec. 2.4.

2.2.1 Event generation

The relevant volume for Cherenkov light production is defined as a cylinder with height and radius of about 3 absorption lengths larger than the instrumented volume (the “can”), limited by the seabed below. The first step in the simulation chain is the generation of particle fluxes incident on the can – neutrinos from astrophysical sources, and the atmospheric muon and neutrino backgrounds – within which a detailed description of particle behaviour and Cherenkov light production is required.

Astrophysical and atmospheric fluxes of (anti-)neutrinos of all three flavours (ν_e , ν_μ , and ν_τ) are simulated with a code propagating neutrinos through the Earth (density profile from [20]) and generating their interactions in rock and sea water. For reactions outside the can, long-range interaction products (muons and taus) are subsequently propagated to the can. Both neutral-current (NC) and charged-current (CC) reactions are simulated. The deep inelastic scattering (DIS) cross-sections, which are dominant in the energy range relevant to this study, are implemented using the LEPTO code [21]. The CTEQ6D table of parton distribution functions is used, and the resulting behaviour – especially in the small- x region – validated up to 10 PeV. Quasi-elastic scattering and resonance production are also taken into account, by using RSQ [22] below 300 GeV. Reactions of $\bar{\nu}_e$ with electrons in the atmosphere are relevant in the energy regime of resonant W production (“Glashow resonance”) around 6.3 PeV and are simulated according to the leading-order electroweak cross sections. The propagation of muons in rock and water is performed with MUSIC [23]. Tau leptons, which have a life time of 2.9×10^{-13} s and thus typically travel only very short distances before decaying, are propagated by assuming them to be minimally ionising particles, and decayed using TAUOLA [24].

Atmospheric muons, produced in cosmic-ray interactions in the atmosphere, can penetrate to the detector volume if their energy at the sea surface is in the TeV range or above. This is frequently the case, both for single muons, and muon ‘bundles’ up to several hundred muons from a primary cosmic ray event. Atmospheric muons therefore establish an important, high-rate background that is simulated using MUPAGE [25]. Single and multiple atmospheric muon events are generated using a parameterisation of the flux of muon bundles at different depths and zenith angles. In the present analysis, three simulated atmospheric muon event samples are used, with muon bundle energies exceeding 1 TeV, 10 TeV, and 50 TeV, respectively, in order to provide sufficient coverage in the high-energy regime. The corresponding lifetimes of these and the neutrino productions are shown in Fig. 16.

The correlated flux of atmospheric neutrinos and muons from the same primary cosmic ray interaction is simulated with CORSIKA v7.4001 [26], in order to investigate the ‘self-veto’ effect [27] for high-energy studies. GHEISHA [28] and QGSJET01 [29] were respectively used to model low- and high-energy hadronic interactions, and the curvature of the Earth was accounted for. Both muons and neutrinos are recorded at sea-level; muons are propagated to and through the can with MUSIC, while one neutrino from each event is forced to interact. The intention was to estimate the effect of accompanying muons on high-energy atmospheric neutrino events (Sec. 2.2.7) – thus, only events with at least one muon at can level, and one neutrino above 10 TeV, are kept, which excludes all up-coming neutrino events⁴. The resulting event sample forms only a small fraction of all atmospheric muon bundles, but a significant fraction of all downgoing atmospheric neutrino background events above 10 TeV. Therefore, analyses using CORSIKA events down-weight the standard atmospheric neutrino events to avoid double-counting. Additionally, CORSIKA underestimates the expected atmospheric neutrino flux at high energies [16, 17], and this is corrected for as per [30].

⁴Events with only a neutrino or atmospheric muon bundle are already simulated using standard methods




Figure 16: Effective simulated lifetimes for neutrinos: ν_μ NC (cascade-like), ν_μ CC (track-like), for both the IceCube diffuse flux [12] and atmospheric spectra; atmospheric μ events; and cosmic ray (CR) events from CORSIKA, as a function of neutrino / muon-bundle / cosmic-ray energy E . The lifetimes for other neutrino channels (ν_e and ν_τ) are similar to that of ν_μ NC, except for the atmospheric ν_τ events, which have effectively infinite lifetime (since the estimated flux is very small).

2.2.2 Detector response

The particles propagated to the can level, or generated inside the can volume, are then tracked in the sea water using tabulated results from full GEANT 3.21 simulations of relativistic muons and electromagnetic cascades to generate the number of Cherenkov photons detected by the KM3NeT photomultiplier tubes (PMTs). The light production from hadronic or mixed hadronic/electromagnetic cascades is scaled to that from purely electromagnetic cascades according to the energy and type of constituent particles. The program takes into account the full wavelength dependence of Cherenkov light production, propagation, scattering and absorption; and the response of the PMTs as described in Sec. 1 and modelled in [31], including absorption in the glass and optical gel, the PMT quantum efficiency, the reduced effective area for photons arriving off-axis, and the effect of the reflecting expansion cones [32].

Hits from background photons (mostly due to ^{40}K decays in the sea water) in an event are simulated by adding random noise hits with a rate of 5 kHz per PMT. Correlated hits over multiple PMTs on the same optical module from single ^{40}K decays are also included, with $\{2, 3, 4\}$ -fold coincidences at rates of $\{500, 50, 5\}$ Hz per DOM. The singles and coincidence rates as well as the angular dependence are in reasonable agreement with the results from the prototype detection unit deployed at the KM3NeT-It site [3]. An example of the simulated time-distribution of photons detected by a KM3NeT PMT from a 1 TeV muon 50 m from the track is given in Fig. 17.

KM3NeT PMT hits are recorded via the start time and the duration of the signal above a predefined threshold (time-over-threshold, or ToT). This scheme is implemented in the detector simulation, with the simulated response of individual PMTs to photon hits being based on laboratory measurements. The full transit-time distribution is implemented on a per-photon basis, corresponding approximately to a 2 ns Gaussian smearing for the majority of photons. Hit amplitudes are smeared, and the start time and ToT are calculated by accounting for sequences of photo-electrons on PMTs that cannot be resolved in time, saturation effects at around 40 simultaneous photoelectrons, and a maximum ToT readout of 255 ns. After this step, each event contains a complete and unbiased snapshot of all hits recorded during a time window around the event, representing a part of the stream of data sent to shore.

The final stage is to simulate on-shore triggering, as described in Sec. 1.6.1. This process takes filtered L1 hits (photon hits on multiple PMTs within a short time window on the same OM) and generates a trigger if multiple nearby OMs record such events at causally connected times within a spherical (cascade) or cylindrical (track) geometry. Trigger parameters have been tuned so as to minimise false triggers on




Figure 17: Simulated time distribution (relative to the nominal time of the Cherenkov shock front) of photoelectrons detected by a KM3NeT PMT from a 1 TeV muon track with closest approach distance $\rho = 50$ m. The PMT is simulated facing both towards (red) and away (blue) from the track. The optical background rate is also shown for comparison. The PMT response (time-smearing of ~ 2 ns) is not included.




Figure 18: Effective areas of ARCA (two blocks) at trigger level for ν_μ , ν_e , and ν_τ , as a function of neutrino energy E_ν . The effective area is defined relative to an isotropic neutrino flux incident on the Earth, is averaged over both ν and $\bar{\nu}$, and includes both NC and CC interactions. The peak at 6.3 PeV is due to the Glashow resonance of $\bar{\nu}_e$.




Figure 19: Simulation times per triggered event for different simulation stages, for ν_e CC (cascade) events (left), and ν_μ CC (track) events (right). The times are calculated for simulations run on a single dedicated CPU at the Centre de Calcul de l'IN2P3/CNRS.

optical backgrounds, while registering all reconstructible physics events. In the case of ARCA, the real-time trigger rate is dominated by downgoing atmospheric muons, and trigger settings were set to keep the corresponding data rate manageable.

The trigger settings correspond to a coincidence (L1) time window of $\Delta T = 10$ ns, and a minimum number of 5 L1 hits for both the shower trigger and the muon trigger. Only Monte Carlo events which pass either trigger condition are available for further analysis, as is the case for the on-shore trigger. The resulting effective areas are given in Fig. 18.

The simulation times per event for different stages are shown in Fig. 19. The simulation time is dominated by event reconstruction and light propagation, which can reach up to a few seconds per event at high energies. The cascade reconstruction time does not reduce quickly at low energies, since it includes in the likelihood fit PMTs which have no detections.

The MC events simulated with the described codes have been compared with the data from a prototype of the string that was deployed at the Italian site and that took data for about one year [3]. The very good agreement between the data and the MC simulation have demonstrated the high reliability of the MC simulation chain.

2.2.3 Further improvements

The simulation chain for ARCA is mature, but not complete, and several additions will be required for future data analysis. These are:

- The simulation of tau (anti-)neutrinos is performed using some simplifications. Charged-current tau interactions within the Earth are treated as absorbing the neutrino, i.e. the tau 'regeneration' effect is not included. Additionally, only two- and three-body tau decay modes (approximately three quarters of all decays) are currently implemented – the branching ratio of $\sim 17.4\%$ for the decay to a muon is kept constant, while other modes are re-normalised to the remaining 83.6%, and result in almost identical event topologies at high energies.
- The MUPAGE package for generation of atmospheric muons does not contain a prompt component originating from charm decays in cosmic-ray-induced air showers. The flux of atmospheric muons with energies above roughly 10 TeV is therefore underestimated, although likely only by a small amount. A refined simulation has recently been provided in the CORSIKA [26] framework, where the correlations between conventional and prompt muon and neutrino fluxes are adequately included at the event-by-event level. While a production with the new CORSIKA v7.4005 has begun, the high CPU demand has so far prevented this simulation from being fully processed through the Monte Carlo chain and used for analysis.




Figure 20: Event displays for a simulated ν_μ CC event (left) and a contained ν_μ NC event (right), showing only DOMs with a total ToT (summed over all 31 PMTs) of more than 30 ns in a narrow time window. In both cases, the incoming neutrino is indicated by the red line, and the outgoing lepton (muon or neutrino) by the green line. The colour scale gives the hit times with respect to the time of the neutrino interaction, while the size of the circles are proportional to the total ToT on each DOM. DOMs without hits are shown by grey dots.

- Atmospheric muon events which coincidentally arrive simultaneously with neutrino events have not been simulated, since it is anticipated that resolving multiple components will prove feasible for ARCA. An explicit production of coincident muon events will need to be produced in order to verify this.

2.2.4 Event reconstruction

Two broad event classes can be identified in a high energy neutrino telescope: track-like events and cascade-like events:

- The track-like events are generated by muons that are produced in the matter inside or surrounding the detector through CC interactions of ν_μ ($\bar{\nu}_\mu$) and ν_τ ($\bar{\nu}_\tau$). CC reactions of ν_τ ($\bar{\nu}_\tau$) produce a muon with a branching ratio of 17%, when the emerging τ decays into a μ .
- The cascade-like events are produced in the matter near or inside the detector volume through CC interactions of ν_e ($\bar{\nu}_e$) and ν_τ ($\bar{\nu}_\tau$) and in NC interactions of neutrinos of all flavours. CC ν_τ ($\bar{\nu}_\tau$) interactions produce cascade events with a branching ratio of 83%.

These two event classes produce very different time-space hit patterns in the detector. The cascade-like events are characterised by a very dense hit pattern close to the neutrino interaction point. A significant fraction of the neutrino energy is released in a hadronic shower (and, in the case of ν_e ($\bar{\nu}_e$) CC interactions, the rest in an electromagnetic cascade), thus allowing for a good estimate of the neutrino energy. A track-like event is characterised by the Cherenkov light from the emerging muon that can travel large distances through Earth rock and sea water. The spatial hit pattern in this case is closely related to the muon direction, thus allowing for a precise measurement of the latter. Typical hit patterns for track-like and cascade-like events are shown in Fig. 20

Starting from the ANTARES experience, algorithms that reconstruct direction, energy and interaction vertex of the neutrinos from the muon tracks or the showers have been developed. These have been optimised for 'pure' track events (ν_μ CC events far from the detector, where only a single energetic muon is observed) and for cascade events (ν NC and ν_e CC events, where only a cascade is observed), respectively. Thus their performance on more complicated event topologies is not optimal; prospects for improvements are discussed at the end of this section.

Track reconstruction Muons with energies above 1 TeV can reach track lengths of the order of kilometres and have a direction that is nearly collinear with that of the parent neutrino. To reconstruct the muon direction – and consequently the neutrino direction – an algorithm is used that maximises the likelihood




Figure 21: Left panel: median of the angle between the neutrino and the reconstructed muon direction (black line) and between the neutrino and the true muon direction (red line), for selected ν_μ CC events ($\Lambda > -5.8$, see below). The dark and light blue bands represent the 90% and 68% quantiles of the distributions. Right panel: distribution of $\log_{10}(E_{\text{reco}}/E_\mu)$, where E_{reco} is the reconstructed and E_μ is the true muon energy for events with $E_\mu \geq 10$ TeV that satisfy a containment criterion. The red line represents the Gaussian fit.

that the observed space-time PMT hit pattern is consistent with Cherenkov emission from the fitted muon trajectory. An initial hit selection exploits hit coincidences between PMTs in the same optical module or between different optical modules to remove uncorrelated hits from background photons, mostly from ^{40}K decays. The reconstruction of the muon trajectory starts with a linear fit, followed by three consecutive fitting steps, each using the results of the previous one as starting point. A pseudo-vertex position is also estimated, which, however, usually is related to the entry point of the muon in the detector rather than to the location of the interaction vertex; this quantity is useful for background rejection. In addition to the track information (direction and pseudo-vertex) an estimator of the fit quality, Λ , and the number of hits associated with the final track fit, N_{hit} , are determined. The Λ parameter is used in the analysis to reject badly reconstructed events, in particular atmospheric muons mis-reconstructed as up-going. The N_{hit} parameter is related to the muon energy and is used to reject low-energy events that are mainly due to atmospheric neutrino background. A very good angular resolution of about 0.2° is achieved for neutrinos above 10 TeV, see Fig. 21 (left).

The amount of light collected by the PMTs when a muon travels inside the detector is correlated with the muon energy. To estimate the muon energy, a method exploiting this dependence by means of an artificial neural network has been developed. The first step is the selection of events with a reconstructed muon track travelling inside the detector for an adequate distance. The second step is the evaluation of several quantities related with the total event ToT and with the number of DOMs hit. These quantities are used to feed the neural network. The energy resolution obtained for well reconstructed (cut on Λ applied) and contained events is $\simeq 0.27$ units in $\log_{10}(E_\mu)$ for $10 \text{ TeV} \leq E_\mu \leq 10 \text{ PeV}$ (see Fig. 21 right); without containment requirement, the resolution slightly worsens to $\simeq 0.28$ units. Further details on the track reconstruction code can be found in [33].

This energy reconstruction method must be trained on appropriate samples of MC events and is not yet fully integrated in the reconstruction software for ARCA. A simple energy reconstruction using the N_{hit} parameter is embedded in the reconstruction software and gives results of almost equivalent quality. This method is used for the sensitivity studies presented in the following.

2.2.5 Cascade reconstruction

The length of a cascade event depends logarithmically on the cascade energy and is of the order of 10 m in the energy range relevant for ARCA analyses. At the length scale of typical distances between optical modules, cascades thus produce almost point-like signatures, characterised by vertex position, direction, and energy. CC interactions of ν_μ and ν_τ , if they happen in the detector volume, also produce cascades, but the outgoing μ , τ , or τ decay products produce a more complex signature. Hence, cascade reconstruction is




Figure 22: ARCA vertex resolutions for contained ν_e CC events using Algorithm 1, after the event selection of Sec. 2.3.1. Left: resolution in the longitudinal direction, showing the offset from the MC vertex to the shower maximum. Right: directional resolution in the lateral direction, which gives the characteristic accuracy of ~ 0.5 m. For both plots, the black line shows the median value; dark blue shaded regions give the 68% range, while light blue shaded regions give the 90% range.

optimised for ν NC and ν_μ CC interactions, and then the performance is assessed on the latter class. Three independent algorithms have been applied to reconstruct cascade vertex position, direction and energy. The first has been specifically developed for ARCA, and exhibits the best performance. The second and third have been adapted from ANTARES analyses and have outputs which prove useful in event classification and background discrimination. All three are described here, although only the performance of the first is shown.

The first algorithm (Algorithm 1) has been specifically developed to exploit the information provided by the KM3NeT multi-PMT DOM. The hit selection is designed to be simple and to allow for a fast reconstruction. Hits on the same PMT within 350 ns are merged using the time of the first hit, and coincidences of two merged hits within 20 ns on a single DOM are used for the vertex fit. This fit minimises time-residuals assuming a spherically expanding shell of light about an assumed cascade maximum. The offset of this fit from the MC true vertex position in the longitudinal direction (Fig. 22 left) mostly measures the shower elongation, while the offset in the lateral direction (Fig. 22 right) measures the accuracy, reaching a precision well below 0.5 m in the high-energy regime.

The direction and energy are reconstructed using maximum-likelihood methods, applied to the merged hits as described above. All PMT hits within -100 ns to $+900$ ns of the expected Cherenkov light-front from the vertex fit are used. Thus each PMT only has a 0.2% chance of receiving a random hit from the optical background. Rather than fitting the ToT (\sim charge) measurement from each PMT, the algorithm simply fits the probability of a PMT recording one or more photons within this time-window, making the procedure highly robust. This probability is estimated from simulations as a function of PMT distance and pointing direction to the shower, angle from the shower axis to the PMT position, and electromagnetic-equivalent cascade energy. The strong geometrical dependence in hit probabilities allows for a very high reconstruction quality: nearby PMTs facing towards the cascade, close to the Cherenkov angle, will tend to have a hit probability of unity, while distant PMTs facing away from the cascade, far from the Cherenkov angle, will tend to have a hit probability of zero.

For contained events above 50 TeV, the $\pm 1\sigma$ energy and median direction resolutions achieved with this method are roughly 10% and 2° respectively, with no loss of efficiency. The resolutions after the selection cuts described in Sec. 2.3.1 are shown in Fig. 23. For energy above 60 TeV, corresponding to the approximate low-energy threshold of the cut-and-count diffuse-flux analysis (see Sec. 2.3.1), the 1σ energy resolution is characteristically 5%, while the median directional resolution is 1.5° . This energy resolution is close to the limit imposed by variations in the hadronic cascade component (mostly due to the variable inelasticity), which yields less Cherenkov light ($\sim 90\%$ at 100 TeV) than the electromagnetic component [34].

The second algorithm fits the vertex position from the positions and the arrival times of the PMT hits using an M-estimator procedure and applies selection cuts on the resulting quality parameters. The cascade direction is determined from the average direction of hits with respect to the vertex position, the energy




Figure 23: ARCA resolutions for contained ν_e CC events using Algorithm 1, after the event selection of Sec. 2.3.1. Left: energy resolution, right: directional resolution. For both plots, the black line shows the median value; dark blue shaded regions give the 68% range, while light blue shaded regions give the 90% range.

is estimated from the observed ToT values, taking into account the expected relative intensities at given PMT positions. The third algorithm starts from a simple vertex estimation based on large-amplitude hits, followed by a hit selection using this vertex and causality relations and finally by two sequential, independent log-likelihood fits yielding first the vertex position and then the energy and direction of the event. The algorithms yield similar accuracy and are fully efficient for events passing the cuts. While they are less precise than Algorithm 1, they exhibit different responses to non-cascade events, and their output is useful for background suppression. More details on the cascade reconstruction codes presented here can be found in [35].

2.2.6 Prospects for improved reconstruction

The main reconstruction goal of ARCA is to precisely determine the parameters of track-like and cascade-like events, and the methods presented above have been developed with this in mind. New reconstruction algorithms tuned on ν_μ CC and ν_e CC events are in the testing phase and first results are very promising. Also reconstructions tuned for different event classes that present more complex topologies are in the development phase. In particular:

- Improved track and cascade reconstructions

The track and cascade reconstructions described above are first-generation algorithms developed for ARCA, and there are good prospects for improvements in both. In fact, when the reconstruction algorithms were developed the full PMT response was not yet being implemented in the simulation chain. Reconstructions based on a more-detailed knowledge of the detector are currently in development or in the testing phase.

In particular, the best current cascade reconstruction (Algorithm 1) uses very little timing information to fit the cascade energy and direction, and no information from individual PMT signal magnitudes (all time-over-threshold values treated equally). A new cascade reconstruction algorithm that exploits this information in detail is under development. First estimates indicate that a cascade resolutions of 1° may be attainable with improved efficiency.

Additionally, a new track reconstruction algorithm has recently been developed. From initial values obtained by a rigorous scan of the full solid angle, the likelihood is maximised using a multi-dimensional probability distribution function of the arrival time of Cherenkov light from the muon. In Fig. 24, the angular resolution reached for ν_μ CC events is reported, showing that an angular resolution better than 0.1° is reached for events with energy higher than 100 TeV.

However, these reconstructions have not yet been processed through the full Monte Carlo chain described in Sec. 2.2, and hence are not used in the analyses presented here. However, since the




Figure 24: ARCA resolutions for ν_μ CC events using the new track reconstruction algorithm. The black line shows the median value; dark blue shaded regions give the 90% range, while light blue shaded regions give the 68% range. Quality cuts that remove most of the atmospheric muons are applied.

atmospheric background for point-source studies (Sec. 2.3.3) reduces with the square of the angular resolution, using these reconstructions is expected to significantly improve the sensitivity of such studies in the near future.

- τ ‘double-bang’ events

A τ produced in a ν_τ CC interaction will on average travel 4.89 cm/TeV before decaying. If the decay is not into a μ ($\sim 83\%$ probability), the decay products will create a cascade-like signature offset from the first interaction vertex. At sufficiently high energies ($E \gtrsim 100$ TeV), this second cascade will be offset from the first by distances significantly larger than the precision of the cascade reconstruction, creating a ‘double bang’. Identifying such double-bang events would be a clear signature of the flavour of the neutrino primaries.

A preliminary investigation, conducted assuming an initial hadronic cascade (‘bang’) energy of $E_1 = 0.2E_{\nu_\tau}$, an outgoing tau of energy $E_\tau = 0.8E_{\nu_\tau}$, a tau decay length $\ell_\tau = 4.89 \text{ m} \cdot E_\tau / (100 \text{ TeV})$, and a second ‘bang’ energy of $E_2 = 0.67E_\tau$, showed that cascade reconstruction Algorithm 1 could identify both events when separated by 10 m or more, i.e. for ν_τ at ~ 250 TeV and above. It is expected that an even closer separation will be resolvable.

- Starting track events

A ν_μ CC interaction in the detector volume will produce a cascade at the interaction vertex, and an outgoing μ ; ν_τ CC events with subsequent $\tau \rightarrow \mu\nu\nu$ decays will produce a similar signature. Such interactions typically do not manifest themselves as either well-reconstructed cascade or track events, due to the presence of the other component. An optimal reconstruction method would separate both components and reconstruct them simultaneously, allowing for improved energy and direction resolution on the neutrino primary, and a better event selection.

- Muon bundles

Groups of muons from the core of an extended air shower (EAS) exhibit a signature very similar to that of a single high-energy muon in the detector. However, their stochastic energy-loss pattern is much more uniform, and their lateral spread is non-negligible at the characteristic spatial resolution scale of ARCA. Currently, muon bundles are reconstructed using a single-muon hypothesis. Identifying such events can be used to reduce the background for studies searching for an excess of single energetic down-going muons, either from an astrophysical ν_μ flux, or from the prompt decay of charm particles in EAS.




Figure 25: Simulated signature of a self-vetoed event: a ~ 1 PeV atmospheric neutrino (creating the cascade-like signature) accompanied by a muon bundle. Only DOMs with a total ToT (summed over all 31 PMTs) of more than 30 ns in a narrow time window are shown. The colour scale gives the hit times relative to the shower core impacting the sea surface – only some of the muons at sea level (shown as blue lines) penetrated to the detector depth. The size of the circles are proportional to the total ToT on each DOM. DOMs without hits are shown by grey dots.

- Coincident EAS

The rate of downgoing μ from EAS above ARCA that produce a detectable signature in the detector is expected to be about 50 Hz. With a typical event duration of $5 \mu\text{s}$, approximately one in 20,000 events (once every 6–7 minutes) will have a coincidental muon present. Current simulations only model particles for individual EAS, and current reconstruction methods only return at most a single track or cascade event. Observe that this effect is much less important for ARCA than it is for IceCube: the increased detector depth reduces the rate of coincident downgoing muonic background, and the better time-resolution afforded by the low scattering in sea water allows photons from different sources to be separated within a much narrower time window.

2.2.7 Background suppression

Backgrounds from atmospheric muons, as well as random coincidences of hits from ^{40}K decays, are reduced to acceptable levels by applying selection cuts on the event reconstruction quality, the reconstructed zenith angle for track-like events and quantities related to the event energy (such as the number of hits) or event topologies (e.g. using Boosted Decision Tree techniques – see Sec. 2.3). For point-source studies, the main method of reducing the background event rate of both muon and neutrino events is via the excellent angular resolution afforded by seawater, since the background rate reduces with the square of the resolution. However, in particular for studies of a diffuse flux, the most problematic remaining source of background is the atmospheric neutrino flux.

Self-veto of down-going atmospheric neutrinos The interactions of cosmic rays with the atmosphere generates extensive air showers (EAS) where both neutrinos and muons are produced. Despite the ~ 3 km overburden of water, muons with an energy in the TeV range and above can reach the detector, either singly, or in multiples (muon ‘bundles’), particularly under low zenith angles. These muon bundles can be used to ‘veto’ any accompanying neutrinos, allowing for a strong reduction of the down-going atmospheric neutrino background. This technique has been proposed in [27], where it is predicted in the context of an IceCube-like detector that atmospheric neutrinos above 10 TeV and with zenith angles less than 60° can be vetoed with almost 100% efficiency. More detailed calculations in [36] suggest a somewhat lower, but still significant, veto probability.

Events simulated by CORSIKA (see Sec. 2.2) have been used to estimate the self-veto probability, and some preliminary results for the high-energy diffuse analysis using the cascade channel (see Sec. 2.3.1) are




Figure 26: Effect of the self-veto on down-going $\bar{\nu}_e$ events as a function of energy and zenith angle, showing the yearly rate both before (left) and after (right) the self-veto effect is taken into account. Figure taken from [30] (Fig. 4).

shown in [30]. In the case of ARCA, accompanying muons make neutrino-induced cascade events less cascade-like, so that while these events are not explicitly vetoed (as would be the case with IceCube), their topology is such that they appear more background-like than signal-like in sensitivity studies targeting down-going cascade events (see Sec. 2.3). An example of such an event is given in Fig. 25.

An effective ‘veto’ effect can be demonstrated by using the (less sensitive) ‘cut-and-count’ analysis method, as shown in Fig. 26. Shown are the distributions of atmospheric $\bar{\nu}_e$ CC events in the event selection both before (left) and after (right) the self-veto effect has been taken into account. The total effect is a reduction of the down-going atmospheric neutrino events in the selection by a factor of about two, or $\sim 25\%$ in the all-sky background, with higher-energy events close to the zenith being more efficiently rejected. It is difficult, however, to compare this estimate with those of [27] and [36] due to the different event samples, rejection methods, and detector depths. It should also be noted that the current estimate suffers from low statistics, and that the analysis was not optimized with the self-veto effect being taken into account. Hence, the final self-veto efficiency is expected to be higher, and improvements of the results for searches of both diffuse and point-like astrophysical neutrino fluxes in the cascade channel are anticipated.

2.3 Sensitivity studies

In this section, studies of the sensitivity of ARCA to diffuse fluxes and point-like sources are presented. All the analyses take into account (anti-)neutrinos of all flavours (ν_μ , ν_e , and ν_τ) in equal proportions and their CC and NC interactions, as simulated according to Sec. 2.2. Each analysis proceeds in the following steps:

1. A preselection of the events to reject most of the atmospheric background, mostly by cuts on parameters that are provided by the reconstruction algorithms or that are related to the total ToT or number of hits.
2. A multivariate analysis based on the Boosted Decision Tree algorithm (BDT) from the ROOT TMVA package [37], applied to the preselected events for a more stringent background rejection. This step is not applied in all analyses.
3. A ‘cut-and-count’ analysis method for a fast evaluation of the discovery potential and a rough estimate of the number of events from background and signal. This method consists of maximising of the Model Discovery Potential (MDP) (see e.g. the methods of [38,39]) by placing cuts on simulated observables to obtain clean event samples.
4. A maximum likelihood method applied to the event sample resulting from step 2 to calculate the discovery potential at different significance levels. All quoted significances arise from this method – however, since only loose cuts are applied for the likelihood maximisation in order to retain the maximum possible information, the resulting event sample is very broad. Therefore, the expected

numbers of events passing cuts are quoted using the cut-and-count method above, reflecting the number of high-quality signal candidate events.

For the last step, the likelihood ratio function:

$$\text{LR} = \sum_{k=1}^n \log \frac{\frac{n_{\text{sig}}}{n} \cdot P_{\text{sig}}(X_k) + \left(1 - \frac{n_{\text{sig}}}{n}\right) \cdot P_{\text{back}}(X_k)}{P_{\text{back}}(X_k)} \quad (1)$$

is employed, where n_{sig} is the estimated number of signal events, n is the total number of events (and hence, implicitly, $n - n_{\text{sig}} = n_{\text{back}}$ is the number of background events), and P_{sig} and P_{back} are the probability distribution functions (PDF) for signal and background events, respectively. The LR is maximised by altering n_{sig} to obtain LR_{max} . The PDFs are functions of one or more parameters X , such as the BDT output if it is applied, and/or other parameters related to the specific analysis.

Pseudo-experiments are performed and LR is maximised for each pseudo-experiment. The distributions of LR_{max} when simulated signals events are present are compared to distributions in the background-only case to evaluate the significances of each simulated observation.

Unlike in the high-energy starting event (HESE) analysis of IceCube [10], no explicit veto to remove atmospheric muon contaminations is used for KM3NeT/ARCA. Rather, the methods of steps 2 and 4 above assign to each event likelihoods based on the observed event topology, which is well-preserved in sea water due to the low light scattering.

2.3.1 Isotropic diffuse neutrino flux

The detection and detailed investigation of the astrophysical flux observed by IceCube is one of the main physics goals of ARCA during KM3NeT Phase-2.0. In the following, an estimate of the time to detect this flux at the 5σ level is presented.

This study has been optimised assuming that the IceCube signal originates from an isotropic, flavour-symmetric neutrino flux following a power law spectrum with a cut-off at a few PeV. The cutoff – or a steeper spectrum – is implied by the observation of events with a deposited energy exceeding 1 PeV and the absence of events at about 6.3 PeV associated with the Glashow resonance (W production in scattering of $\bar{\nu}_e$ on electrons). The single-flavour energy spectrum has been parameterised as:

$$\Phi(E_\nu) = 1.2 \times 10^{-8} \cdot \left(\frac{E_\nu}{\text{GeV}}\right)^{-2} \cdot \exp\left(-\frac{E_\nu}{3 \text{ PeV}}\right) \text{ GeV}^{-1} \text{ cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}. \quad (2)$$

Since the first IceCube discovery [10], several new analyses with updated event samples and different event selection strategies have been published [13, 18, 40]. In these analyses various compatible parameterisations for the cosmic neutrino flux have been proposed. To check the robustness of our results with respect to the diffuse neutrino flux assumed we have also calculated the significance of the KM3NeT/ARCA observation to the following diffuse flux from [41], which is similar to the results recently reported in [12]:

$$\Phi(E_\nu) = 4.11 \times 10^{-6} \cdot \left(\frac{E_\nu}{\text{GeV}}\right)^{-2.46} \cdot \exp\left(-\frac{E_\nu}{3 \text{ PeV}}\right) \text{ GeV}^{-1} \text{ cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}. \quad (3)$$

Note that for this steeper spectrum, a cut-off to suppress the Glashow resonance signature is not necessarily required by observations, but is kept here in order to avoid biasing the analysis by maximising the selection of such events. In Fig. 27 these fluxes are presented together with the atmospheric neutrino fluxes for comparison.

In the following, the sensitivity studies for diffuse fluxes are presented for the cascade channel and for the track channel.

Cascade channel Events simulated as described in Sec. 2.2 have been reconstructed with the three available cascade reconstruction codes discussed in Sec. 2.2.4.

The first selection cut requires the containment of the reconstructed vertex in a cylindrical volume around the detector centre, with radius $r < 500$ m and height $z < 200$ m. The effect of this cut is illustrated in Fig. 28. It rejects most of the atmospheric muons which, coming from above, have the reconstructed




Figure 27: Diffuse signal fluxes according to Eq. 2 (full blue line) and Eq. 3 (dashed blue line) as a function of the neutrino energy. For comparison the atmospheric neutrino fluxes are also reported (Honda flux [16] for the conventional component and the Enberg flux [17] for the prompt component, see Sec. 2.1.2).




Figure 28: Reconstructed vertex position (r and z are the radial and vertical distances from the detector centre) for atmospheric muon events (left top) and ν_e CC events (right top). The black dashed lines show the instrumented volume of one building block. The red line shows the selected fiducial volume defined by $r < 500$ m and $z < 200$ m. Left bottom: cumulative distribution of ToT_{evt} (see text) for events contained in the fiducial volume, for the different event classes. Right bottom: cumulative distribution of the BDT output ρ for the preselected events (see text) for the different event classes. Vertical lines represent the selection cuts applied.




Figure 29: Ratio of the numbers of selected events and triggered events at each step of the cascade analysis (see text) for atmospheric muons (left) and ν_e CC neutrino reactions (right) as a function of the primary energy.

vertex in the upper part of the detector. The containment cut reduces the fiducial volume by about 20% with respect to the instrumented volume, although this is compensated for by the included region below the instrumented volume.

The corresponding rejection efficiency is reported in Fig. 29 (red line). A further reduction of the background is obtained by removing low-energy events. The event ToT, i.e. $\text{ToT}_{\text{evt}} = \sum_{k=1}^N \text{ToT}_k$ with N being the number of causally connected hits selected by the cascade reconstruction algorithm, is related to the energy deposited in the detector. The cumulative ToT_{evt} distribution is shown in Fig. 28 (bottom left panel). A cut $\text{ToT}_{\text{evt}} > 12 \mu\text{s}$ is applied and rejects most low-energy atmospheric muons and a large part of the atmospheric neutrino background, which is concentrated at lower energies. The corresponding rejection efficiency is reported in Fig. 29 (green points).

As shown in Fig. 29 (left panel) the number of reconstructed atmospheric muons is still too large. To further reduce this background, a BDT algorithm was applied to the preselected event sample. As input for the BDT training, several quality parameters from the available shower and track reconstruction algorithms are used. The BDT is then trained to discriminate tracks from showers using simulated datasets of atmospheric muons and ν_e CC events as training samples. The cumulative distribution of the resulting discrimination parameter ρ and the cut applied on ρ are shown in Fig. 28 (right bottom panel).

A first estimate of the discovery potential, obtained with the cut-and-count approach, yields final event selection cuts of $\rho > 0.5$ and $E_{\text{rec}} > 10^{4.7} \text{ GeV} \approx 50 \text{ TeV}$. The corresponding rejection efficiency is reported in Fig. 29 (blue line). With these cuts the background due to atmospheric muons is almost completely rejected for the presently available simulation livetime of three years at high energies. Tab. 3 reports the number of events per 5 years for ARCA at each step of the analysis for the different event samples. Most of the selected events are ν_e and ν_τ events, due to the higher cascade energy deposition of the CC interactions as compared to the NC channel, and because the BDT identifies ν_μ CC events as less “shower-like” due to the presence of the outgoing muon. With these event rates, a 5σ discovery of the IceCube flux in the cascade channel can be achieved with 50% probability after 1.3 years of ARCA operation. The MC neutrino energy distribution for the final cut-and-count event sample is shown in Fig. 30. The final cuts preferentially select events in the neutrino energy range from about 50 TeV to about 2 PeV ($10^{4.7} \text{ GeV} < E_\nu < 10^{6.3} \text{ GeV}$).

To further improve the evaluation of the discovery potential, the maximum likelihood method (step 4 above) has been applied to the preselected events. The PDF functions in Eq. 1 are functions of the reconstructed energy E_{rec} and the BDT output ρ . The resulting significance is reported in Fig. 34 as a function of the number of observation years. With the KM3NeT/ARCA detector the assumed signal flux will be detectable at 5σ in the cascade channel in about one year of observation time.

The estimate of the significance depends on the assumed background and in particular on the model assumed for the description of the conventional and the prompt components of the atmospheric neutrino flux (see Sec. 2.1.2). For the cascade channel the maximum variation of the significance, reported as a red band in Fig. 34, has been obtained assuming the maximum and minimum flux values of the prompt atmospheric neutrino component reported in [17]. Moreover, the significance has also been estimated taking into account

	reconstruction level	after preselection cuts	after final cuts
μ_{atm}	2.4×10^7	5.5×10^4	6
ν_{atm}^{μ}	1.0×10^5	49	20
ν_{atm}^e	7.1×10^3	23	19
ν_{cosm}^{μ}	352	34	11
ν_{cosm}^e	304	49	41
ν_{cosm}^{τ}	250	34	26

Table 3: Expected number of events for the ARCA detector (2 building blocks) for the different event samples in 5 years of observation time. The cosmic events correspond to the source flux of Eq. 2.




Figure 30: Number of events per year for one building block as a function of the true neutrino energy for events with the MDP cuts. The black vertical lines show the energy range where the 90% of the signal is expected.

the new prompt calculation reported in [19] (see Sec. 2.1.2). In this case, the time to discover the diffuse flux is reduced by about 30%.

Track channel Since energetic muons can have very long tracks (a 10 TeV muon has a path length of $\approx 5\text{--}6$ km in water), muon neutrinos with interaction points far from the instrumented volume can be detected, thus making the effective volume much larger than the geometrical detector volume. The main challenge in using the track channel is to distinguish these events from atmospheric muons. Here we follow the traditional approach to reject atmospheric muons by using the Earth as a shield, and select track-like events that come from below the horizon, or a few degrees above it. In this analysis a cut on the reconstructed zenith angle $\theta_{\text{rec}} > 80^\circ$ is applied.

In Fig. 31 the ratio of the numbers of selected events and triggered events is reported (red lines) for atmospheric muons (left panel) and ν_{μ} CC events (right panel). The atmospheric muon rate at energies below 10^6 GeV is reduced by more than one order of magnitude. Most of the remaining atmospheric muons are mis-reconstructed as up-going or are near the horizon.

To remove the mis-reconstructed events an additional cut on the quality parameter Λ (see Sec. 2.2.4) was applied. In Fig. 32 (left panel) the cumulative Λ distribution is shown for atmospheric muons (black




Figure 31: Ratio of the numbers of selected events and triggered events at each step of the track analysis (see text) for atmospheric muons (left) and ν_μ CC neutrino reactions (right) as a function of the primary energy.




Figure 32: Cumulative distribution of the Λ parameter (left) and of the number of hits, N_{hit} , (right) for events with $\theta_{\text{rec}} > 80^\circ$. The cosmic neutrino signal corresponds to the flux given in Eq. 2. The vertical lines indicate the final cut values applied in the analysis (see text).

	reconstruction level	after preselection cuts	after final cuts
μ_{atm}	2.7×10^8	1.1×10^7	≈ 3
ν_{atm}^{μ}	1.6×10^6	8.0×10^7	18.8
ν_{atm}^e	6.0×10^4	4.9×10^4	0.2
ν_{cosm}^{μ}	1.9×10^3	977	27.9
ν_{cosm}^e	600	381	2.1
ν_{cosm}^T	655	400	2.6

Table 4: Expected numbers of events for the ARCA detector (2 building blocks) for the different event samples in 5 years of observation time. The cosmic events correspond to the source flux of Eq. 2.

line), for atmospheric neutrinos (dashed line) and cosmic neutrinos (solid line), for events with $\theta_{\text{rec}} > 80^\circ$. For $\Lambda \gtrsim -6$ the atmospheric muon background is reduced to the level of the astrophysical neutrino signal.

To reduce the background due to atmospheric neutrinos, a cut on the number of hits associated with the fitted track, N_{hit} , is applied (see Sec. 2.2.4). N_{hit} is related to the muon energy loss in the detector and thus to the primary neutrino energy. The cumulative N_{hit} distribution is presented in Fig. 32 (right panel).

The final cut values, obtained by maximising the MDP for 5 years of observation time, are $\Lambda > -5.8$ and $N_{\text{hit}} > 591$. The resulting numbers of events and the selection efficiencies are reported in Tab. 4 and Fig. 31, respectively. The number of atmospheric muons surviving the final cuts has been extrapolated from the present statistics (see Fig. 16). The MC neutrino energy distribution for the event sample after final cuts is shown in Fig. 33. The cuts select events in the neutrino energy range from about 80 TeV to about 3 PeV. A discovery at 5σ with 50% probability is achieved in about 3.2 years.

As in the cascade analysis, the maximum likelihood method was applied to the preselected event sample ($\theta_{\text{rec}} > 80^\circ$ and $\Lambda > -5.8$) to further improve the sensitivities. The likelihood ratio (Eq. 1) was calculated for signal and background using PDFs that were mono-dimensional functions of N_{hit} . The resulting significance is reported in Fig. 34 as a function of the observation time. The assumed signal flux can be detected with KM3NeT/ARCA at 5σ in the track channel in about 1.6 years of observation time with 50% probability.

For the track channel, the maximum variation of the significance (reported as a grey band in Fig. 34) has been obtained with the assumed uncertainties in the intensity of the conventional atmospheric neutrino flux (see Sec. 2.1.2).

Combined analysis To combine the results of the cascade and track analyses, up-going and down-going events were analysed separately. For down-going events, where the atmospheric background is very high, only contained events are considered (same preselection cuts as in the cascade analysis). For up-going events, preselection cuts on the reconstructed vertices and ToT_{evt} are used to reject atmospheric muons that are wrongly reconstructed as up-going. BDT discrimination cuts are then applied to both samples of preselected events. The BDTs use parameters coming from both the track and shower reconstruction algorithms and are optimised to reject atmospheric muons. The BDT outputs are used together with the cascade energy estimate in a maximum likelihood approach based on Eq. 1. The final result is reported in Fig. 34 (blue line) as a function of the observation time. Combining the results of the track and cascade analyses, KM3NeT/ARCA is expected to observe the IceCube flux (Eq. 2) in about 6 months with a significance of 5σ with 50% probability.

To investigate the sensitivity of these results to the assumed form of the IceCube diffuse flux, both the cascade and track analyses were repeated for signal fluxes according to Eq. 3 both with and without the 3 PeV cutoff. In each case, the flux normalisation constant, $\Phi_{5\sigma}$, required for a 5σ discovery after 1 year of observation time, was calculated. The results are reported in Tab. 5 in terms of their ratio to the flux normalisation reported by IceCube, Φ_{IC}^0 . Values larger (less) than unity indicate a 5σ discovery time of more (less) than 1 year. The results show that for flux assumptions with a softer spectrum and the same cut-off the main results of our analysis do not change, and in fact a small improvement ($\approx 10\%$) is expected.




Figure 33: Number of events passing the MDP cuts (see text), per year for one building block, as a function of the MC neutrino energy. The black vertical lines show the energy range where 90% of the signal is expected.




Figure 34: Significance as a function of the observation time for the detection of a diffuse flux of neutrinos corresponding to the signal reported by IceCube (Eq. 2) for the cascade channel (red line) and muon channel (black line). The black and red bands represent the uncertainties due to the conventional and prompt component of the neutrino atmospheric flux. The blue line represents the results of the combined analysis (see text).

Φ_{IC}^0 [GeV ⁻¹ cm ⁻² s ⁻¹ sr ⁻¹]	$\Phi_{5\sigma}/\Phi_{\text{IC}}^0$	
	Cascades	Tracks
1.2×10^{-8} (Eq. 2)	0.95	1.30
4.11×10^{-6} (Eq. 3)	0.80	1.20
4.11×10^{-6} (Eq. 3 without cutoff)	0.75	0.92

Table 5: Ratios between the flux normalisation needed for a 5σ discovery in KM3NeT/ARCA within 1 year with 50% probability and the different parameterisations of the IceCube flux (see text).

2.3.2 Diffuse neutrino flux from the Galactic plane

One of the most promising potential source regions of a diffuse astrophysical neutrino flux is the Galactic Plane (GP). Neutrinos are expected to be produced in the interactions of the galactic cosmic rays with the interstellar medium and radiation fields, with a potentially significant excess with respect to the expected extragalactic background. The observation of diffuse TeV γ -ray emission from the GP [42, 43], which is expected to arise from the same hadronic processes that would produce high-energy neutrinos, strongly supports this hypothesis. Also Fermi-LAT observes, after the subtraction of known point-like emitting sources, a broad diffuse emission from the GP, with a spectrum consistent with a significant hadronic component [44].

Recently, also related to the observed IceCube high-energy neutrino events, new phenomenological models for the diffuse galactic neutrino emission have been proposed [45–49]. In particular, in [50] a non-uniform cosmic-ray (CR) transport model with a radially dependent diffusion coefficient has been adopted to explain the high-energy diffuse γ -ray emission along the whole GP, as well as the hardening of CR spectra measured by PAMELA and AMS-02 around 250 GeV and two possible CR cut-offs, at 5 PeV and 50 PeV, compatible with KASCADE and KASCADE-Grande observations. In [47], these authors estimate that the astrophysical flux detected by IceCube in both the HESE [11] and diffuse muon [51] analyses is still dominated by an extragalactic diffuse component, with galactic emission respectively accounting for 15% and 10% of events. Using this model, a detailed prediction of the neutrino emission from the inner galactic plane, i.e. for $|l| < 30^\circ$ and $|b| < 4^\circ$ (b and l being the Galactic latitude and longitude, respectively) has been obtained (see Fig. 4 of [50]). This flux is adopted here to estimate the performance of the KM3NeT/ARCA detector in searching for neutrinos from the GP. The selected region is entirely located in the Southern hemisphere. The estimated one-flavour neutrino flux has been parameterised as:

$$\frac{d\phi}{dE_\nu} = 5 \times 10^{-6} \left(\frac{E_\nu}{1 \text{ GeV}} \right)^{-2.3} \cdot \exp \left(-\sqrt{\frac{E_\nu}{1 \text{ PeV}}} \right) \text{ GeV}^{-1} \text{ cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}. \quad (4)$$

An analysis similar to that described for the all-sky diffuse track channel has been performed to estimate the KM3NeT/ARCA sensitivity to this flux. Events were preselected requiring the zenith angle to be $\theta_{\text{rec}} > 80^\circ$ and to point to the sky region with $|l| < 30^\circ$ and $|b| < 4^\circ$. This sky region is visible to the KM3NeT/ARCA detector for events up to 10° above the horizon for about 77% of the time. The final cut values, obtained by minimising the MDP, were $\Lambda > -5.8$ and $N_{\text{hit}} > 181$. The numbers of events from the selected GP region are found to be 2.8 background events (muons and neutrinos) and 3.4 from the source flux of Eq. 4 in one year of ARCA operation. The significance as a function of the observation time has been evaluated by the maximum-likelihood method and is reported in Fig. 35 (left panel). A discovery at 5σ with 50% probability can be achieved in about 5 years of observation time with the KM3NeT/ARCA detector. The discovery flux at 5σ and 3σ is reported as a function of the observation time in Fig. 35 (right panel). As for the diffuse flux analysis, a reduction on the number of the observation years is expected if the cascade channel is included in the analysis. This work is at the moment on-going.




Figure 35: Significance (left) and 5σ and 3σ discovery fluxes (right) as a function of the observation time for the detection in the track channel of a diffuse flux of neutrinos from a selected region of the GP near the Galactic Center (see text).

2.3.3 Point-like neutrino sources

Due its good angular resolution, KM3NeT/ARCA is a very promising instrument for the detection of point-like sources. In particular, its location in the Northern Hemisphere will allow the study of most Galactic sources, as well as extragalactic sources (which are expected to be approximately uniformly distributed over the sky) using up-going muon track events. In this section the sensitivity of the ARCA detector to point-like sources will be discussed. In particular the two following physics cases will be analysed:

- Neutrino emission by the supernova remnant (SNR) RX J1713 and the pulsar wind nebula (PWN) Vela-X, which are at present the Galactic objects exhibiting the most intense high-energy emission [52–54]. For these sources, the zenith position, angular extension, and neutrino flux parameterisation are extracted from the measured high-energy γ -ray spectra. In both cases, the expected neutrino spectra are evaluated from the γ spectrum under the hypothesis of a transparent source and 100% hadronic emission. Although PWN are commonly assumed to be powered by e^-/e^+ winds, they will entrain ions from the ambient medium, possibly accelerating them to very high energies.
- Sources without significant angular extension, emitting a benchmark E^{-2} neutrino spectrum. These can be viewed as characteristic of extragalactic sites of hadronic acceleration (e.g. AGN) with cut-offs expected at very high energies. While the actual spectra of individual neutrino sources is not expected to follow a simple E^{-2} power-law, and may exhibit features such as a peak at PeV energies, or a harder spectra extending to EeV energies [55], the projected sensitivity to an E^{-2} flux gives a good indicator of ARCA's ability to study such extragalactic sources with higher-energy fluxes.

For the detection of neutrinos from point-like sources, the best performance is expected from a search for track-like events. In fact, as discussed in Sec. 2.2.4, with long muon tracks an angular resolution of about $\sim 0.2^\circ$ can be achieved. To remove the unavoidable down-going atmospheric muon background, events are selected that contain tracks reconstructed as up-going.

At the latitude of the Mediterranean Sea, selecting tracks that are reconstructed below or a few degrees above the horizon implies a reduction of the visibility for source declinations above -40° , as shown in Fig. 36. On the other hand, it is possible to view Northern-sky sources below $+50^\circ$ of declination, giving a total of $\approx 3.5\pi$ sr sky coverage.

Galactic sources SNR RX J1713.7-3946 (short: RX J1713) is a young shell-type supernova remnant that has been observed by H.E.S.S. in several campaigns [56, 57]. The γ rays are emitted from a relatively large circular region with a radius of about 0.6° and a complex morphology, with an energy spectrum that extends up to 100 TeV. The source, at a declination of $-39^\circ 46'$, is visible for 80% of the time when selecting tracks with reconstructed zenith angle $\theta_{\text{rec}} > 78^\circ$. For the present analysis, homogeneous emission from a circular




Figure 36: KM3NeT/ARCA visibility as a function of source declination for the muon-track analysis, for 2π downward coverage, i.e. tracks below the horizon (black line); tracks up to 6° above the horizon (blue line); and tracks up to 10° above the horizon (red line).

region around the measured declination with a radial extension of 0.6° has been assumed. The neutrino flux adopted has been derived from the measured γ -ray spectrum and has been parameterized following [58]:

$$\frac{d\phi}{dE_\nu} = 16.8 \times 10^{-15} \left[\frac{E_\nu}{1 \text{ TeV}} \right]^{-1.72} \cdot \exp \left(-\sqrt{\frac{E_\nu}{2.1 \text{ TeV}}} \right) \text{ GeV}^{-1} \text{ cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}. \quad (5)$$

This energy spectrum is shown in Fig. 37 (black line).

In the point source analysis for track-like events, all simulated events (ν_μ , ν_e , ν_τ , μ_{atm}) have been reconstructed with the track reconstruction code described in Sec. 2.2.4.

Since the maximum elevation for a source at the declination of RX J1713 is $\sim 14^\circ$, and to maximise the signal-to-background ratio, events were preselected requiring that the reconstructed track has a zenith angle $\theta_{\text{rec}} > 78^\circ$ and a radial distance from the center of the source of $\alpha < 10^\circ$. The numbers of events at reconstruction level and after the preselection cuts are shown in Tab. 6. Even after the preselection, the numbers of events due to neutrino and muon atmospheric background largely exceed the number of expected signal events from the source. The atmospheric muons can be efficiently removed by imposing a cut on the Λ parameter as shown in Fig. 38. Finally, a BDT trained to discriminate signal events from neutrino background is applied.

The MDP is then maximised by adjusting the cut on the BDT output value. The number of events per 5 years of observation time surviving these cuts is indicated in Tab. 6, together with the number of events expected at each step of this analysis. The ratio between these event numbers and the number of triggered events is reported as a function of the neutrino energy for ν_μ CC interactions in the right panel of Fig. 39.

The significance has been evaluated with an unbinned method [59] by maximising the likelihood ratio of Eq. 1, with PDFs expressed as functions of the BDT output (Fig. 40). The result shows that a 3σ significance can be reached in about 4 years of observation time.

The same analysis has been applied to Vela-X, which is one of the nearest and most intense PWNe (Pulsar Wind Nebulae), and has been extensively studied in TeV γ rays by the H.E.S.S. Collaboration [60,61]. Vela-X is located at a declination of $-45^\circ 36'$. The neutrino spectrum has been estimated from the differential energy spectrum using the prescription in [62–64] for an integration radius of 0.8° around the source center and was parameterized as:

$$\frac{d\phi}{dE_\nu} = 7.2 \times 10^{-15} \cdot \left[\frac{E_\nu}{1 \text{ TeV}} \right]^{-1.36} \cdot \exp \left(-\sqrt{\frac{E_\nu}{7 \text{ TeV}}} \right) \text{ GeV}^{-1} \text{ cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}. \quad (6)$$

This spectrum is shown in Fig. 37 (red line). The source has been simulated as a homogeneously emitting




Figure 37: $\nu_\mu + \bar{\nu}_\mu$ energy spectra for RX J1713 (Eq. 5, black line) and Vela-X (Eq. 6, red line).

	reconstructed tracks	after preselection cuts	after final cuts
μ_{atm}	2.7×10^8	1.5×10^5	≈ 2.0
ν_{atm}^μ	1.6×10^6	1.2×10^4	11.6
ν_{atm}^e	6.0×10^4	545	0
ν_{RXJ}^μ	33.4	23.5	8.1
ν_{RXJ}^e	12.5	0.8	0
ν_{RXJ}^T	12.3	2.55	0.57

Table 6: Expected event numbers in 5 years of observation time for KM3NeT/ARCA (2 blocks) at different stages of the RX J1713 track analysis. The number of surviving atmospheric muons after the final cuts has been extrapolated from the present statistics.




Figure 38: Distributions of the zenith angle θ_{rec} of atmospheric muons at generation level (yellow area), reconstruction level (red line), and after the preselection cut and the cut on Λ (blue histogram).




Figure 39: Ratios between the numbers of selected events and triggered events at each step of the analysis for atmospheric ν_μ CC neutrinos (left panel) and ν_μ CC neutrinos from the source (right panel) as a function of the neutrino energy.




Figure 40: Distributions of the BDT output for background neutrino events and signal events for the SNR RX J1713 analysis.




Figure 41: Significance as a function of the observation time for the detection of the Galactic sources RX J1713 and Vela-X. The bands represent the effect of the uncertainties on the conventional component of the atmospheric neutrino flux.

disk of 0.8° radius.

The expected sensitivity of ARCA to Vela-X is shown in Fig. 41 as a function of the observation time. Owing to the good visibility of the Galactic Plane, a significance of 3σ can be reached in less than 3 years of observation time. The bands show the variation of the significance due to the uncertainty on the normalisation of the conventional part of the atmospheric neutrino spectrum (see Sec. 2.1.2).

Sources with a spectrum $\propto E^{-2}$ The flux required for a 5σ discovery has also been calculated for a generic point-like source with a spectrum $\propto E^{-2}$. In the preselection sample only events with $\theta_{\text{rec}} > 80^\circ$ have been selected. In this analysis, at present, the BDT procedure has not been applied, since the larger difference in the slopes of the atmospheric and source neutrino energy spectra eases discrimination between them.

After the preselection, an unbinned method has been applied that maximises the likelihood ratio of Eq. 1, with PDFs as functions of the two parameters N_{hit} (related to the energy of the neutrinos) and α , the angular distance from the source center. The 5σ discovery flux is reported in Fig. 42 as a function of the declination for 3 years of observation time, corresponding to the exposure for the current IceCube result. The upper limit of ANTARES is also reported for comparison.

ARCA's expected resolution on cascades of $\sim 1.5^\circ$ (see Sec. 2.2.5) allows us to also use this channel for a point-source search, as recently demonstrated by ANTARES [67]. Since discriminating downgoing cascade events from the muonic background is easier than for tracks, cascade searches have a 4π sr coverage, making this detection channel especially important for sources with an otherwise limited visibility. First preliminary results for the cascade channel for generic point-like sources with an E^{-2} spectrum will also be presented in this section.

The sensitivity of KM3NeT/ARCA to point-like sources has been evaluated using cascade events. In this analysis all simulated events have been reconstructed with both the track and cascade reconstructions. To remove the atmospheric muons, which are the main source of background, a preselection of events was performed, leading to the two event samples:

- Sample A: Events reconstructed as down-going with the track reconstruction. Cuts similar to the cascade diffuse analysis have been applied:
 - Geometrical containment cuts $z < 250$ m and $r < 500$ m (see Fig. 28);
 - Reconstructed track zenith $\theta_{\text{rec}} < 80^\circ$;




Figure 42: KM3NeT/ARCA 5σ discovery potential as a function of the source declination (red line) for one neutrino flavour, for point-like sources with a spectrum $\propto E^{-2}$ and 3 years of data-taking. For comparison, the corresponding discovery potential for the IceCube detector [65] (blue line), and upper limits on particular sources for the ANTARES detector [66] (blue squares) are also shown.

- $ToT_{\text{evt}} > 6 \mu\text{s}$ (see Fig. 28);
- $\Lambda < -5.8$.
- Sample B: Events reconstructed as up-going with the track reconstruction. The following cuts are applied:
 - Geometrical containment cuts $z < 324 \text{ m}$ and $r < 450 \text{ m}$ (see Fig. 28);
 - Reconstructed track zenith $\theta_{\text{rec}} > 80^\circ$;
 - $ToT_{\text{evt}} > 4 \mu\text{s}$ (see Fig. 28).

The containment cuts mainly select cascade events that have the interaction vertex inside the detector volume and remove track-like events. Remaining track-like events are rejected by the Λ cut in Sample A (removing well-reconstructed atmospheric muons) and with the ToT cut that removes lower-energy tracks with the vertex inside the instrumented volume. In both samples, most of the selected source events are cascade-like events, the track “contamination” being of order 10%.

Since the cut-and-count method has not been applied in this case (i.e. no cut on the distance between the reconstructed direction and the source center has been applied, with events reconstructed closer to the source appearing more source-like), the unweighted number of signal and background events passing the above cuts is not meaningful, and is not reported.

The same BDT procedure described in Sec. 2.3.1 for the diffuse cascade analysis, to discriminate tracks from showers, has been applied to the two samples. An optimal cut on the BDT output variable was found to be $\rho > 0.5$.

The discovery potential has been obtained by performing an unbinned log-likelihood search. The likelihood takes into account the energy and directional information of each event reconstructed with the cascade reconstruction. In order to take into account the two different event samples, the following likelihood ratio, similar to that one of Eq. 1, has been considered:

$$\text{LR} = \prod_{j=1}^2 \prod_{i=1}^{N_j} \left[\frac{n_{\text{signal}}^j}{N_j} \cdot S_i^j + \left(1 - \frac{n_{\text{signal}}^j}{N_j} \right) \cdot B_i^j \right], \quad (7)$$

where j indicates the data sample and i indicates the event in that sample. S_i^j and B_i^j are the PDFs for the signal and background of the j^{th} sample and are evaluated as functions of the reconstructed cascade energy and of the distance from the source center. N_j is the total number of events in the j^{th} sample. The estimates number of signal events n_{signal}^j in each sample is related to the total number n_{signal} by the relative




Figure 43: Relative fraction of events in Sample A (dark blue) and in Sample B (light blue) as a function of the source declination (left). Angular resolution of events of the preselected samples for a source at declination $\delta = 45^\circ$. The black line represents the median of the distribution. The dark and light blue bands show the 90% and 68% quantiles of the distribution (right).

contribution $n_{\text{signal}}^j = C(\delta) \cdot n_{\text{signal}}$. In Fig. 43 (left panel) the relative percentage of events, $C(\delta)$, of the two selected samples with $\rho > 0.5$ is shown as a function of the declination.

The discovery flux at the 5σ level is reported in Fig. 44 as a function of the declination (red line) for 3 years of observation time, and is compared with the discovery flux obtained for the track analysis. For declinations higher than 50° , where the visibility for up-going tracks is very poor or null, a competitive value w.r.t. the present IceCube value can be obtained (see Fig. 42).

The cascade angular resolution of the preselected events for $\delta = 45^\circ$ is reported in Fig. 43 right panel, and shows that an average angular resolution of about 2° can be reached. This includes all events passing the cuts described above (no cuts on reconstructed angle to the source), showing that a very good angular resolution is obtained.

Similarly to the diffuse analysis, improvements in point-source sensitivity are expected when combining the events from the track and cascade channels, especially for sources located in the Northern sky.

Potential improvements in point-like-source searches An improvement in the sensitivity for the search for neutrinos from point-like sources is expected when the two new reconstruction algorithms (one for track and one for cascade events, see Sec. 2.2.4), that at moment are being tested, are applied to the MC data set. Additionally, the first tests indicate a higher number of reconstructed events (higher efficiency) in addition to a better angular resolution.

The search for neutrino sources can be also improved by grouping potential sources together in a procedure that is known as “source stacking”. This is usually applied to sources of the same class. In our case, several sources otherwise too weak to be investigated individually are present in the Galactic and extragalactic region. This technique has not been yet applied, but an improvement is expected both for the search of Galactic PeV sources (SNR, PWN, etc.) and for extragalactic sources (AGN).

2.3.4 Further physics opportunities

In addition to the central science targets of neutrino astronomy, i.e. investigating high-energy cosmic neutrinos and identifying their astrophysical sources, KM3NeT/ARCA will offer a wide spectrum of further physics opportunities, of which a selection is sketched in the following. Corresponding physics analyses have been pioneered by the IceCube and ANTARES collaborations.

- **Gamma-ray bursts (GRB)**

There is strong evidence that long-duration gamma-ray bursts (GRBs) are produced from relativistic jets formed in the collapse of a massive star [68]. Shocks generated either within the jet, or when the jet collides with surrounding material, are potential cosmic ray acceleration sites, with an associated neutrino flux from subsequent interactions and decays [69]. The short duration of GRBs (seconds




Figure 44: Discovery potential at 5σ with 50% probability for point-sources with an E^{-2} spectrum, for cascade events (red line) for three years of observation time. For comparison the discovery potential for the track analysis is also shown (blue curve).

to minutes) allows a narrow neutrino-search time-window, effectively reducing the background when compared to a standard point-source search. This has allowed ANTARES and IceCube to constrain the properties of GRB jets [70, 71]. KM3NeT/ARCA will increase the sensitivity of such searches similarly to that for E^{-2} point-sources (Sec. 2.3.3).

- **Multimessenger studies:**

KM3NeT/ARCA will be part of a global alert system able to tag synchronous observations of different experiments, observing e.g. γ rays or gravitational waves, that in themselves are not significant but become so when combined. Another branch of multimessenger studies is the creation of alerts for optical, radio or X-ray telescopes to follow up “suspicious” neutrino observations, such as a doublet of events from the same celestial direction during a short time period. As per the ANTARES TAToO program [72], KM3NeT/ARCA will monitor more than half the sky, and will be able to generate alerts with high angular precision within seconds. As ultra-high-energy cosmic rays are also expected to retain some directional information, correlation studies with the arrival directions of events detected by e.g. the Pierre Auger Observatory will also be possible.

- **Cosmic ray physics:**

KM3NeT/ARCA will register a huge number of high-energy atmospheric muons that reflect the direction of impact of the primary cosmic-ray (CR) particle with sub-degree precision. This data set will allow us to investigate inhomogeneities of the CR flux and to complement the corresponding sky maps by IceCube and dedicated CR experiments.

A further opportunity might be the detailed investigation of muon bundles that could, via their multiplicity and divergence, be related to the chemical composition of CRs.

- **Particle physics with atmospheric muons and neutrinos:**

The high-energy end of the atmospheric muon and neutrino spectra are expected to be dominated by prompt processes, i.e. the production of charm or bottom hadrons in the primary CR reactions in the atmosphere and their subsequent fast decay to leptons. Little is experimentally known about these reactions, and theoretical modelling is difficult since it involves QCD processes at the border line of the non-perturbative regime. Identifying and measuring the muons and neutrinos from these processes would shed light on the underlying reaction mechanisms.

- **Tau neutrinos:**

The capability to identify tau neutrino reactions at energies beyond a few 100 TeV (see Sec. 2.2.4) will not only allow for constraining the flavour composition of high-energy cosmic neutrino fluxes, but

might also provide an additional handle to investigate prompt neutrino fluxes (see above), which are the only CR reactions for which a significant production probability for tau neutrinos is expected.

A further interesting phenomenon of tau neutrinos is their regeneration after CC reaction in the Earth through the subsequent tau decay (relevant for energies above a few 10 TeV). The observation of this phenomenon would be interesting in itself, but might in addition signal new particle physics, e.g. in the context of supersymmetry.

- **Dark matter:**

Even though the existence of Dark Matter is considered proven and its particle nature very likely, there is no direct or indirect evidence for the properties of these particles. Should they have masses in the TeV range or above, neutrinos from self-annihilation reactions could be the first Dark Matter signal ever detected. ANTARES and IceCube have already proven the ability of neutrino telescopes to significantly constrain Dark Matter properties, with searches targeting accumulations in the Sun [73, 74], the Galactic Centre [75, 76] and halo [77], and nearby galaxies [78]. The corresponding investigations with KM3NeT/ARCA data will – as with all indirect searches – be particularly sensitive to Dark Matter particles with spin-dependent scattering cross sections on nuclei. The study of neutrino fluxes from the Galactic center and halo, nearby galaxies and galaxy clusters could also provide constraints on scenarios that invoke the decay of very heavy (\sim PeV) dark matter to explain the high-energy neutrino excess observed by IceCube [79–82].

- **Exotics:**

There is a variety of hypothesised stable or quasi-stable particles that would leave an identifiable, characteristic signature when crossing the detector. Amongst these are magnetic monopoles (for which ANTARES and IceCube have already performed a search [83, 84]), strangelets, Q-balls, and nuclearites.

- **Violation of Lorentz invariance:**

Violation of Lorentz invariance (LIV) could lead to oscillation-like interference patterns of atmospheric neutrinos in the energy range of TeV and above. Additionally, LIV would produce a time-delay between neutrinos and photons from distant, time-variable sources (in particular, GRBs), allowing LIV to be tested by multimessenger studies.

2.4 Investigation of systematic effects

The simulation chain described in Sec. 2.2 assumes standard values for the detector geometry, water optical properties, bioluminescent rates, and also a perfectly calibrated detector. In the context of KM3NeT/ARCA sensitivity studies, the term ‘systematic effects’ is used broadly to cover all potential deviations from the standard simulated dataset, and this section describes a series of dedicated studies aiming to estimate their potential influence on ARCA event reconstruction and sensitivity to astrophysical neutrino fluxes.

Each systematic was simulated using a data-set of 10% that of the standard simulation, with the systematic being inserted at the latest possible point in the chain to ensure the least influence of random variation between the sets. For example, changing the water scattering length required re-simulating the hit-time distribution on the PMTs, while reducing the PMT effective area was performed by keeping 90% of the detected photons from the standard simulation. Most systematic effects were simulated as a 10% change, and thus did not reflect the expected size of the resulting effect, but rather were used to estimate the change $dX/d\text{Sys}$ in some relevant quantity X as a function of the systematic Sys.

In each case, the effects of the systematics were first analysed using the ‘golden channel’ approach, i.e. by applying the track reconstruction of Sec. 2.2.4 to ν_μ CC events, and applying the cascade reconstruction Algorithm 1 of Sec. 2.2.4 to ν_e CC events and with the cut-and-count method. Only when a significant effect was found was the systematic applied to the full analysis chains: the point-like track analysis of Sec. 2.3.3, and the diffuse cascade analysis of Sec. 2.3.1.

Similar effects have been considered for KM3NeT/ORCA, particularly in the case of $\vec{\nu}_e$ reconstruction (Sec. 3.4.6). However, the effects of systematics on the mass hierarchy sensitivity of KM3NeT/ORCA are treated via the inclusion of nuisance parameters in the calculation described in Sec. 3.6, rather than using fully resimulated data.

Effect	Tracks		Cascades	
	$\Delta E/E$	$\Delta\theta$	$\Delta E/E$	$\Delta\theta$
$\Delta\lambda_{\text{abs}} = \pm 10\%$	$\pm 8\%$	$\pm 0.1^\circ$	$\pm 30\%$	$< 0.1^\circ$
$\Delta\lambda_{\text{scat}} = \pm 10\%$	$\pm 0.6\%$	$\pm 0.1^\circ$	$< 1\%$	$< 0.1^\circ$
$\Delta A_{\text{eff}} = \pm 10\%$	$\pm 5\%$	$< 0.1^\circ$	$\pm 10\%$	$< 0.1^\circ$
1% missing DOMs		0.01°	$< 0.1\%$	$+0.02^\circ$
1 missing DU		0.01°	$< 0.1\%$	$+0.025^\circ$

Table 7: Estimated effects of systematics on event reconstruction accuracy, evaluated on the event samples from E^{-2} point-source searches for tracks (Sec. 2.3.3) and the diffuse flux search (Sec. 2.3.1). For each sample, the worsening in angular resolution $\Delta\theta$, and percentage change in the mean reconstructed energy $\Delta E/E$, are given for the changes listed in the first column. The magnitude of the effects does not reflect the final expected uncertainty.

2.4.1 Optics: water properties and DOM response

The absorption and scattering of light in seawater has been measured at the KM3NeT-It site to within an accuracy of approximately 10% [85]. The dominant uncertainty is the contribution due to particulates, whereas the scattering and absorption from pure seawater (salt and water) is well-determined. To simulate this effect, the particulate contribution only has been varied so that the scattering/absorption lengths (λ_{scat} and λ_{abs} respectively) vary by $\pm 10\%$ at wavelengths near 400 nm. It is expected that *in-situ* measurements using the KM3NeT calibration system [86] will be able to significantly improve on this knowledge.

The major uncertainty in the response of a DOM to incident photons is the total effective area, A_{eff} , to Cherenkov photons. This is modelled in GEANT simulations with a high degree of accuracy, as described in [31], and has been measured using ^{40}K coincidences *in-situ* with a precision of $\sim 1\%$ [3]. In order to model a significant effect, simulations were produced with A_{eff} varied by $\pm 10\%$ for all photon wavelengths and incident angles.

The effects of these systematic uncertainties on reconstruction accuracy are summarised in Tab. 7, showing the change in reconstruction variables for each percent systematics uncertainty. The most significant effect for the cascade channel is on the energy reconstruction due to a change in λ_{abs} , since in the high-energy regime, only after a large distances do PMTs cease to become saturated, so that the energy reconstruction depends on the response after several absorption lengths. In no case was the direction reconstruction affected, since the Cherenkov peak (which contains most of the directional information) remains unobscured by these effects.

In the case of muon reconstruction, changes in absorption and PMT efficiency have similar effects on energy reconstruction, which is smaller than in the case of cascade reconstruction due to the inherent uncertainties. Systematic effects on direction reconstruction depends on the muon (\sim neutrino) energy. The difference in the track direction w.r.t the standard value is constant above ≈ 1 TeV (values quoted in the table), and increases with decreasing energy below 1 TeV ($\approx 0.1 - 0.2^\circ$ at 100 GeV) as expected, since here the reconstruction is photon-limited. Unlike the case of cascade reconstruction, an increase in water quality, or a larger effective area of the PMTs, improves the directional reconstruction, by increasing the number of Cherenkov photons directly reaching the PMTs.

The effects given in Tab. 7 describe the best estimates of future systematic effects as a function of future uncertainties in the quantities shown. Another relevant measure is: what range of future performances of ARCA is possible given the current uncertainties in these parameters? For this, the systematic effects above were propagated through the simulation chain, allowing reconstructions, cuts, etc. to be re-optimised, i.e. assuming the new value of the changed parameter is known. Effects were analysed in the context of the diffuse flux search using cascades, and the RX J1713 source search using the track channel. Results are given in Tab. 8. Note that for the diffuse analyses, the effects are small, since detection efficiency to both signal and background are affected equally.

Effect	Diffuse (cascades)	Diffuse (tracks)	RX J1731 (tracks)
$\Delta\lambda_{\text{abs}} = -10\%$	3.5%	0%	6.5%
$\Delta\lambda_{\text{scat}} = -10\%$	<1%	1.5%	1.5%
$\Delta A_{\text{eff}} = -10\%$	4%	3%	1.5%
10% less DOMs	1.5%	3.0%	
1 missing DU	0.15%	0.1%	

Table 8: Effects of systematics on the expected sensitivity of the analyses shown, in terms of the change in 5σ discovery flux after one year. E.g., a reduction in λ_{abs} of 10% is expected to increase the one-year 5σ discovery flux of the diffuse cascade analysis by 3.5%.

2.4.2 Detector calibration and alignment

The suite of calibration and alignment systems described in [86] have a finite accuracy, and differences from the true DOM positions and orientations might reduce the precision of reconstruction.

Due to the mechanical structure of KM3NeT detection units, the major degrees of freedom for DOM motion are the position in the horizontal plane, and rotation about the vertical axis. The accuracy of acoustic positioning (position in the horizontal plane) is expected to be 20 cm (corresponding to a hit time uncertainty of about 1 ns in water), while the internal compass for each DOM will measure the rotation angle to within 3° .

To simulate each effect, a false detector was generated with each DOM randomly deviated using Gaussian distributions of width equal to the expected accuracies above, and these were used by reconstruction routines on events generated with the standard simulation chain.

No detectable effects were observed in the accuracy of either the cascade or track reconstruction. This is partially due to the accuracy of the calibrations, partially the uniform coverage of the DOMs (which make errors in the pointing direction less relevant), and partially the robust nature of the reconstruction algorithms themselves. In the case of orientation angle, the uncertainty was artificially increased until, at 9° (three times the expected uncertainty), negligible degradation (too small to be measured) in the angular reconstruction accuracy of track-like events was observed. Hence, no further investigation was undertaken.

2.4.3 Ageing effects

As KM3NeT ages, some loss of performance due to the degradation or loss of key parts is expected. The effects of PMT ageing are covered by the A_{eff} estimate above. An additional simulation was performed to estimate the effect of both lost DOMs and entire DUs, with the standard simulation re-run once with a random 10% sample of DOMs turned off, and once with DUs randomly removed. The effects on both reconstruction and future sensitivity were estimated assuming that the failed units were known, which will be the case due to continual monitoring. The results are shown in Tab. 7 and Tab. 8. In general, the effects are most important for low-energy muon tracks.

2.5 Detector geometry studies

The detector geometry of ARCA was optimised in preliminary studies to target Galactic sources such as RX J1713. While some limits on the final layout are imposed through engineering considerations — in particular, the maximum length of detection units — the horizontal spacing between detection units can be increased or decreased within a relatively broad range. The discovery by IceCube of a diffuse flux extending above 100 TeV [10] now motivates revisiting the question of the optimal horizontal spacing. In particular, a larger spacing would be expected to be more optimal when targeting high-energy events.

In order to characterise the effects of a larger spacing, the analyses described above have also been performed by considering a detector block with 120 m spacing between the detection units, giving an approximate 78% increase in detector volume. The results are tentative since the analyses have not been fully re-optimised to the alternative geometry. Nonetheless, the change in performance gives an indication of the utility of increasing the horizontal spacing.

Analysis channel	Sensitivity change: 90 m → 120 m
Cascades: diffuse (Eq. 2)	+27%
Muons: diffuse (Eq. 2)	+20%
Muons: point-sources (E^{-2} , $\delta = -60^\circ$)	+18%
Muons: RX J1713 (TeV cutoff)	-20%

Table 9: Sensitivity changes in the different analysis channels when increasing the string distance from 90 m to 120 m. The percentages indicate the variation of the fluxes detectable in one year; positive signs indicate a gain in sensitivity with a 120 m spacing. The precise calculation is $\Phi_{90\text{ m}}^{5\sigma}/\Phi_{120\text{ m}}^{5\sigma} - 1$, where $\Phi_{90\text{ m}}^{5\sigma}$ and $\Phi_{120\text{ m}}^{5\sigma}$ are the 5σ discovery fluxes after one year for 90 m and 120 m spacings respectively. Note that all numbers are approximate, being estimated using analyses which have not been fully optimised (see text).

These tentative results are summarised in Tab. 9. They have been performed with 10% of the data set, using the fast cut-and-count method. An improvement of about 20–30% in the discovery flux is observed in the search for a diffuse flux for both channels for the detector layout with increased string spacing. Note that in the cascade channel the sensitivity gain is significantly below the increase of the instrumented volume; one of the reasons is a decrease of the signal detection and reconstruction efficiency with increasing string distances. For Galactic sources, which present a lower neutrino energy spectrum, the change is of the same order, but as expected, in the reversed direction.

It is therefore expected that the final optimum, taking into account all channels and their physics priorities, is in or close to the range explored in this first geometry investigation. Clearly, the choice of optimum configuration depends on the targeted science goals — a larger spacing is better for high-energy diffuse fluxes, and smaller spacings for point-like sources with a low-energy cutoff.

3 Oscillation Research with Cosmics in the Abyss (ORCA)

3.1 Introduction

Important progress has been made in the past two decades on determining the fundamental properties of neutrinos. A variety of experiments using solar, atmospheric, reactor and accelerator neutrinos, spanning energies from a fraction of MeV to tens of GeV, have provided compelling evidence for neutrino oscillations, implying the existence of non-zero neutrino masses (see e.g. [87] and the review by Nakamura and Petcov in [88] for recent insights on the subject).

In the standard 3ν scheme, the mixing of the neutrino flavour eigenstates (ν_e, ν_μ, ν_τ) into the mass eigenstates (ν_1, ν_2, ν_3) is described by the PMNS matrix U which is a product of three rotation matrices related to the mixing angles θ_{12} , θ_{13} and θ_{23} and to the complex CP phase⁵ δ :

$$U = \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{pmatrix} \times \begin{pmatrix} c_{13} & 0 & e^{-i\delta}s_{13} \\ 0 & 1 & 0 \\ -e^{i\delta}s_{13} & 0 & c_{13} \end{pmatrix} \times \begin{pmatrix} c_{12} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (8)$$

where $c_{ij} \equiv \cos \theta_{ij}$ and $s_{ij} \equiv \sin \theta_{ij}$.

Oscillation experiments are not sensitive to the absolute value of neutrino masses but they provide measurements of the squared-mass splittings $\Delta m_{ij}^2 = m_i^2 - m_j^2$ ($i, j = 1, 2, 3$). In the 3ν scheme, there are two independent squared-mass differences; one is responsible for oscillations observed in solar and long-baseline reactor experiments ($\Delta m_{sol}^2 \simeq 7.5 \times 10^{-5} \text{ eV}^2$), while the other impacts the atmospheric neutrino sector ($\Delta m_{atm}^2 \simeq 2.5 \times 10^{-3} \text{ eV}^2$).

At present, the values of all mixing angles and squared-mass differences in the 3ν oscillation scheme can be extracted from global fits of available data with a precision better than 15%, the largest remaining uncertainty being currently on $\sin^2 \theta_{23}$ and its possible octant (i.e. whether θ_{23} is smaller or larger than $\pi/4$) [89–91]. Tab. 10 summarizes the best fit values of the oscillation parameters and the associated 3σ uncertainties as published in [89]⁶.

The recent observation of $\bar{\nu}_e$ disappearance in several short-baseline reactor experiments [92–94] has provided the first high-significance measurement of the mixing angle θ_{13} which drives the $\nu_\mu - \nu_e$ transition amplitude with large mass splitting. The relatively large value of this parameter, $\sin^2(2\theta_{13}) \simeq 0.1$, is an asset for the subsequent searches for the remaining major unknowns in the neutrino sector, and in particular for the determination of the neutrino mass hierarchy (NMH).

The ordering of neutrino mass eigenstates has indeed not been determined so far. After fixing $\Delta m_{21}^2 = (\Delta m^2)_{sol} > 0$, two solutions remain possible depending on the sign of Δm_{31}^2 : the **normal hierarchy** (NH: $m_1 < m_2 < m_3$) and the **inverted hierarchy** (IH: $m_3 < m_1 < m_2$), as can be seen from Fig. 45.

From a theoretical point of view, the determination of the NMH is of fundamental importance to constrain the models that seek to explain the origin of mass in the leptonic sector and the differences in the mass spectrum of charged quarks and leptons [95]. More practically, it has also become a primary experimental goal because the NMH can have a strong impact on the potential performances of next-generation experiments with respect to the determination of other unknown parameters such as the CP phase δ (related to the presence of CP-violating processes in the leptonic sector), the absolute value of the neutrino masses, and their Dirac or Majorana nature (as probed in neutrinoless double beta decay experiments, or $0\nu\beta\beta$). From the astrophysical point of view, the NMH impacts e.g. neutrino flavour conversion in supernovae [96, 97]. Finally, the NMH also affects the precise determination of the PMNS matrix parameters, as can be seen from Tab. 10, which summarizes the current best fit values and their 3σ uncertainties under both hierarchy hypotheses.

While the combination of $0\nu\beta\beta$ and direct neutrino mass experiments with cosmological constraints on $\sum_\nu m_\nu$ might have an indirect sensitivity to the NMH, most of the efforts currently focus on the determination of NMH via neutrino oscillation experiments (see e.g. Sec. 3.1 of [87] for an overview of the subject). One option uses medium-baseline (~ 50 km) reactor experiments such as JUNO and RENO-50, which probe the ν_e oscillation probability at low energies ($\sim \text{MeV}$) where matter effects are negligible [98]. These experiments may be sensitive to the NMH through the interference effects arising from the combination of the fast

⁵We have omitted here the two additional Majorana phases ξ and ζ which are irrelevant in oscillation phenomena.

⁶Updates of global fits presented at conferences, yet unpublished, achieve a precision better than 12% on $\sin^2 \theta_{23}$.




Figure 45: Scheme of the two distinct neutrino mass hierarchies. The color code indicates the fraction of each flavour (e, μ, τ) present in each of the mass eigenstates (ν_1, ν_2, ν_3). From [87].

oscillations driven by Δm_{31}^2 and Δm_{32}^2 . Such a measurement however requires an extreme accuracy both in the energy resolution and in the absolute energy scale calibration.

Another appealing strategy consists in probing the impact of matter effects in both the ν_μ survival probability and in the rate of $\nu_\mu \leftrightarrow \nu_e$ appearance at the atmospheric mass scale. As will be detailed in the next subsection, this option requires long oscillation baselines and matter effects that essentially affect the ν_e -component of the propagation eigenstates, making it possible to determine whether the ν_1 and ν_2 states are lighter or heavier than ν_3 . The ν_e appearance channel is the main focus of current (such as NOvA [99] and T2K [100]) and next-generation (such as CHIPS [101], LBNE [102], LBNO [103] or more recently DUNE [104]) accelerator neutrino experiments. In atmospheric experiments, such as ICAL at INO [105], HyperKamiokande [106], PINGU [107] and ORCA, both channels are important due to the much longer baselines providing stronger matter effects. This strategy has been extensively discussed both for magnetized detectors [108–120] and for water-Cherenkov detectors [114, 121–127], including more recently the specific case of the Mton-scale underice/sea detectors PINGU and ORCA [128–139].

In the 3ν framework, the $\nu_\mu \leftrightarrow \nu_e$ and $\nu_\mu \leftrightarrow \nu_\mu$ transition probabilities in vacuum can be approximated by the following formulae:

$$P_{3\nu}(\nu_\mu \rightarrow \nu_e) \approx \sin^2 \theta_{23} \sin^2 2\theta_{13} \sin^2 \left(\frac{\Delta m_{31}^2 L}{4E_\nu} \right) \quad (9)$$

$$P_{3\nu}(\nu_\mu \rightarrow \nu_\mu) \approx 1 - 4 \cos^2 \theta_{13} \sin^2 \theta_{23} (1 - \cos^2 \theta_{13} \sin^2 \theta_{23}) \sin^2 \left(\frac{\Delta m_{31}^2 L}{4E_\nu} \right) \quad (10)$$

where E_ν is the neutrino energy and L stands for the oscillation baseline. These relations establish the direct link between the transition probabilities and the value of θ_{13} ; they also show that the transitions in vacuum are actually insensitive to the sign of Δm_{31}^2 .

This sign can however be revealed once matter effects come into play along the neutrino propagation path [140, 141]. Contrarily to the other flavours, the ν_e component can indeed undergo charged-current (CC) elastic scattering interactions with the electrons in matter and consequently acquire an effective potential $A = \pm\sqrt{2}G_F N_e$, where N_e is the electron number density of the medium, G_F is the Fermi constant and the $+(-)$ sign is for ν_e ($\bar{\nu}_e$). In the case of neutrinos propagating in a medium with constant density, the transition probabilities now read (adapted from [142]):

$$P_{3\nu}^m(\nu_\mu \rightarrow \nu_e) \approx \sin^2 \theta_{23} \sin^2 2\theta_{13}^m \sin^2 \left(\frac{\Delta m^m L}{4E_\nu} \right) \quad (11)$$

Parameter (hierarchy)	Best fit	3σ range
$\sin^2(\theta_{12})/10^{-1}$ (NH or IH)	3.08	2.59 – 3.59
$\sin^2(\theta_{13})/10^{-2}$ (NH)	2.34	1.76 – 2.95
$\sin^2(\theta_{13})/10^{-2}$ (IH)	2.40	1.78 – 2.98
$\sin^2(\theta_{23})/10^{-1}$ (NH)	4.37	3.74 – 6.26
$\sin^2(\theta_{23})/10^{-1}$ (IH)	4.55	3.80 – 6.41
δ_{cp}/π (NH)	1.39	-
δ_{cp}/π (IH)	1.31	-
$\Delta m_{21}^2/10^{-5} \text{ eV}^2$ (NH or IH)	7.54	6.99 – 8.18
$\Delta m_{\text{large}}^2/10^{-3} \text{ eV}^2$ (NH)	2.43	2.23 – 2.61
$\Delta m_{\text{large}}^2/10^{-3} \text{ eV}^2$ (IH)	2.38	2.19 – 2.56

Table 10: The best fit values and 3σ ranges of the mixing parameters from [89], in the normal (NH) or inverted (IH) hierarchy hypothesis. For the large squared-mass difference the following convention is used: $\Delta m_{\text{large}}^2 = \Delta m_{31}^2 - \Delta m_{21}^2/2$ with $+\Delta m_{\text{large}}^2$ for NH and $-\Delta m_{\text{large}}^2$ for IH.

$$\begin{aligned}
 P_{3\nu}^m(\nu_\mu \rightarrow \nu_\mu) &\approx 1 - \sin^2 2\theta_{23} \cos^2 \theta_{13}^m \sin^2 \left(\frac{(\Delta m_{31}^2 + \Delta^m m^2)L}{8E_\nu} + \frac{AL}{4} \right) \\
 &\quad - \sin^2 2\theta_{23} \sin^2 \theta_{13}^m \sin^2 \left(\frac{(\Delta m_{31}^2 - \Delta^m m^2)L}{8E_\nu} + \frac{AL}{4} \right) \\
 &\quad - \sin^4 \theta_{23} \sin^2 2\theta_{13}^m \sin^2 \left(\frac{\Delta^m m^2 L}{4E_\nu} \right)
 \end{aligned} \tag{12}$$

as a function of the effective neutrino mixing parameters in matter:

$$\sin^2 2\theta_{13}^m \equiv \sin^2 2\theta_{13} \left(\frac{\Delta m_{31}^2}{\Delta^m m^2} \right)^2 \tag{13}$$

$$\Delta^m m^2 \equiv \sqrt{(\Delta m_{31}^2 \cos 2\theta_{13} - 2E_\nu A)^2 + (\Delta m_{31}^2 \sin 2\theta_{13})^2}, \tag{14}$$

where A is positive for neutrinos and negative for antineutrinos. Both the amplitude and the phase of the oscillations can therefore be affected by matter effects. From Eq. 14, the resonance condition is met when the effective mixing is maximal, i.e $\Delta^m m^2$ is minimal. This happens for the case of the NH (IH) in the neutrino (antineutrino) channel at the energy:

$$E_{\text{res}} \equiv \frac{\Delta m_{31}^2 \cos 2\theta_{13}}{2\sqrt{2}G_F N_e} \simeq 7 \text{ GeV} \left(\frac{4.5 \text{ g/cm}^3}{\rho} \right) \left(\frac{\Delta m_{31}^2}{2.4 \times 10^{-3} \text{ eV}^2} \right) \cos 2\theta_{13}. \tag{15}$$

where ρ is the matter density of the medium. For neutrinos passing through the Earth's mantle (core) the resonance will appear around 7 GeV (3 GeV), which explains why atmospheric neutrinos are an appropriate probe for these effects, in association with the large baselines available.

As an illustration, oscillation curves $P(\nu_x \rightarrow \nu_\mu)$ ($x=e,\mu$) obtained with the ORCA software tools (using the PREM model [143] of the Earth density layers) are shown in Fig. 46 for various zenith angles θ (i.e various baselines) as a function of the neutrino energy, both for neutrinos and antineutrinos. In each case, both NMH hypotheses are represented. The strongest impact of the NMH to the oscillation probabilities is in the resonance region $E_\nu \sim (4 - 8) \text{ GeV}$. In the region $\cos \theta \lesssim -0.85$ and $E_\nu < 7 \text{ GeV}$, the effect of the resonant enhancement of the oscillations [108, 121–123, 129, 144–152] for the neutrino trajectories crossing




Figure 46: Oscillation probabilities $\nu_\mu \rightarrow \nu_\mu$ (blue lines) and $\nu_e \rightarrow \nu_\mu$ (red lines) as a function of the neutrino energy for several values of the zenith angle (corresponding to different baselines). The solid (dashed) lines are for NH (IH). For neutrinos (left) and for antineutrinos (right).

the Earth's core can also be seen. Above ~ 15 GeV, the $\nu_e \rightarrow \nu_\mu$ transition probability becomes very small and differences from distinct NMHs tend to disappear as well⁷.

Fig. 46 shows that to first order, the effect for neutrinos in the NH scheme is the same as for antineutrinos in the IH scheme. Nevertheless, and even in the case of non-magnetized detectors (such as ORCA) which do not distinguish ν 's and $\bar{\nu}$'s event-by-event, a net asymmetry in the combined $(\nu + \bar{\nu})$ event rates between NH and IH for a given flavour can be observed. This mainly comes from the fact that in the GeV energy range relevant for atmospheric neutrinos, the CC cross section is different (by about a factor of 2) for neutrinos and antineutrinos, as can be seen from Fig. 47. The relative contribution of ν_e and ν_μ in the steeply falling atmospheric neutrino spectrum, as shown in Fig. 48, also affects the number of events of each flavour that can be expected at the detector level.

Convoluting the oscillation probabilities with the atmospheric neutrino fluxes and the neutrino-nucleon cross section, one can construct bidimensional plots of event rates as a function of the neutrino energy E_ν and cosine of the zenith angle θ . Such an ‘‘oscillogram’’ is represented in Fig. 49 for $\nu_\mu + \bar{\nu}_\mu$, for both NMH hypotheses. Integrating over energies above 4 GeV, one typically expects of the order of 4650 ν_μ -induced events and 2850 ν_e -induced events per year in a 1 Mton detector. The phase space region where the differences between NH and IH are most visible clearly depends on the three ingredients mentioned hereabove; but other factors also come into play, related both to intrinsic effects (such as the physics of the neutrino interaction) and to the detector performance (such as energy and angular resolutions), that will blur the oscillogram patterns and partly wash out the asymmetry effect.

An intrinsic uncertainty in the neutrino energy and direction arises from the kinematics of the neutrino interaction. At the relevant energies, the outgoing lepton can no longer be considered as collinear with its parent neutrino, as can be seen from Fig. 50⁸. This smearing can conveniently be expressed in terms of the

⁷This justifies the approximation of a 2-flavour $\nu_\mu \rightarrow \nu_\tau$ oscillation scheme adopted by high-energy atmospheric neutrino experiments so far [153, 154].

⁸Note that the angle and energy resolutions can be improved by combining information from the leptonic and hadronic parts of the interaction to better reconstruct the kinematics.




Figure 47: Total neutrino (left) and antineutrino (right) CC cross sections per nucleon (for an isoscalar target) divided by neutrino energy and plotted as a function of the energy. Also shown are the various contributing processes: quasielastic scattering (dashed), resonance production (dot-dashed), and deep inelastic scattering (dotted). Taken from [155].




Figure 48: Atmospheric neutrino flux in its different flavour components: absolute values (left) and ratios (right). Taken from [139].

Bjorken inelasticity parameter

$$y = \frac{E_\nu - E_l}{E_\nu} \quad (16)$$

where l stands for a charged lepton and which represents the fraction of energy transferred to the associated hadronic shower. Since the cross section for neutrino and antineutrino behave differently as a function of y , measuring the inelasticity of the neutrino interaction could provide some statistical separation between the ν and $\bar{\nu}$ channels and therefore enhance the sensitivity to the NMH [131]. This effect could be best exploited in the muon channel, where the lepton track and the hadronic shower can in principle be more easily identified than in the other channels; the difference in the muon angular spread for 10 GeV ν_μ and $\bar{\nu}_\mu$ is illustrated in Fig. 51. Preliminary studies performed for ORCA using flavour identification tools are presented in Sec. 3.5 and could be the starting point for a statistical separation between ν_μ 's and $\bar{\nu}_\mu$'s, providing additional enhancement of the sensitivity to NMH in the track and in the shower channel.

The kinematic smearing described here is only one among other sources of systematics directly related to the physical processes at play; fluctuations in the development of the particle cascades, and in the production and propagation of the associated Cherenkov light, must also be taken into account. These effects are discussed in more detail in [156].

Uncertainties in the neutrino oscillation parameters can also degrade the sensitivity to the NMH. These uncertainties are taken into account when evaluating the ORCA sensitivity to the NMH (see Sec. 3.6.1).




Figure 49: Neutrino “oscillograms”: $\nu_\mu + \bar{\nu}_\mu$ event rate (in units of $\text{GeV}^{-1} \cdot \text{y}^{-1} \cdot \text{sr}^{-1}$ in log scale) as a function of the neutrino energy and cosine of the zenith angle, for a 1 Mton target volume. The left (right) plot shows the distribution for the normal (inverted) mass hierarchy.

Other sources of systematics such as the uncertainties on the atmospheric spectra, the uncertainties of the Earth matter density profile, or the unknown δ_{CP} phase are further discussed in Sec. 3.6.4. Detector-related effects, and in particular the energy and angular resolution, are presented along with the description of the event selection and reconstruction performances in Sec. 3.3.

In order to identify, for each flavour, the phase space region where the effects are larger and therefore the discrimination more powerful, asymmetry variables can be defined such as

$$\mathcal{A} = \frac{N_{IH} - N_{NH}}{\sqrt{N_{NH}}} \quad (17)$$

which was used in [128], or

$$\mathcal{A}' = \frac{N_{IH} - N_{NH}}{N_{NH}} \quad (18)$$

where N_{NH} and N_{IH} are the number of expected events at a given angle and energy for NH and IH respectively. \mathcal{A}' essentially reflects the asymmetry of oscillation probabilities and does not depend on the effective mass of the detector, while \mathcal{A} is useful to provide an estimation of the significance of the hierarchy measurement by summing over all oscillogram entries, as proposed by [128]. This approach should however be taken with care as it typically overestimates the discrimination power of the experiment. Alternative approaches discussed in [136, 157–159], and providing a more rigorous statistical treatment, are followed in Sec. 3.6.

An example of asymmetry plots (following the definition of Eq. 18) for ν_μ and ν_e obtained with ORCA software tools and a smearing on energy and angle is shown in Fig. 52. It is clear that the region where the asymmetry is more evident is above 5 GeV. The plots also indicate that comparable levels of asymmetry are reached in both ν_μ and ν_e charged-current interaction channels. Most first-stage studies have concentrated on the ν_μ channel (and on the detection of the associated muon) to determine the sensitivity to NMH, anticipating on the larger statistics in the muon channel and the worse angular resolution of deep sea (/ice) Cherenkov detectors for shower-like events (as produced by ν_e) [107, 127, 128, 130, 132].

In the course of the study it has however been pointed out that this approach may have been too conservative and that the shower channel, and in particular the ν_e -induced events, could also provide a significant contribution to the total sensitivity to NMH⁹. To first order, the atmospheric flux of ν_e of energy E_ν which reach the detector after crossing the Earth along a given trajectory, $\Phi_{\nu_e}(E_\nu, \theta)$ is given by [122, 123]:

$$\Phi_{\nu_e}(E_\nu, \theta) = \Phi_{\nu_e}^0 P(\nu_e \rightarrow \nu_e) + \Phi_{\nu_\mu}^0 P(\nu_\mu \rightarrow \nu_e) \quad (19)$$

$$\simeq \Phi_{\nu_e}^0 [1 + (\sin^2 \theta_{23} r - 1) P_{2\nu}] \quad (20)$$

⁹Experiments like Super-Kamiokande and the proposed Hyper-Kamiokande have indeed mainly focused on the electron neutrino channel, because of the good resolutions they can achieve for this topology in the few GeV energy range [106, 160].




Figure 50: Distribution of the angular difference between the outgoing lepton and the parent neutrino as a function of the neutrino energy, for neutrinos (left column) and antineutrinos (right column) of each flavour. [plot obtained with ORCA tools based on GENIE]




Figure 51: Scattering angle $\phi_{\nu,\mu}$ as a function of Bjorken y for neutrinos (left) and antineutrinos (right) for (anti)neutrino energies of 10 GeV. The distributions are normalized to 1. [plot obtained with ORCA tools based on GENIE]

where $\Phi_{\nu_{e(\mu)}}^0 = \Phi_{\nu_{e(\mu)}}^0(E_\nu, \theta)$ is the $\nu_{e(\mu)}$ flux at the production point in the atmosphere,

$P_{2\nu} = \sin^2 2\theta_{13}^m \sin^2\left(\frac{\Delta m_{31}^2 L}{4E_\nu}\right)$ is the ν_e disappearance probability in a 2ν scheme, and

$$r \equiv r(E_\nu, \theta) \equiv \frac{\Phi_{\nu_\mu}^0(E_\nu, \theta)}{\Phi_{\nu_e}^0(E_\nu, \theta)}. \quad (21)$$

As can be seen from Fig. 48, the ratio r is close to 2 around 2 GeV and below, which tends to suppress the oscillations. This is referred to as the "screening effect" in [128]. However, in the energy range of a




Figure 52: Asymmetry (as defined by Eq. 18) between the number of $\nu + \bar{\nu}$ CC interactions expected in case of NH and IH, expressed as a function of the energy and the cosine of the zenith angle. The right (left) plot applies to muon (electron) neutrinos. A smearing of 25% is applied on the energy. On the angle, a smearing $\sigma_\theta = \sqrt{\frac{m_p}{E_\nu}}$ is applied.

few GeV the ratio increases, which could on the contrary enhance the asymmetry as stated in [129]. The final asymmetry level in the ν_e channel will also depend on the value of the mixing angle θ_{23} ; it could in particular be further enhanced if θ_{23} is found to be in the second octant (i.e. $\theta_{23} > 45^\circ$). The status of electron neutrino studies within ORCA is summarized in Sec. 3.4.

3.2 Simulations

3.2.1 Benchmark detector

The detector geometry used in the Monte Carlo simulations for KM3NeT/ORCA follows the design as described in Sec. 1.2. The simulated ORCA detector corresponds to one building block of 115 string detection units (DUs) with 18 digital optical modules (DOMs) each. The DOMs are made of glass spheres that are designed to resist the hydrostatic pressure of the deep sea environment, each one containing 31 photomultiplier tubes (PMTs) of 3 inch diameter and the related electronics.

The ORCA simulated detector characteristics rely on reasonable assumptions based on the expertise acquired in the KM3NeT collaboration. To reduce the energy threshold, both the vertical and the horizontal spacing must be reduced with respect to the high-energy KM3NeT design (KM3NeT/ARCA). Vertically this can essentially be done at will whereas horizontally there are limitations due to the deformation of the lines by the sea currents and the unfurling procedure of the strings. For a line with 6 m vertical spacing and 18 modules the maximum deviation at the top of the line is about 10 m (corresponding to a sea current of about 30 cm/s). In addition, the accuracy with which a string can be placed on the sea bottom is from ANTARES experience a few meters. A 20 m distance is therefore assumed to be feasible. The collaboration therefore decided to start the simulation study with a detector consisting of 2070 optical modules distributed on 115 DUs placed at a distance of about 20 m from each other (accounting for the positioning uncertainty at deployment), in a circular pattern of radius 106 m (Fig. 53). The DUs host 18 DOMs with 6 m vertical spacing. In this geometry, whose optimum vertical inter-DOM spacing is investigated in this document, the first floor is 50 m distant from the seabed. The detector has a total instrumented volume of about $3.6 \times 10^6 \text{ m}^3$ (equivalent to $\sim 3.7 \text{ Mt}$ for sea water). It is located at the KM3NeT-France site (2450 m depth).

3.2.2 Event generation and characterization

This section describes the software packages used for the generation of Monte Carlo events. Additionally, a selection of event observable distributions is used to characterize their typical fundamental and detector physics phenomenology.

The employed software packages generate atmospheric muons and atmospheric neutrinos. Several

codes have been developed for the KM3NeT project and older codes, that were developed by the ANTARES collaboration, have been modified to take into account the KM3NeT DOM characteristics. The codes simulate the particle interactions with the medium surrounding the detector, light generation and propagation as well as the detector response. In the simulation chain a volume surrounding the instrumented volume, called "can", is defined. The can volume is a cylinder with height and radius exceeding the instrumented volume by about 3 absorption lengths for the atmospheric muon background simulation and by 40 m for the neutrino generation. Generated particles are propagated inside the can and Cherenkov light is generated.

Neutrino and antineutrino induced interactions in sea water in the energy range from 1 to 100 GeV have been generated with a software package based on the widely used GENIE [161,162] neutrino event generator. Electron and muon neutrino events are weighted to reproduce the conventional atmospheric neutrino flux following the Bartol model [163].

All particles emerging from a neutrino interaction vertex are propagated with the GEANT4 based software package KM3SIM [164] that has been developed by the KM3NeT collaboration. It generates Cherenkov light from primary and secondary particles in showers and simulates hits taking into account the light absorption and scattering in water as well as the DOM and PMT characteristics.

The background due to downgoing atmospheric muons is generated with the MUPAGE [25,165] program. MUPAGE provides a parameterised description of the underwater flux of atmospheric muons including also multi-muon events. The parameterised muon flux was obtained starting from full simulations with HEMAS [166] and cosmic ray data. These muons are tracked inside the can with the code KM3 which generates and propagates the light produced by the muons and their secondary particles, taking into account the optical properties of the water. For the photon propagation, the code uses tables containing parameterisations obtained from a full GEANT3 simulation. The code simulates the PMT hit probabilities and the response of the PMTs. The PMT photocathode area, quantum efficiency and angular acceptance, as well as the transmission of light in the optical module glass sphere and in the optical gel are taken into account.

In order to reproduce the randomly distributed background PMT hits due to the Cherenkov light from β -decays of ^{40}K , single photoelectron hits can be added to the hits induced by charged particles inside a chosen time window. Also the hits in coincidence due to ^{40}K between two PMTs inside the same DOM are taken into account.




Figure 53: Footprint of the ORCA benchmark detector (top view), with 115 strings (20 m spacing) with 18 OMs each (6 m spacing). The instrumented volume is $3.6 \times 10^6 \text{ m}^3$ (cylinder: $R=106 \text{ m}$, $z=102 \text{ m}$)

First measurements of the optical background rate indicate a single PMT noise rate of 8 kHz and two-fold coincidence noise of about 340 Hz, for details see [5]. For the simulation results described below a conservative optical background light estimation has been used. An uncorrelated hit rate of 10 kHz per PMT and time-correlated noise on each DOM (500 Hz twofold, 50 Hz threefold, 5 Hz fourfold and 0.5 Hz fivefold) was added. The simulated time-correlated noise rates due to ^{40}K decays have been verified with a complete simulation based on GEANT4.

Fig. 54 shows linearly increasing distributions of the total number of hit PMTs and DOMs as a function of the muon energy for events for which almost all produced light is contained within the instrumented volume. Also shown is a comparison of the simulated light yield for the older KM3 and the more recent KM3SIM code. Both simulations agree quantitatively very well. Roughly 15 detected photons, i.e. PMT hits, and 8 different hit DOMs can be expected per GeV of a contained muon. About 2/3 of the DOMs are hit by scattered light, 1/3 by direct light, while 40% of all hit PMTs register unscattered light.

Fig. 55 compares the number of hit PMTs (DOMs) due to the Cherenkov light emission from a muon, an electromagnetic and a hadronic shower as a function of their respective energy. An electromagnetic shower will cause roughly 12 hits per GeV while a hadronic shower is, as can be expected due to the Cherenkov thresholds of the comparably massive hadrons involved, much dimmer with 7 hits per GeV, i.e. an electromagnetic shower of about 5 GeV energy is almost as bright as a hadronic shower with 10 GeV. The DOM hit multiplicity scales in a similar manner, but somewhat more favorably for hadronic showers due to the on average greater opening angle as compared to electron positron pair cascades.




Figure 54: Median and 15 % / 85 % quantiles (vertical bars) for the distribution of the number of PMTs with hits (left) and DOMs with at least one hit (right) generated by a muon as a function of its energy. Shown are results for two different simulation packages (KM3 (v4r5) and KM3SIM); for the simulation with KM3 the light yield is differentiated into 'direct', 'scattered' and 'all' light.

The inelasticity parameter y of a neutrino interaction on the nucleon critically determines the reaction kinematics as can be seen in Fig. 56 and Fig. 51. At energies below 10 GeV the different strengths of the different interaction channels, quasi-elastic, resonant and deep inelastic, are visible in the y -distributions and result in a higher average inelasticity for neutrinos ($\langle y \rangle \approx 0.5$) compared to antineutrinos ($\langle y \rangle \approx 0.35$). The scattering angle $\phi_{\nu,\mu}$ between the incoming neutrino and the outgoing muon shows a strong dependency and increase with increasing reaction inelasticity. The lower average inelasticity for antineutrinos leads to on average also lower scattering angles. This indicates the discrimination potential of this parameter and the importance to get access through event reconstruction.

Muons from hadronic showers

Employing detailed GEANT3 based simulations, the muon production within the hadronic shower has been studied. A significant contribution of muons with path lengths in excess of the hadronic shower extension, i.e. with energies of at least one or several GeV, could complicate and probably deteriorate the particle flavour identification capabilities (see Sec. 3.5). However, as is shown in [167] and summarized in the following, GeV muons from hadronic showers affect only about 1 % of the events.




Figure 55: Median and 15% / 85% quantiles (vertical bars) for the distribution of the number of PMT (left) and DOM (right) hits generated by a muon, an electromagnetic and a hadronic shower as a function of their respective energy. The KM3SIM package has been used.




Figure 56: Distribution of the interaction inelasticity parameter y as a function of the neutrino energy for neutrinos (left) and antineutrinos (right). Each energy bin is normalized to 1.

In Fig. 57 the Cherenkov photon emission positions along and perpendicular to the hadronic shower direction are shown for simulated shower energies of $E_{\text{had}} \approx 5$ GeV (left) and $E_{\text{had}} \approx 20$ GeV (right). Each Cherenkov photon is weighted with its wavelength dependent detection probability taking into account the PMT quantum efficiencies and the absorption and scattering in sea water. In total 3400 (4000) ν_e CC events with $8 < E_{\nu}/\text{GeV} < 12$ ($30 < E_{\nu}/\text{GeV} < 50$) are used to extract and superimpose their hadronic showers. For both cases only a few muon tracks can be seen to emit light significantly beyond the hadronic shower extension.

Most muons in the hadronic shower come from pion decays. However, pions with energies in the GeV range will likely interact before they decay, as the hadronic interaction length for pions in water is approximately 1 m. In order to study the muon production from charged pions in greater detail, 10^4 charged pions with energies of $E_{\pi} = 2, 5, 10$ GeV have been simulated in sea water. The mean number of muons $\langle N_{\mu} \rangle$ and the fraction of simulated events with at least one muon $N_{\mu \geq 1}$ are summarized in Tab. 11. The energy spectrum and cumulative energy distribution of the most energetic muon is shown in Fig. 58. For all three pion energies the fraction of events producing a muon with more than 1 GeV (2 GeV) is below 2% (1%).




Figure 57: Simulated Cherenkov photon emission positions along and perpendicular to the direction of hadronic showers from ν_e CC events. 3400 superimposed hadronic showers with $E_{\text{had}} \approx 5$ GeV (left). 4000 superimposed hadronic showers with $E_{\text{had}} \approx 20$ GeV (right).

Simulation	$\langle N_\mu \rangle$	$N_{\mu \geq 1}$
π^+ 10 GeV	2.79	0.96
π^+ 5 GeV	1.44	0.84
π^+ 2 GeV	0.93	0.73
π^- 10 GeV	2.22	0.89
π^- 5 GeV	0.91	0.61
π^- 2 GeV	0.42	0.38

Table 11: Mean number of muons $\langle N_\mu \rangle$ and fraction of simulated pions producing at least one muon $N_{\mu \geq 1}$.




Figure 58: Energy spectrum (left) and cumulative energy distribution (right) of the most energetic muon from charged pion π^\pm simulations with energies of $E_\pi = 2, 5, 10$ GeV. In total, 10^4 events for each pion energy are simulated. The peak visible at $E_\mu \approx 250$ MeV is due to the decay of kaons produced at rest in pion induced hadronic interactions.

3.2.3 Implementation of different vertical spacings

The different investigated detector configurations have the same footprint as the benchmark detector (cf. Fig. 53) but different vertical spacings: 6 m, 9 m, 12 m and 15 m.

For the simulation of the neutrino signal the same simulation as for the benchmark detector with 6 m vertical spacing was used and the different vertical spacings are achieved by masking parts of the detector. Doing so the instrumented volume stays the same for all detector configurations, but the DOM density changes. In order to compare the effective volume of the different detector configurations assuming the same number of DOMs for each vertical spacing, the effective volumes of the masked detectors are scaled accordingly (factor of 1/1.5/2/2.5).

Masking

For 6 m vertical spacing all simulated DOMs in the benchmark detector are used. For 9 m vertical spacing every third DOM on each DU is masked. To make the masked detector as homogeneous as possible, the DUs are masked in three different schemes (1st scheme: masking DOM 1, 4, ...; 2nd scheme: masking DOM 2, 5, ...; and 3rd scheme: masking DOM 3, 6, ...). Neighboring DUs use different masking schemes. For the other vertical spacings a similar masking procedure is applied.

It should be noted that the surface to volume ratio for the masked detectors is larger than it would be for a full detector with 18 DOMs per DU. Therefore, the presented results overestimate possible surface-related effects.

3.2.4 Triggering

As described in Sec. 1.6.1, muon and shower events are extracted from the real-time data stream using causality conditions. In the case of ORCA, with a simulated 10 kHz uncorrelated single noise rate per PMT and about 500 Hz time-correlated noise from ^{40}K decays on each DOM, the estimated L1 rate (coincidences on the same DOM in a short time-window) per optical module is about 1.5 kHz.

The trigger algorithms described in Sec. 1.6.1 were optimised for ORCA by considering the effective volume and the event purity. The effective volume is the volume in which a neutrino interaction would trigger the event to be written to disk and the event purity is the fraction of triggered events that contain a neutrino interaction or at least one atmospheric muon. The trigger rate from neutrino interactions is $\mathcal{O}(\text{mHz})$ and is negligible compared to the rate from atmospheric muons ($\mathcal{O}(40 \text{ Hz})$).

The trigger settings correspond to a L1 time window of $\Delta T = 10 \text{ ns}$, a maximum angle between the PMT axes of 90 degrees (L2), and a minimum number of L1 hits of 3 for the shower trigger and 4 for the muon trigger¹⁰. Both triggers run in parallel and one of them or both must fire to flag an event (logical *OR*). For the different considered vertical spacings the distance parameters (R and D) of the muon and shower triggers have been adjusted such that each of the triggers has a rate of $\sim 10 \text{ Hz}$ from pure noise. The rate of atmospheric muon events is evaluated at a depth of 2450 m using the simulations described in Sec. 3.2 and amounts to about 36 Hz (6 m) - 55 Hz (15 m) depending on the vertical spacing of the ORCA detector.

In order to estimate the trigger rates, dedicated simulations for each vertical spacing have been performed, i.e. the detector masking described in Sec. 3.2.3 has not been applied. Trigger rates from pure noise and atmospheric muons are summarised in Tab. 12 for the various vertical spacings. The trigger event purity is 65% - 73%.

It should be noted that during periods of high bioluminescence, the trigger conditions (minimum number of L1 hits and distance parameters R and D) can be tightened in order to reduce the output data rate and match the available data transfer bandwidth.

The effective volume at trigger level for 6 m vertical spacing is shown for different neutrino flavours in Fig. 59 (left) as a function of neutrino energy. Events are weighted to reproduce the conventional atmospheric neutrino flux following the Bartol model [163] and only upgoing neutrinos are considered. The effective volume is smaller for $\bar{\nu}_{\tau}$ NC and $\bar{\nu}_{\tau}$ CC than for $\bar{\nu}_{e,\mu}$ CC events as the outgoing neutrinos are

¹⁰ Muon and shower triggers with larger minimum numbers of L1 hits in conjunction with larger distance parameters R and D have also been studied. However, these triggers show smaller effective volumes than those used in this document.

detector configuration vertical spacing [m]	trigger configuration		trigger rates [Hz]		event purity
	R [m]	D [m]	pure noise	atm. muons	
6	35	40	19	36	0.65
9	39	43	18	41	0.69
12	42	46	19	47	0.71
15	44	50	20	55	0.73

Table 12: Expected trigger rates from pure noise and atmospheric muons for the trigger configurations used for different detector configurations with 6 m, 9 m, 12 m and 15 m vertical spacing.

invisible to the detector. For $\bar{\nu}_{e,\mu}$ CC events the effective volume is larger than for $\nu_{e,\mu}$ CC due to the lower average inelasticity and the resulting higher average light yield (at the considered energies hadronic showers have a smaller average light yield than electromagnetic showers).

The effective volume depends also on the neutrino direction as Fig. 59 (right) shows for ν_e CC events. Other neutrino flavours exhibit a similar zenith angle dependency. For vertical upgoing events ($\cos\theta_\nu \approx 1$) the effective volume rises more steeply with energy than for horizontal events ($\cos\theta_\nu \approx 0$) as more PMTs are oriented downward than upward in an DOM and the density of DOMs is higher in vertical than in horizontal direction.




Figure 59: Effective volume at trigger level (left) for 6 m vertical spacing as a function of neutrino energy for different neutrino flavours (upgoing events only) and effective volume at trigger level for ν_e and $\bar{\nu}_e$ CC events as a function of neutrino energy and cosine of the zenith angle (right).

The effective volumes at trigger level for $\bar{\nu}_e$ CC and $\bar{\nu}_\mu$ CC events for different vertical spacings are shown in Fig. 60 as a function of neutrino energy. For 9 m, 12 m and 15 m vertical spacing the simulation of the benchmark detector with a 6 m spacing is masked and the resulting effective volumes are scaled to the same number of DOMs per DU as described in Sec. 3.2.3. Further details on the triggering studies can be found in [167].

3.3 Muon neutrino studies

This section presents the strategy adopted to reconstruct muon neutrino charged current events with ORCA, and its current performance. All results shown in this section are based on the Monte Carlo simulations presented in the previous sections.

3.3.1 Muon direction reconstruction

The track reconstruction algorithm presented here permits to estimate muon (and consequently neutrino) directions using the combined information of the PMT spatial positions and the Cherenkov photon arrival times. The reconstruction code used is based on the strategy developed for the ANTARES telescope and




Figure 60: Effective volume at trigger level for different vertical spacings (6 m/9 m/12 m/15 m) as a function of neutrino energy for upgoing ν_e and $\bar{\nu}_e$ CC (solid lines) and ν_μ and $\bar{\nu}_\mu$ CC events (dashed lines).

described in reference [168]. This algorithm has been modified to exploit the multi-PMT peculiarities taking into account the directional sensitivity of the KM3NeT optical module.

After an initial hit selection, requiring space-time coincidences between hits, the reconstruction proceeds through four consecutive fitting procedures, each using the result of the previous one as starting point. Each fitting stage improves the result, but the last fit produced, that provides the most accurate result, works well only if the input parameters of the muon track are not too far from the true track parameters. Moreover, the efficiency of the algorithm is improved with a scanning of the entire sky in steps of 3° starting from the prefit track, thus generating 7200 tracks. A scheme of the overall procedure is shown in Fig. 61.




Figure 61: Schematic depiction of the reconstruction algorithm.

As described in Sec. 3.2, the optical background induced by ^{40}K decays has been simulated adding an uncorrelated hit rate of 10 kHz per PMT and a time-correlated hit rate of 500 Hz per DOM (two coincident hits in different PMTs inside the same DOM). To remove the hits from ^{40}K decays, the requirement of




Figure 62: Efficiency and purity of the hit selection adopted by the track reconstruction algorithm as a function of the interacting neutrino energy.

space-time coincidences between hits is used, since hits due to optical background are mostly uncorrelated.

In particular, the hit selection proceeds by first selecting all the local coincidences, i.e. coincidences of hits within the same DOM, in a time window of 10 ns and for which the PMTs involved are less than 90° apart. Among them, a cluster is selected such that any hit in the cluster is causally related to all the remaining ones, according to the following causality relation:

$$|\Delta t| < d/c_{water} + 20\text{ns} \quad (22)$$

where Δt is the time difference between the two hits, d is the distance between the two PMTs and c_{water} is the group velocity of light in water. The cluster of hits obtained is further extended by including the yet unselected hits which fulfill all the following conditions:

- are causally connected to at least 75% of all the hits in the cluster,
- are closer than 50 m to at least 40% of all the hits in the cluster,
- are all causally connected among them.

The latter extension procedure is iterated twice. The resulting performance of the hit selection for ν_{μ} -CC events in terms of efficiency and purity is shown in Fig. 62, where the efficiency is the fraction of signal hits selected among all the signal hits, whereas the purity is the fraction of signal hits among all the selected ones. The resulting set of hits is referred to as *Selected hits* in the following.

The *Selected hits* serve as input of the first step of the track reconstruction procedure, referred to as “linear prefit”, which is a linear fit through the positions of the hits. Once a first estimate of the track is obtained, the evaluation for each hit of the expected angle of incidence θ_i of the photon on the PMT is possible. An “angular selection” is then applied discarding all the hits with $\cos \theta_i > -0.5$. The prefit is then repeated with the new hit set.

Additional starting tracks are obtained by rotating the prefit track by step of 3° over the whole sky. For each starting track, the two fits called *M-estimator fit* and *Pdf fit* are performed. These fits are based on the maximum likelihood method and use probability distribution functions (PDF) that depend on the time residuals, i.e. the difference between the time of the hits and the expected times according to the track hypothesis and Cherenkov light emission.

The *M-estimator fit*, is a maximum likelihood fit based on a function that describes the data for small time residuals. The behaviour of this function for large residuals is a trade-off between a reproduction of the data and the ease of finding the global maximum. The input set of hits, called *M-estim hits* is chosen among the *Selected hits* with conditions on the time residual and the orthogonal distance from the starting track and discarding all hits with $\cos \theta_i > -0.5$. The PDF used for the *Pdf fit* has been parameterised by fitting a set of spectra obtained from Monte Carlo simulations of muons traversing the detector without including

background hits. The input track is the track resulting from the *M-estimator fit* and the input hits are chosen among the *M-estim hits* with conditions on the time residual and the orthogonal distance from the M-estimator track and discarding all hits with $\cos\theta_i > -0.5$. Once the fitting procedures are performed for each starting direction, the solution with the highest likelihood per degree of freedom is chosen as the best one. A further adjustment of the track direction is then achieved with the *Final fit*, using the best track as starting point. This fit relies on the maximum likelihood method and the PDF is obtained taking into account the contributions from both the background hits and the signal hits. The quality of the final reconstructed track is estimated by the quantity:

$$\Lambda = \frac{\mathcal{L}}{N_{\text{hits}} - 5} \quad (23)$$

where N_{hits} is the number of hits used in the final fit and \mathcal{L} is the maximum value of the likelihood.

3.3.2 Neutrino energy estimate

The neutrino energy estimation is performed in two steps: first the muon energy is estimated by reconstructing the muon track length and the interaction vertex, then the neutrino energy is estimated depending on the reconstructed muon length and the number of hits used by the track reconstruction algorithm. These two procedures are described in detail in the following sections.

Reconstruction of the muon track length and the interaction vertex

A dedicated algorithm for the muon energy estimate has been developed relying on the length of the reconstructed muon track. In case of events interacting sufficiently close to the instrumented volume, this algorithm also reconstructs the neutrino interaction vertex as the starting point of the reconstructed muon track.

The estimate of the track length and of the vertex position proceeds through different phases:

1. The detected photons are projected back to the track according to the Cherenkov angle. The first track length estimate, l'_μ is then defined as the distance between the position of the first and last projected photon on the track. The first projected photon is the first vertex estimate \mathbf{V}' . If the muon is generated inside or near the instrumented volume, \mathbf{V}' is an estimate of the interaction vertex, otherwise it indicates the first photon seen by the detector. For these reasons in the following the vertex estimate will be referred to as the "pseudo-vertex" estimate.
2. Some specific features of the hits from the hadronic shower are identified and used to select a set of hits around the first pseudo-vertex estimate.
3. The selected hits are fitted with the hypothesis that they originate isotropically from a single point. This fit gives a second pseudo-vertex estimate \mathbf{V}'' and a second track length estimate l''_μ .
4. The final pseudo-vertex estimate \mathbf{V} is chosen between the first and the second according to the likelihood value of the fit. The corresponding l_μ is kept.

In the following each stage is described in details. The reader only interested in the obtained performances can jump to Sec. 3.3.3.

The procedure to estimate \mathbf{V}' and l'_μ is sketched in Fig. 63. It is assumed that the track direction has already been reconstructed and a subset of hits correlated to the track, called *track-hits*, have been selected. From the position P_i and the time t_i of each hit, the corresponding photon emission point P'_i and the emission time t'_i can be easily calculated. The emission points P'_i are ordered on the basis of their occurrence time t'_i , and the first point P'_0 is the first pseudo-vertex estimate \mathbf{V}' . If P'_n is the last emission point identified, $|P'_n - P'_0|$ corresponds to the first track length estimate l'_μ . Due to the contamination of the optical background and hadronic shower photons, a strict selection is needed to identify the *track-hits*. The following conditions are applied to perform this hit selection:

- A maximum orthogonal distance from the reconstructed track of 50 m;
- A time residual with respect to the reconstructed track in the interval [-10;10] ns;
- $\cos\theta_i < 0$, where θ_i is the expected angle of incidence of the photon on the PMT ($\cos\theta_i = -1$ corresponds to a photon hitting head-on the PMT);




Figure 63: Schematic of the track length estimation procedure. The horizontal arrow indicate the reconstructed muon trajectory, the open circles the PMTs which recorded a hit and the solid dots their projections on the muon trajectory according the Cherenkov hypothesis.

- a minimum density of one point P'_i for each 2 meters along the track segment $\overrightarrow{P'_0 P'_n}$.

The percentage of background hits contained in the set of *track-hits* is below 2%. The set of *track-hits* contains 60–70% of the total amount of hits coming from the track. On the other hand, the contamination due to the hits produced by the hadronic shower increases with the inelasticity y . For low values of y , the largest part of the neutrino energy is transferred to the muon and almost all the selected hits are hits produced by the true muon track. In this case the purity of the *track-hits* reach about 98%. When $y \approx 1$ the hadronic shower takes almost all the neutrino energy and most of the detected hits are due to the shower. Consequently, the purity of the selection decreases to about 20%, the track length is overestimated and the estimated vertex position is some meters away from the real interaction vertex. In such a case, the particles produced at the vertex may even travel backwards with respect to the muon direction. To overcome this problem, a study of the distribution in time and space of hits produced at the interaction vertex has been performed, with the goal of identifying specific features in the reconstruction phase which could be used to distinguish hadronic shower hits among hits due to the optical background and to the muon.

The parameters analysed are the distance d from the estimated pseudo-vertex to the hit position, the transverse and longitudinal projection of d with respect to the reconstructed muon track direction, called k and l respectively. Moreover, the time evolution of the shower hits can also be studied. Under the simplistic assumption that all the hits are emitted from the vertex at a time t_V , a hit with distance d from the vertex should occur at a time $t_V + d/v$, if v is the speed of light in the medium. A “time residual” can be thus defined as $\Delta t = t_i - (t_V + d/v)$, where t_i is the time of the hit. Finally, the conditions applied to select hits from the shower are: $l < 120$ m, $k < 100$ m, $|\Delta t| < 50$ ns, and $(k - l)/k > -2$. The first two conditions are intended to reject the optical background hits and identify a region where the shower is likely to be. The other two are used to distinguish the shower hits from the hits due to the muon track.

The used cuts are chosen in order to distinguish as much as possible shower hits from muon and background hits but trying to keep the few hits that are produced by the shower at low energy. Hits selected in this way are called *shower-hits*. In this hit set the contamination due to the background hits is around 2-3%. The purity of the *shower-hits* increase with the inelasticity reaching about 75% when $y \sim 1$. The set of *shower-hits* contains about 50% of the total number of hits coming from the shower. To find the vertex position a maximum likelihood fit applied to the selected *shower-hits*. A function obtained from the Δt distribution for the simulated shower hits is used as PDF and the final estimate of vertex position is chosen among the first emission point and the result of the fit. Once the vertex has been identified, the track length is scaled according to the distance from the estimated vertex and the last back projected photon on the track. The muon energy is estimated as $E_R = 0.24 I_\mu^{ec}$ GeV, if the estimated track length I_μ^{ec} is expressed in meters.

A selection of the events based on containment conditions is needed for this analysis. Such conditions are based on the results of the reconstruction. In particular, all the events for which the muon reconstructed vertex lies within a volume defined by $|z| < 52$ m and $r < 107$ m, which roughly corresponds to the




Figure 64: Normalised distributions of the time residuals for muon neutrino charged current events with energy higher than 5 GeV and whose vertex is reconstructed within the instrumented volume for 4 Bjorken y intervals, with respect to the reconstructed muon track on the left (track hypothesis) and with respect to the reconstructed vertex on the right (shower hypothesis). For the blue curve, the peak at TimeResidual = 0 is less sharp due to the lower resolution of the vertex reconstruction at low Bjorken y interactions, caused by the lower amount of light emitted at the interaction vertex.

instrumented volume, are selected.

Bjorken y estimation

The Bjorken y is estimated on the basis of the distribution of the time residuals of the *selected hits* (cf. Sec. 3.3.1) with respect to the reconstructed track and with respect to the reconstructed vertex, according to a track and a shower hypothesis respectively (Fig. 64). The simulated angle between the outgoing muon and the outgoing hadronic shower $\phi_{lep, had}$ is shown in Fig. 65 for neutrinos with $E_\nu \approx 10$ GeV. The rationale for this approach relies on the fact that a different repartition of the total neutrino energy among the hadronic shower and the muon influences the distribution of the time residuals. The estimation is performed among four Bjorken y intervals: 0-0.25, 0.25-0.50, 0.50-0.75 and 0.75-1. For each interval, the log-likelihood of the time residuals is calculated for a track hypothesis, if the tested Bjorken y is < 0.5 , or a shower hypothesis, if the tested Bjorken y is > 0.5 . The Bjorken y interval corresponding to the highest likelihood is chosen. The performance of the algorithm is shown in Fig. 66.

Neutrino energy estimation

The estimation of the neutrino energy is obtained by combining the estimated track length and estimated Bjorken y with the number of hits used by the muon track reconstruction. Depending on the reconstructed track length, the true neutrino energy can be related to the number of hits used by the track reconstruction (n_{hits}^{fit}), or equivalently the number of degrees of freedom of the fit $NDoF_{fit} = n_{hits}^{fit} - 5$. The relation between $NDoF_{fit}$ and the energy of the interacting neutrino, for a certain interval of reconstructed muon track length, is obtained by fitting the median distribution of E_ν as a function of $NDoF_{fit}$. In order to further improve the accuracy, two different estimations are used, taking into account the reconstructed Bjorken y being higher or lower than 0.5.




Figure 65: Angle between the outgoing muon and the outgoing hadronic shower $\phi_{\mu,\text{had}}$ from $\nu_{\mu}\text{CC}$ interactions with $9.5 < E_{\nu}/\text{GeV} < 10.5$ as a function of inelasticity y . The kinematics looks very similar for $\nu_e\text{CC}$ interactions. The features at $0 < y < 0.2$ are due to the different neutrino interaction channels (cf. Fig. 56). Each column of bins in y is normalized to 1 (left). Distribution of the opening angle $\phi_{\mu,\text{had}}$ in $\nu_{\mu}\text{CC}$ interactions for three different inelasticities y and neutrino energies (right).




Figure 66: Distribution of the reconstructed Bjorken y (horizontal axis) for four bins of true Bjorken y (colour code). Each histogram is normalised to 1. The plot refers only to muon neutrinos interacting by charged current (no antineutrinos).

3.3.3 Performance

The simulations have been performed with a 3.7Mt detector composed of 115 vertical detection units, horizontally separated by $\sim 20\text{m}$, each housing 18 optical modules, vertically separated by 6m. In order to optimise the vertical separation of the optical modules within a detection unit, the performances of the reconstruction algorithm have been also studied with configurations that mimic vertical spacings of 9m, 12m and 15m by switching off optical modules, cf. Sec. 3.2.3. For the 9m configuration, one optical module out of three is switched off, thus alternating vertical spacings of 6m and 12m, whereas for the 12m configuration one optical module out of two is switched off. The 15m configuration is obtained by alternating vertical spacings of 12m and 18m. The instrumented volume is the same for all the mentioned configurations.

Fig. 67 shows the performances for events reconstructed as upgoing, whose vertex is reconstructed within the instrumented volume, with quality cut of the reconstruction algorithm of $\Lambda > -5.0$ (see Eq. 23). The top left plot shows the median distance between the true and estimated vertex position, $distance(P_{vertex}^{true}, P_{vertex}^{reco})$, as a function of the neutrino energy. The value of the $distance(P_{vertex}^{true}, P_{vertex}^{reco})$ is of the order of a few meters for all reconstructed events. The top right plot in Fig. 67 shows the resolution on the reconstructed neutrino zenith angle and the bottom plot shows the fractional energy resolution, which is defined as $|E_{\nu} - E_{rec}|/E_{\nu}$.




Figure 67: Median resolution as a function of the true neutrino energy, for various vertical spacings, of: the distance between the true interaction vertex and reconstructed one (top left), the absolute value of the difference between the reconstructed zenith angle and the true neutrino zenith (top right), and the fractional energy error (bottom). For muon neutrino and antineutrino events weighted according to the atmospheric spectrum, reconstructed as upgoing, with vertex reconstructed within the instrumented volume and $\Lambda > -5.0$.

Another parameter needed to evaluate the reconstruction performance as well as to calculate the sensitivity for the measurement of the neutrino mass hierarchy is the detector effective volume. The effective volume V_{eff} can be defined as the volume of a 100% efficient detector for observing neutrinos that interact within that volume, for a set of specified quality cuts. In the simulation adopted, described in Sec. 3.2, all the neutrinos interacting within a volume larger than the instrumented volume and surrounding the detector,




Figure 68: Top left: effective mass for ν_μ and $\bar{\nu}_\mu$ as a function of the neutrino energy for events reconstructed as upgoing and whose vertex is reconstructed inside the instrumented volume with a reconstruction quality $\Lambda > -5.0$, for 4 different vertical spacings. Top right: effective masses for 6 m/9 m/12 m/15 m scaled by a factor 1/1.5/2/2.5. Lower plot: ratio between neutrino and antineutrino effective masses as a function of the neutrino energy.

that can be referred to as generation volume V_{gen} , are kept for the subsequent steps of the simulation and, eventually, the reconstruction. The effective volume is then obtained by scaling V_{gen} with the ratio of the reconstructed events N_{rec} (or selected according to a given criterion) and the generated events N_{gen} :

$$V_{\text{eff}}(E_\nu, \theta_\nu) = \frac{N_{\text{rec}}(E_\nu, \theta_\nu)}{N_{\text{gen}}(E_\nu, \theta_\nu)} V_{\text{gen}}. \quad (24)$$

Assuming a seawater density of 1.025 g/cm^3 , the effective volume is converted into an effective mass M_{eff} . The M_{eff} calculated for events with a quality parameter $\Lambda > -5.0$ and whose vertex is reconstructed within the instrumented volume is plotted in Fig. 68 as a function of the neutrino energy and for various intervals of the direction of the incoming neutrino.

3.4 Electron neutrino studies

This section describes the methodology and performance of a reconstruction strategy that has been developed for neutral-current and charged-current shower-like events in ORCA [167]. Electron neutrino events will play a crucial role for the envisaged mass hierarchy measurement, good angular and energy resolutions are therefore mandatory.

3.4.1 Phenomenology of shower events

Charged-current (CC) interactions of electron (anti)neutrinos $\bar{\nu}_e$ with nucleons constitute a very important signal class for the neutrino mass hierarchy measurement. They result in a particle shower:

$$\bar{\nu}_e + N \rightarrow e^\pm + h, \quad (25)$$

where N refers to the target nucleon and h to the hadronic system in the final state. The outgoing electron initiates an *electromagnetic shower* while the hadronic system develops into a *hadronic shower* with a possibly complex structure of hadronic or electromagnetic subshowers, depending on the decay modes of individual particles in the shower.

In the following, the energy E_{had} and momentum \vec{p}_{had} of the hadronic shower are defined by the difference of the respective energy and momentum of the neutrino and the electron:

$$(E_{\text{had}}, \vec{p}_{\text{had}}) = (E_{\nu}, \vec{p}_{\nu}) - (E_e, \vec{p}_e). \quad (26)$$

The inelasticity y ("Bjorken y ") of the reaction is defined as:

$$y = \frac{E_{\text{had}}}{E_{\nu}} = \frac{E_{\nu} - E_e}{E_{\nu}}. \quad (27)$$

In events induced by the neutral-current (NC) weak interaction of a neutrino on a nucleon only a hadronic shower is visible.

Kinematics

The kinematics for ν_e CC interactions is similar to that presented in Fig. 65. The angle is minimal for $y = 0.5$ with a mean value of roughly 25° . For $y \rightarrow 0$ ($y \rightarrow 1$) the angle between the incoming neutrino and the outgoing hadronic shower (lepton) becomes larger, leading to larger $\phi_{lep,\text{had}}$. For increasing neutrino energies the angle $\phi_{lep,\text{had}}$ becomes smaller.

Light production in showers

Some information about the Cherenkov light production of showers can be found in the literature, e.g. in [169] and references therein. Mostly, however, previous studies have focused on energies well above those relevant for ORCA. Therefore, the most important characteristics of showers, as obtained from Monte-Carlo simulation studies, in the relevant energy range for ORCA are briefly summarised in the following.

In general, an electromagnetic shower consists of a cascade of e^\pm emitting photons via bremsstrahlung, which interact with matter and again produce e^\pm -pairs via pair production. The evolution of a hadronic shower is similar but the initial particles are hadrons and the developing cascade will show significantly larger fluctuations as it is dominated by particle decays. In water the electromagnetic and nuclear interaction length are roughly 36 cm and 83 cm [88], respectively. Therefore, compared to muon tracks, showers appear in first approximation as a point-like burst of light in the detector. The light is emitted by charged particles with energies above their Cherenkov threshold.

The longitudinal and transverse light emission profile of electromagnetic and hadronic showers can be seen in Fig. 69. For the energies of interest the brightest point of a shower is offset roughly 1-2 m in the shower direction. The longitudinal extension of the showers increases with $\log(E)$. In spite of the larger interaction length the longitudinal offset for hadronic showers is smaller than for electromagnetic showers with the same shower energy $E_e = E_{\text{had}}$, since they are initiated by several hadrons, each with an energy below E_{had} , and the initial hadrons have different directions reducing the longitudinal extension when projecting onto the shower axis. The transverse extension of the showers is negligible compared to the longitudinal.

Although an electromagnetic shower consists of many e^\pm -pairs with rather short path lengths and overlapping Cherenkov cones, the small pair opening angle preserves the Cherenkov angle peak of the effective angular light distribution which results in a single Cherenkov ring in a projection onto a plane perpendicular to the shower axis. Similarly, each hadronic shower particle with energy above the Cherenkov threshold will produce a Cherenkov ring. Therefore, hadronic showers show a huge variety of different signatures due to the various possible combinations of initial hadron types, their momenta and the diversity of their hadronic interactions in the shower evolution.

Two simulated electron neutrino event examples with $E_{\nu} \approx 10$ GeV and $y \approx 0.5$ each are shown in Fig. 70. The Cherenkov photon ring from the electron is clearly visible together with fainter rings from hadronic shower particles. Due to the large scattering length in water, the angular profile of the emitted light is well conserved over large distances, which leads to the different visible, distinct Cherenkov rings.

While electromagnetic showers show only negligible fluctuations in the number of emitted Cherenkov photons and in the angular light distribution, hadronic showers show significant intrinsic fluctuations in the relevant energy range. These intrinsic fluctuations of hadronic showers and the resulting limitations for the




Figure 69: Light emission profiles of electromagnetic (red) and hadronic showers (blue) with 4 GeV and 40 GeV energy, depicted in shower direction (left, longitudinal) and perpendicular to the shower direction (right, transverse).




Figure 70: Two different simulated ν_e CC events with $E_\nu \approx 10$ GeV and $y \approx 0.5$ in the upper and lower row. Each event is rotated in such a way that the electron is in z -direction. Left: Illustration of the particles produced in the two events. Each arrow represents one particle. The arrow direction and length correspond to the particle momentum in the p_y - p_z -plane, and the arrow color indicates the particle type. Middle and right: Photon distributions in sea water recorded on shells at 20 m and 50 m around the neutrino interaction vertex. Each photon is weighted with the solid angle averaged effective area of a PMT for the photon wavelength. The Cherenkov ring from the electron is centered around $(0, 0)$ with an opening angle of 42° , as the electron moves in z -direction.

energy and angular resolutions have been studied in detail, see [156].

In hadronic showers also muons can be produced via charged pions, which can lead to a wrong flavour classification of the event (see Sec. 3.5). The relevance of muons leaking out of hadronic showers and their energy distribution has been studied in detail, see Sec. 3.2.2.

The averaged angular light distribution for electromagnetic and hadronic showers is shown in Fig. 71 for $E_e = E_{\text{had}} = 5$ GeV. For both shower types the probability to detect at least one photon within one DOM

(DOM-hit probability) is maximal at the Cherenkov angle of 42° , but it is more peaked for electromagnetic than for hadronic showers. At smaller distances the Cherenkov peak becomes washed out due to the extension of the shower in conjunction with the small lever arm for the definition of the angle with respect to the shower direction. Note that the light distribution for a single hadronic shower event will not be as smooth as shown in these plots due to the distinct Cherenkov rings from each hadron.




Figure 71: DOM-hit probability (probability to detect at least one photon in an entire multi-PMT optical module) at a distance of 20 m (left) and 50 m (right) away from the brightest point for showers with $E_e = E_{\text{had}} = 5\text{GeV}$ as a function of the angle θ between shower direction and the vector from the brightest point to the DOM centre.

Sensitivity to the reaction inelasticity y

Electromagnetic and hadronic showers induced by neutrino interactions in the energy range relevant for the neutrino mass hierarchy measurement show slightly different light emission characteristics in the detector. Due to the large scattering length in water these differences are conserved over sufficiently large distances, so that information from a large detector volume can contribute to the discrimination between both shower types. In electron neutrino charged-current events, in which both an electromagnetic and a hadronic shower are present at the same time and partly overlapping, the angular separation $\phi_{e,\text{had}}$ of both showers can help to distinguish between them. This can make an estimation of the reaction inelasticity y in $\bar{\nu}_e\text{CC}$ events feasible. Additionally, it might allow for a partial separation of $\bar{\nu}_e\text{CC}$ and NC events on a statistical basis.

However, with an ORCA-like detector¹¹ it seems impossible to distinguish a shower induced by a single electron from a shower induced by a single hadron, since both resulting Cherenkov light cones will be of the same intensity for the same particle energy. Fig. 70 (bottom) shows a simulated example event, in which the electron ($E_e = 4.77\text{ GeV}$) and the pion ($E_\pi = 3.71\text{ GeV}$) induce Cherenkov rings of similar intensity.

The most intense Cherenkov ring in $\bar{\nu}_e\text{CC}$ events is seen in most cases from the electron, as can be inferred from the distribution of the inelasticity parameter y in Sec. 3.2 and keeping in mind that the hadronic shower energy E_{had} is often shared between many hadrons. A measure for the intensity of a Cherenkov ring E_x^{cher} induced by a particle x with energy E_x can be defined by:

$$E_x^{\text{cher}} = \begin{cases} E_x - m_p, & \text{if particle } x \text{ is a baryon} \\ E_x, & \text{else} \end{cases} \quad (28)$$

where m_p is the proton mass.

In the example event in Fig. 70 (bottom), the most intense Cherenkov ring is from the electron ($E_e^{\text{cher}} = E_e$) and the relative intensity is $E_e/E_\nu = 4.77\text{ GeV}/9.82\text{ GeV} = 0.49$, while the leading Cherenkov ring in the hadronic shower $E_{\text{had}}^{\text{cher}}$ is from the pion ($E_\pi^{\text{cher}} = E_\pi$) with a relative intensity of $E_\pi/E_\nu =$

¹¹Detector with a spacing between optical sensors of several metres up to few tens of metres.

$3.71 \text{ GeV}/9.82 \text{ GeV} = 0.38$. The distribution of E_e/E_ν , the leading $E_{\text{had}}^{\text{cher}}/E_\nu$ in the hadronic shower and the leading $E_{\text{tot}}^{\text{cher}}/E_\nu$ of the total event is shown in Fig. 72 for $\bar{\nu}_e$ CC events with $9 \text{ GeV} < E_\nu < 11 \text{ GeV}$. Additionally, the distribution of leading $E_{\text{had}}^{\text{cher}}/E_\nu = E_{\text{tot}}^{\text{cher}}/E_\nu$ in $\bar{\nu}_e$ NC events with $9 \text{ GeV} < E_{\text{had}} < 11 \text{ GeV}$ is shown.

Generically, the measurable inelasticity y is given by $1 - E_{\text{tot}}^{\text{cher}}/E_\nu$. Therefore, it is expected that all measured events will show an inelasticity of $y \lesssim 0.8$, as $E_{\text{tot}}^{\text{cher}}/E_\nu \gtrsim 0.2$ for all neutrino interaction types (cf. Fig. 72). This is even the case for NC events, which are in principle very similar¹² to $\bar{\nu}_e$ CC events with the same neutrino energy as the hadronic energy in the NC events and an inelasticity of $y = 1$.




Figure 72: Distribution of E_e/E_ν (red), the leading $E_{\text{had}}^{\text{cher}}/E_\nu$ in the hadronic shower (blue) and the leading $E_{\text{tot}}^{\text{cher}}/E_\nu$ of the total event (back). Top left: ν_e CC events with $9 \text{ GeV} < E_\nu < 11 \text{ GeV}$. Top right: $\bar{\nu}_e$ CC events with $9 \text{ GeV} < E_\nu < 11 \text{ GeV}$. Bottom: ν_e and $\bar{\nu}_e$ NC events with $9 \text{ GeV} < E_{\text{had}} < 11 \text{ GeV}$.

3.4.2 Shower reconstruction algorithm

A neutrino induced shower-like event is characterised by 8 free parameters: vertex position \vec{x}_{tx} and time t_{tx} , energy E , direction \hat{e}_s and inelasticity y . The shower direction is characterised by 2 angles.

The shower reconstruction is performed in two steps. In the first step the vertex is reconstructed based on the recorded time of the PMT signals, commonly called hits, and in the second step the direction, energy and inelasticity are reconstructed based on the number of hits and their distribution in the detector. In both steps a maximum likelihood fit is performed for many different starting shower hypotheses and the solution with the best likelihood is chosen.

This factorization of the fitting procedure works well due to the homogeneity of water and its large scattering length which allows for a precise vertex reconstruction independent of the shower direction.

¹²Small differences due to different characteristics of hadronic showers induced by W or Z bosons.

Vertex reconstruction

The majority of the Cherenkov light from electromagnetic and hadronic showers is emitted within a few metres around the neutrino interaction vertex, cf. Sec. 3.4.1. Therefore, direct hits from a shower are characterised by a small time residual t_{res} :

$$t_{\text{res}} = t_{\text{hit}} - t_{\text{vtx}} - d/c_{\text{water}}, \quad (29)$$

where d is the distance between the vertex position \vec{x}_{vtx} and the PMT position, t_{vtx} is the vertex time and c_{water} is the speed of light in water. The vertex position and time are here defined as the brightest point and its corresponding time in the shower evolution and not by the neutrino interaction itself, because the brightest point is what is actually seen by the detector.

The vertex reconstruction is performed in two successive maximum likelihood fits. For both fits, the likelihood for the vertex hypothesis $(t_{\text{vtx}}, \vec{x}_{\text{vtx}})$ is given by:

$$L = \sum_{\text{hits}} g(t_{\text{res}}|t_{\text{vtx}}, \vec{x}_{\text{vtx}}), \quad (30)$$

where $g(t_{\text{res}}|t_{\text{vtx}}, \vec{x}_{\text{vtx}})$ is a function of the hit time residuals for a given shower hypothesis.

The first vertex fit (prefit) is designed to be very robust against noise hits and an imprecise initial vertex hypotheses. The initial hit selection is optimised for low energetic shower-like events and is described below together with the choice of the initial vertex hypothesis. In the prefit, the following function g is used:

$$g(t_{\text{res}}) = 1/\sqrt{4 + (t_{\text{res}}/\text{ns})^2} \quad (31)$$

Based on the initial hit selection in total 15 starting vertex hypotheses for the prefit are generated. The fitted vertex with the best likelihood is chosen as result of the prefit.

The second vertex fit is more precise but needs a hit selection with higher signal purity and a good starting vertex hypothesis. The result of the prefit is used to generate in total 10 starting vertex hypotheses (result of the prefit and 9 vertex hypotheses around it with time shifts of ± 25 ns and position shifts of 5 m in a random direction). A rather pure signal hit selection is achieved by selecting hits according to the following criteria:

- $10 \text{ m} < d < 80 \text{ m}$
- $-50 \text{ ns} < t_{\text{res}} < 50 \text{ ns}$
- $-1 < \cos(\psi) < 0.1$

where ψ is the angle between the PMT direction (vector normal to the photocathode plane) and the vector from the vertex to the PMT, i.e. only PMTs which are orientated towards the vertex and can be hit by unscattered photons are taken into account. In this fit, for function g a distance d dependent t_{res} distribution, obtained from simulated $\bar{\nu}_e \text{CC}$ events, is used and shown for three distances in Fig. 73. With increasing distances the peak of direct hits becomes broader due to scattering and dispersion, and the hit probability decreases due to absorption leading to a relative increase of the noise level.

The fitted vertex with the best likelihood and within 10 m and 50 ns around the result of the prefit is chosen as final vertex.

Initial hit selection for first vertex fit

For the initial selection of shower-like hits the following hit patterns are defined:

- L1** coincidence between hit times of 2 PMTs on the same DOM in a time window $\Delta t \leq 10$ ns
- L2** L1 with an angle between the hit PMTs smaller than 90° , note that these are the same definition as used in the triggers, cf. Sec. 3.2.4
- L3** coincidence between hits on 3 PMTs on the same DOM in a time window $\Delta t \leq 10$ ns
- V2L2** coincidence between two L2 hits on different DOMs which are closer than 35 m and within a time window $\Delta t \leq 10 \text{ ns} + t_D$, where t_D is the time required by the light to travel the distance D between the two DOMs




Figure 73: Time residual distribution for three different distances d between the vertex and the PMT, obtained from simulations of fully contained ν_e CC and $\bar{\nu}_e$ CC events with 5 kHz noise rate. The time residual is defined with respect to the brightest point of the shower. The distributions are normalized so that the maximum is 1.

T0L0 coincidence between two hits on adjacent or next-to-adjacent DOMs on the same string in a time window $\Delta t \leq 10, \text{ns} + t_D$

The general strategy is to find first a *reference hit* that is very likely a signal hit and close to the neutrino interaction vertex. The position/time of this reference hit is then used as an initial vertex hypothesis to select additional hits based on their time residual and further requirements to suppress noise hits.

Firstly, the largest cluster of causally connected L2 hits is selected by requiring $\Delta t \leq D/c_{\text{water}} + 10 \text{ ns}$ for all L1 hits within the cluster. From these causally connected L2 hits the subset of hits that additionally satisfy the L3 or V2L2 criteria is selected. These L3 or V2L2 hits are ranked according to their hit multiplicity (number of coincidences on the same DOM) as well as the number and multiplicity of causally connected hits in the vicinity of 25 m. The most signal-like hit is chosen as 'reference hit'.

Secondly, all hits around the reference hit are selected that are closer than 100 m, within a time window of $-250 \text{ ns} < t_{\text{res}} < 10 \text{ ns}$ and causally connected with most L3 or V2L2 hits. The loose lower time cut allows for distances up to about 50 m between the true neutrino interaction vertex and the reference hit, e.g. because the neutrino interaction is outside the detector volume. The drawback of this relatively large time window is a contamination with noise hits. Therefore, hits are discarded that do not satisfy the L1 criterion, or are either causally connected with an adjacent L3 or V2L2 hit on the same string or fulfill the T0L0 criterion in addition to being causally connected with a L3 or V2L2 hit in the vicinity of 25 m. The hits selected by this procedure are used in the first vertex fit and the position/time of the 15 most signal-like hits are used as initial vertex hypotheses.

Reconstruction of energy, direction and inelasticity

Once the shower vertex is fixed, the remaining parameters which can be fitted are the shower energy E , direction \hat{e}_s and the reaction inelasticity y . In principle all of these parameters can be inferred from the angular light distribution (cf. Fig. 71): the shape is sensitive to the inelasticity y , the integral is in first order proportional to the energy (as the light yield is in first order proportional to the shower energy) and the direction in which this angular light emission profile is present gives the shower direction.

In the following, the shower energy E , direction \hat{e}_s and inelasticity y are reconstructed using a maximum likelihood fit based on the probability that the hit pattern is created by a trial shower hypothesis $\vec{\alpha} = (t_{\text{vtx}}, \vec{x}_{\text{vtx}}, E, y, \hat{e}_s)$.

As discussed in Sec. 3.4.1, the electron mostly is the dominant particle in ν_e CC events and produces the

brightest Cherenkov ring. Therefore, the reconstruction is designed to find the electron direction \hat{e}_e and not the neutrino direction.

The final hit selection, the definition of the likelihood function and the fitting procedure are described in the following.

Final hit selection

Based on the result of the vertex fit, hits are selected according to the following criteria:

- $10 \text{ m} < d < 80 \text{ m}$
- $-25 \text{ ns} < t_{\text{res}} < 25 \text{ ns}$
- $-1 < \psi < 0.1$

For simplification¹³, all PMT-hits on the same DOM are merged and the times of the individual hits are not taken into account, so that the event is quantified by $N_{\text{hits}}^{\text{DOM}}$ for each DOM. For the fit all DOMs with $10 \text{ m} < d < 80 \text{ m}$ are taken into account, that includes also the DOMs without any selected hit.

Likelihood

Ignoring shower-to-shower fluctuations, the probability $P(N_{\text{hits}}^{\text{DOM}})$ to detect $N_{\text{hits}}^{\text{DOM}}$ on a given DOM depends on: E , y , distance d between the vertex and the DOM, angle θ between shower direction \hat{e}_s and the vector \vec{d} from the vertex to the DOM, and the DOM orientation. The DOM orientation can be described by a single angle β between \vec{d} and the DOM direction, because the angular acceptance of the entire DOM (sum of all PMT angular acceptances) shows in first order a rotational symmetry due to the multi-PMT structure, cf. Sec. 1. All of these quantities are illustrated in Fig. 74.




Figure 74: Illustration of the quantities relevant for the probability $P(N_{\text{hits}}^{\text{DOM}})$.

The likelihood is computed as follows:

$$L = \prod_{\text{selected DOMs}} P(N_{\text{hits}}^{\text{DOM}}(E, y, d, \theta, \beta) | \vec{\alpha}) . \quad (32)$$

To define the probability $P(N_{\text{hits}}^{\text{DOM}})$ two auxiliary quantities are introduced: the number of expected photons $\langle N_\gamma \rangle$ and the variance $\text{var}(\langle N_\gamma \rangle)$ of the $\langle N_\gamma \rangle$ distribution. To take fluctuations in the hadronic shower into account, the variance of the expected number of photons has been introduced.

The dependency on the DOM orientation and the distance are parameterized. For the number of expected photons the dependency on the DOM orientation is assumed to follow the angular acceptance of the entire DOM. For the final hit selection, the attenuation of $\langle N_\gamma \rangle$ with distance is well described by $\exp(-d/\lambda_{\text{att}}(d))$.

¹³Besides reducing the computation time for the fit, this simplification is justified by the fact that each DOM in principle measures the intensity of the shower event at a given position. As the PMTs on the same DOM are nearly at the same position and direct light from a shower arrives at the DOM nearly at the same time, the information of the individual PMTs — which PMT is hit at which time — is not needed. Of course, the multi-PMT structure is needed to estimate the shower intensity from the number of hit PMTs $N_{\text{hits}}^{\text{DOM}}$.

d^{-2} , where the first term describes the effective attenuation due to absorption and scattering and the latter term describes the geometrical reduction of solid angle coverage of the DOM on a sphere with radius d . The effective attenuation length has been derived from a fit to the MC simulations and can be parameterized as $\lambda_{\text{att}}(d) = a + b \cdot d$ with $a = 32.2$ m and $b = 0.034$.

Taking these parameterizations into account, $\langle N_\gamma \rangle$ and $\text{var}(\langle N_\gamma \rangle)$ depend on (E, y, θ, d) . Although the d dependency is already taken into account via the parameterization above, the shape of the θ distribution changes with distance (see Fig. 71) so that a coarse binning in d is needed.

The probability density function $P(N_{\text{hits}}^{\text{DOM}})$ depends on $(N_{\text{hits}}^{\text{DOM}}, \langle N_\gamma \rangle, \text{var}(\langle N_\gamma \rangle), \beta)$. The quantities $\langle N_\gamma \rangle$, $\text{var}(\langle N_\gamma \rangle)$ and the probability $P(N_{\text{hits}}^{\text{DOM}})$ are obtained from MC simulations of $\bar{\nu}_e$ CC events.

An example distribution of the expected number of photons $\langle N_\gamma \rangle$ as a function of the angle θ for different inelasticity y intervals is shown in Fig. 75. As the angle θ is defined with respect to the electron direction, a clear Cherenkov peak of the electron at 42° is visible. With higher inelasticity y this peak becomes fainter due to less energetic electrons, while the number of expected photons in the 'off-peak region' ($\theta \gtrsim 60^\circ$) increases due to the more energetic hadronic showers. Therefore, these PDF tables gain sensitivity to the reaction inelasticity y from the ratio of the peak to the off-peak region.




Figure 75: Number of expected photons $\langle N_\gamma \rangle$ as a function of the angle θ between the shower direction (which is the electron direction) and the vector from the vertex to the DOM for different inelasticity y intervals, and for shower energies of $8 \text{ GeV} < E < 9 \text{ GeV}$ and at distances of $40 \text{ m} < d < 50 \text{ m}$.

Fitting procedure

For technical reasons, each event is fitted with 9 different fixed inelasticity y assumptions¹⁴ ($y = [0 - 0.1, 0.1 - 0.2, \dots, 0.7 - 0.8, 0.8 - 1]$). For each fixed y the likelihood maximisation is performed for 5 different starting shower hypotheses. The initial shower hypothesis is calculated from the selected hits. The direction is estimated by the sum of all vectors from the vertex to the DOM weighted by $N_{\text{hits}}^{\text{DOM}}$ and the energy is estimated empirically by $(\sum N_{\text{hits}}^{\text{DOM}} - 20)/4$. The other four seeds are randomly chosen perpendicular to the first starting shower hypotheses with the same energy.

Finally, the result with the best likelihood of all 45 fits is selected. Thus, the final result has a discrete value for the reconstructed inelasticity y .

3.4.3 Event selection

The final event selection criteria are:

¹⁴The last inelasticity y bin (0.8-1) is chosen larger than the other bins, as the MC statistics decreases for $y \rightarrow 1$.

- $E_{\text{reco}} > 1 \text{ GeV}$
- result of both vertex fits is similar in space and time: distance $< 4 \text{ m}$ and time difference $< 20 \text{ ns}$
- a minimum of 7 (3) out of the 15 (10) reconstructed vertices from different seeds in the first (second) vertex fit are similar to the selected vertex of this fit: distance $< 2 \text{ m}$ and time difference $< 10 \text{ ns}$
- $\text{cov}_{20^\circ} \geq 0.4$,
- $\text{cov}_{45^\circ} \geq 0.4$, $\text{cov}_{60^\circ} \geq 0.4$ and $\text{cov}_{75^\circ} \geq 0.4$

The final hit selection (including also hits with $d < 10 \text{ m}$) must again also fulfill the shower trigger, cf. Sec. 3.2.4. The *coverage* cov_x is defined as the fraction of directions on a cone with the opening angle x around the reconstructed direction at the reconstructed vertex position that satisfy the following containment condition: $L_{\text{inVol}} > 20 \text{ m}$, where L_{inVol} is the path length inside the instrumented volume for distances away from the vertex between 10 m and 70 m .

This coverage cut is introduced to ensure that a reasonable fraction of the expected hit pattern from the reconstructed shower is contained in the instrumented volume. Therefore the coverage cut is in principle a containment cut for the reconstructed vertex depending on the reconstructed shower direction.

3.4.4 Reconstruction performance

The performance of the shower reconstruction is studied on MC simulations described in Sec. 3.2 and events are selected according to the criteria described in Sec. 3.4.3. The vertical spacing between the DOMs is 6 m if not stated otherwise. For all following results the events are weighted to reproduce the conventional atmospheric neutrino flux following the Bartol model [163].

Performance for charged-current electron neutrino events

Effective volume: The effective volume for upgoing $\nu_e \text{CC}$ and $\bar{\nu}_e \text{CC}$ events is shown in Fig. 76 as a function of neutrino energy for different neutrino zenith angle ranges. Depending on the zenith angle the plateau reaches 3.8 Mm^3 (horizontal), 3.6 Mm^3 (vertical upgoing) and around 3.7 Mm^3 for all upgoing ν_e and $\bar{\nu}_e$. The turn-on is slightly steeper for vertical upgoing than for horizontal events as more PMTs are oriented downward than upward in a DOM. 90% of the plateau is reached around $E_\nu = 8 \text{ GeV}$ (7 GeV) for ν_e ($\bar{\nu}_e$). The turn-on is slightly steeper for $\bar{\nu}_e$ than for ν_e due to the lower average inelasticity.




Figure 76: Effective volumes in Mm^3 (10^6 m^3) as a function of neutrino energy for different true neutrino $\cos(\text{zenith})$ ranges, where $\cos(\text{zenith})=1$ means vertical upgoing and $\cos(\text{zenith})=0$ means horizontal. The solid black line corresponds to upgoing neutrinos weighted according to the Bartol atmospheric neutrino flux model. Left: $\nu_e \text{CC}$. Right: $\bar{\nu}_e \text{CC}$.




Figure 77: Longitudinal and perpendicular distance between the neutrino interaction position and the reconstructed vertex position for all selected events with $E_\nu = 2 - 30$ GeV (left). Longitudinal distance for $6 \text{ GeV} < E_\nu < 7 \text{ GeV}$ and $0.2 < y < 0.4$ fitted with a Gaussian (middle). Perpendicular distance for the same E_ν and y range fitted with ‘distance \times a Gaussian’ (right).

Vertex resolution: The distance between the neutrino interaction position and the reconstructed vertex position is shown Fig. 77 (left) for all selected events in the energy range of $E_\nu = 2 - 30$ GeV.

The distance is split into a longitudinal (in neutrino direction) and a component perpendicular to neutrino direction. An offset in neutrino direction is clearly visible and expected, since the brightest point of the shower is reconstructed and not the interaction vertex position. As discussed in Sec. 3.4.1, the brightest point of the shower is offset by $0.5 \text{ m} - 2 \text{ m}$ in the relevant energy range (cf. Fig. 69). The longitudinal and perpendicular distances are fitted with Gaussian functions for different neutrino energy and inelasticity bins. Exemplary, the distributions and the Gaussian fits are shown for $6 \text{ GeV} < E_\nu < 7 \text{ GeV}$ and $0.2 < y < 0.4$ in Fig. 77 (middle and right).

The mean of the Gaussian fit to the distribution of the longitudinal distances corresponds to the shift between the brightest point and the neutrino interaction. The vertex resolution corresponds to the resolution on the brightest point and is given by the fitted widths. The longitudinal and perpendicular width can be combined into a 3-dimensional resolution on the vertex by $\sigma_{3D} = \sqrt{\sigma_{\text{long}}^2 + \sigma_{\text{perp.}}^2}$. The combined vertex resolution is about $0.5 \text{ m} - 1 \text{ m}$ and is dominated by the longitudinal vertex resolution. This precise vertex reconstruction justifies the factorization of the shower reconstruction into a vertex reconstruction and a shower energy, direction and inelasticity reconstruction.

The fitted mean longitudinal vertex shift (in meter) is shown in Fig. 78 as a function of E_ν and Bjorken y . The increasing distance of the reconstructed shower brightest point from the interaction vertex with increasing neutrino energy is clearly visible.




Figure 78: Mean value in meter (from Gaussian fits) of the longitudinal distance between the neutrino interaction vertex and the reconstructed brightest point as a function of E_ν and y .

Direction resolution: The median neutrino direction resolution (the angle between reconstructed direction and neutrino direction) as a function of neutrino energy is shown in Fig. 79 for different neutrino zenith angle ranges and for ν_e and $\bar{\nu}_e$ separately. For events weighted with the Bartol flux model the median directional resolution is better than 10° for energies above 8.5 GeV for ν_e CC and above 5.5 GeV for $\bar{\nu}_e$ CC events. The resolution is slightly better for vertical upgoing than for horizontal neutrinos as more PMTs are oriented downward than upward in a DOM.

As the reconstruction is designed to find the electron direction, the resolution is better for $\bar{\nu}_e$ than for ν_e due to the smaller average inelasticity for $\bar{\nu}_e$ leading on average to a smaller intrinsic scattering angle between the neutrino and the electron. The median intrinsic scattering angle, the median resolution with respect to the electron direction and the neutrino direction as a function of neutrino energy are shown in Fig. 80. For the relevant energy range the median electron direction resolution is smaller than the intrinsic scattering angle and the median neutrino direction resolution, verifying that the reconstruction actually has the ability to find the electron in $\bar{\nu}_e$ CC events.

Fig. 81 shows the median electron direction resolution as a function of electron energy for different true inelasticity y ranges. The reconstruction of the electron direction is only slightly affected by the additional light from the hadronic shower up to $y \approx 0.5$. For $y \gtrsim 0.6$ the reconstruction can additionally be confused by high energetic particles in the hadronic shower producing a brighter Cherenkov ring than that from the electron. Due to momentum conservation the most energetic particles produced in neutrino interactions tend to have smaller scattering angles with respect to the neutrino direction. Therefore, by sometimes reconstructing the dominant particle from the hadronic shower the median neutrino direction for ν_e CC events is slightly better than the intrinsic scattering angle between neutrino and electron for neutrino energies above ~ 5 GeV, as can be seen from Fig. 80.




Figure 79: Median neutrino direction resolution (angle between reconstructed direction and neutrino direction) as a function of neutrino energy for different true neutrino $\cos(\text{zenith})$ ranges, where $\cos(\text{zenith})=1$ means vertical upgoing and $\cos(\text{zenith})=0$ means horizontal. The black line corresponds to upgoing neutrinos weighted according to the Bartol flux model. ν_e CC (left) and $\bar{\nu}_e$ CC (right).

Inelasticity resolution: The resolution on the inelasticity y for a low, medium and high y range is shown in Fig. 82 (left) for $6 \text{ GeV} < E_\nu < 12 \text{ GeV}$. The distributions of the reconstructed inelasticity y_{reco} and true inelasticity y_{true} for ν_e CC and $\bar{\nu}_e$ CC events are shown in Fig. 82 (right) for $6 \text{ GeV} < E_\nu < 12 \text{ GeV}$. The absence of $y_{\text{reco}} > 0.8$ can be explained by dominant particles in the hadronic shower mimicking a lower inelasticity as discussed in Sec. 3.4.1. The accumulation of events at low y_{reco} is larger than expected from the MC inelasticity distribution and visible for ν_e CC and $\bar{\nu}_e$ CC events. This is a feature of the reconstruction algorithm.

Due to the sensitivity to y the y_{reco} distribution is different for ν_e and $\bar{\nu}_e$ CC events leading to a separation power between both channels. This sensitivity to y can also be used to separate $\bar{\nu}_e$ CC events from $\bar{\nu}_e$ NC events.




Figure 80: Median intrinsic scattering angle (red crosses), median electron direction resolution (blue diamonds) and the median neutrino direction resolution (black filled circles) as a function of neutrino energy for upgoing ν_e CC (solid marker) and $\bar{\nu}_e$ CC (hollow marker) events weighted according to the Bartol flux model.




Figure 81: Median electron direction resolution (angle between reconstructed direction and electron direction) as a function of electron energy for upgoing ν_e and $\bar{\nu}_e$ CC events weighted according to the Bartol flux model. Different marker colors and styles represent different true inelasticity y ranges.

Energy resolution: In Fig. 83 (left) the reconstructed energy is shown as a function of the neutrino energy for $\bar{\nu}_e$ CC events weighted according to the Bartol flux model. The reconstructed energy is systematically higher than the neutrino energy. Therefore, an energy correction depending on the reconstructed zenith angle θ_{reco} , inelasticity y_{reco} and reconstructed energy E_{reco} is applied. The corrected reconstructed energy $E_{\text{reco}}^{\text{corr}}$ is given by

$$E_{\text{reco}}^{\text{corr}} = f(y_{\text{reco}}, \theta_{\text{reco}}, E_{\text{reco}}) \cdot E_{\text{reco}}, \quad (33)$$

where the 3-dimensional correction function $f(y_{\text{reco}}, \theta_{\text{reco}}, E_{\text{reco}})$ has been calculated from MC such that the median reconstructed energy is equal to the neutrino energy assuming a Bartol flux model. The corrected reconstructed energy as a function of the neutrino energy is shown in Fig. 83 (right).

The difference between reconstructed and neutrino energy in different neutrino energy bins is shown in Fig. 84 for ν_e CC and $\bar{\nu}_e$ CC events separately. These distributions are very well described by Gaussians.




Figure 82: Distribution of reconstructed inelasticity y_{reco} for three different true y ranges ($0 < y < 0.2$, $0.4 < y < 0.6$ and $0.8 < y < 1$) for ν_e CC and $\bar{\nu}_e$ CC events with $6 \text{ GeV} < E_\nu < 12 \text{ GeV}$ weighted according to the Bartol flux model (left). Distributions of the reconstructed inelasticity y_{reco} (solid lines) and true inelasticity y_{true} (dashed lines) for ν_e CC and $\bar{\nu}_e$ CC events (right).




Figure 83: Reconstructed energy as a function of true neutrino energy for ν_e CC and $\bar{\nu}_e$ CC events weighted according to the Bartol flux model (left). Corrected reconstructed energy as a function of true neutrino energy for the same events (right).

The median fractional energy resolution – given as $|E_{\text{reco}} - E_\nu|/E_\nu$ – is better than 18% for neutrino energies above 5 GeV for upgoing ν_e CC and $\bar{\nu}_e$ CC events and is shown as a function of neutrino energy in Fig. 85. The relative energy resolution – given as the RMS of $(E_{\text{reco}} - E_\nu)$ distributions (cf. Fig. 84) over neutrino energy – is better than 26% (24%) for neutrino energies above 7 GeV for upgoing ν_e ($\bar{\nu}_e$) CC events and is shown as a function of visible energy E_{vis} in Fig. 86 (left) together with the resolution for the other shower-like neutrino interaction channels. For $\bar{\nu}_e$ CC events the visible energy is equal to the neutrino energy. The resolution is better for $\bar{\nu}_e$ CC events than for ν_e CC events due to the lower average contribution from the hadronic shower which shows larger fluctuations than electromagnetic showers [156]. Fig. 86 (right) shows the mean relative offset between the mean reconstructed energy and the visible energy. At energies corresponding to the effective volume turn-on region the reconstructed energy is overestimated for $\bar{\nu}_e$ CC and ν_e CC events as only events pass the event selection criteria that appear more energetic than they actually are. Above $\sim 9 \text{ GeV}$ the reconstructed energies are slightly overestimated (underestimated) for $\bar{\nu}_e$ (ν_e) CC due to the smaller light yield of hadronic showers compared to electromagnetic showers.

Performance for shower-like neutrino events

The performance of the shower reconstruction is evaluated separately on different shower-like neutrino interaction event samples: $\nu_{e/\mu}$ NC and $\bar{\nu}_{e/\mu}$ NC events (consisting of $\nu_{e,\mu}$ and $\bar{\nu}_{e,\mu}$, respectively), ν_τ CC and $\bar{\nu}_\tau$ CC events where the τ lepton decays in an electron and ν_τ CC and $\bar{\nu}_\tau$ CC events where the τ lepton decays




Figure 84: Difference between corrected reconstructed energy and neutrino energy in different neutrino energy bins for ν_e CC (red) and $\bar{\nu}_e$ CC (blue) events weighted according to the Bartol flux model. Dashed lines show Gaussian fits.




Figure 85: Median fractional energy resolution ($|E_{\text{reco}} - E_{\nu}|/E_{\nu}$) as a function of neutrino energy for ν_e CC (red) and $\bar{\nu}_e$ CC (blue) events weighted according to the Bartol flux model.

in hadrons.

The effective volume for upgoing shower-like neutrino events is shown in Fig. 87 (left) as a function of neutrino energy. The turn-on is much less steep for $\bar{\nu}_e$ NC and $\bar{\nu}_e$ CC events than for $\bar{\nu}_e$ CC events, as the outgoing neutrinos are invisible to the detector. For $\bar{\nu}_e$ CC events the turn-on is steeper than for $\bar{\nu}_e$ NC events as on average the visible energy in $\bar{\nu}_e$ CC events is larger than in $\bar{\nu}_e$ NC events. In ν NC events the average inelasticity is higher than in $\bar{\nu}$ NC events leading to more energetic hadronic showers and a steeper

turn-on.

The median directional resolution is shown in Fig. 87 (right) as a function of neutrino energy. The directional resolution for $\bar{\nu}_{\tau}\text{NC}$ and $\bar{\nu}_{\tau}\text{CC}$ events is clearly worse than for $\bar{\nu}_{e}\text{CC}$ events as the information of the outgoing neutrinos is unavailable. As the angle between the hadronic shower and the neutrino is smaller for νNC than for $\bar{\nu}\text{NC}$ events due to a higher average inelasticity, the directional resolution is better.

The relative energy resolution – given as RMS over visible energy E_{vis} – for upgoing shower-like neutrino events is shown as a function of E_{vis} in Fig. 86 (left). E_{vis} is defined as the difference between the energy of the incoming neutrino and the outgoing neutrino(s) from the primary neutrino interaction (NC events) or τ -decay ($\bar{\nu}_{\tau}\text{CC}$ events). The resolution is worse for events with higher average contribution from hadronic showers which show larger fluctuations [156].

Due to the smaller light yield of hadronic showers compared to electromagnetic showers, the ratio $\langle E_{\text{reco}} \rangle / E_{\text{vis}}$ is different for each neutrino interaction channel and energy dependent. This can be seen in Fig. 86 (right). The higher the fraction of electromagnetic shower component in the event the higher is the mean reconstructed energy. This leads also to different turn-on behaviours in the effective volume for both shower types, and consequently to different compositions (in terms of electromagnetic and hadronic shower components) of well reconstructed neutrino events. The latter explains the behaviour below $E_{\text{vis}} \lesssim 10$ GeV.

The distribution of the reconstructed inelasticity y_{reco} for $\bar{\nu}_{\tau}\text{NC}$ events with hadronic shower energies of $6 \text{ GeV} < E_{\text{had}} < 12 \text{ GeV}$ is shown in Fig. 88. As expected, the y_{reco} distribution for NC events looks similar to the distribution for $\bar{\nu}_{e}\text{CC}$ events with $0.8 < y < 1$ (cf. Fig. 82), but different to the other y ranges, leading to a separation power between shower-like events from $\bar{\nu}_{e}\text{CC}$ and $\bar{\nu}_{\tau}\text{NC}$ events.




Figure 86: Relative energy resolution $\text{RMS}/E_{\text{vis}}$ as a function of the visible energy E_{vis} for shower-like neutrino interaction channels (left) and mean relative offset in reconstructed energy – given as $\langle E_{\text{reco}} \rangle / E_{\text{vis}}$ – as a function of E_{vis} (right).




Figure 87: Effective volumes (left) and median neutrino direction resolution (right) as a function of neutrino energy for different upgoing shower-like neutrino event types.




Figure 88: Distribution of reconstructed inelasticity y_{reco} for NC events with hadronic shower energies of $6 \text{ GeV} < E_{\text{had}} < 12 \text{ GeV}$ and an arbitrary true inelasticity.

3.4.5 Performance for different vertical spacings

The different detector configurations studied in this section are described in Sec. 3.2.3. The performance for different vertical spacings is studied on upgoing $\bar{\nu}_e$ CC events weighted according to the Bartol atmospheric neutrino flux model. Events are selected according to the same criteria as described in Sec. 3.4.4. For each detector configuration the respective energy correction is applied (cf. Sec. 3.4.4).




Figure 89: Effective volumes for the different vertical spacings (top left). Effective volumes for 6m/9m/12m/15m scaled by a factor 1/1.5/2/2.5 (top right). Ratio of neutrino and antineutrino effective volumes for charged-current electron neutrino events for the different vertical spacings (bottom).

The effective volumes for the masked detectors with different vertical spacings are shown in Fig. 89

(top left). For all detector configurations a similar plateau value is reached, but the turn-on is less steep for smaller DOM density (larger vertical spacing). Assuming the same number of DOMs for each vertical spacing, these effective volumes can be scaled accordingly as shown in Fig. 89 (top right). The ratio of effective volumes for ν_e CC and $\bar{\nu}_e$ CC is shown in Fig. 89 (bottom).




Figure 90: Resolution of the shower reconstruction for different vertical spacings for upgoing ν_e CC and $\bar{\nu}_e$ CC events as a function of neutrino energy. Top left: median neutrino direction. Top right: median fractional energy resolution ($|E_{\text{reco}} - E_\nu|/E_\nu$). Bottom left: relative energy resolution RMS/E_ν . Bottom right: mean relative offset in reconstructed energy $\langle E_{\text{reco}} \rangle / E_\nu$.

In Fig. 90 the resolutions for the different vertical spacings are summarized. The resolution on the neutrino direction, and the energy deteriorates for larger vertical DOM spacings. The performance for other shower-like neutrino events for different vertical spacings is similar as described previously.

3.4.6 Effect of variation in water/PMT properties and noise level on reconstruction performance

The reconstruction performances have been studied for a variation in water properties, PMT quantum efficiencies (QE) and optical background noise. For this purpose, the absorption and scattering lengths λ_{abs} and λ_{scat} have been changed by $\pm 10\%$, while the QE has been changed by -10% – a fuller discussion of these parameters is given in Sec. 2.4. To test the influence of the optical background, the single noise rate is increased from an already conservative 10 kHz to 20 kHz in the whole detector. Bioluminescence does not produce correlated noise apart from random coincidences and can be simulated by increasing single noise rates.

Effect of known parameter variations It is assumed that the true water, PMT and noise properties are known so that they can be accounted for in the reconstruction. The trigger conditions are unchanged

compared to the nominal values¹⁵ and events are selected according to the same criteria as for the nominal values. This study has been performed for the detector with 6 m vertical spacing — similar effects are expected for larger spacings.

The energy and direction resolution for a known variation in water, PMT and noise properties is shown in Fig. 91 and Fig. 92 together with the performance for the nominal values. For all studied variations the direction resolution is unaffected, as the direction resolution is dominated by the intrinsic scattering angle and not by detector effects. The energy resolution deteriorates slightly for a lower number of detected photons, i.e. reduced λ_{abs} or QE.

For 20 kHz single noise rates the resolutions are as good as for 10 kHz, confirming the good signal-to-noise ratio due to small time windows in the hit selections (cf. Sec. 3.4.2) allowed by the large scattering length in water.




Figure 91: Resolution of the shower reconstruction for different water properties (λ_{abs} and λ_{scat}) and quantum efficiencies (QE) for upgoing ν_e CC and $\bar{\nu}_e$ CC events. Median neutrino direction (left) and relative energy resolution RMS/E_ν (right).




Figure 92: Resolution of the shower reconstruction for 10kHz and 20kHz single optical noise rates for upgoing ν_e CC and $\bar{\nu}_e$ CC events. Median neutrino direction (left) and relative energy resolution RMS/E_ν (right).

The effective volumes are shown in Fig. 93. For all studied variations in water, PMT and noise properties a similar plateau value is reached, but the turn-on is less steep for less detected photons, i.e. reduced λ_{abs} or QE. For 20 kHz single noise rates the effective volume is only slightly lower compared to a 10 kHz noise rate.

¹⁵ For 20 kHz single noise rate the trigger rate from pure noise would be too high, so that the trigger conditions would have been tightened. However, the purpose of this study is to demonstrate the robustness of the reconstruction with respect to an increased noise rate.

The negligible deterioration in direction and energy resolution in conjunction with the relatively modest loss in effective volume for an increase in single noise rates by a factor of two¹⁶ compared to the nominal assumed rate of 10 kHz demonstrates the robustness of the reconstruction against higher noise rates. Consequently, it is expected that the assumed performance can be achieved for most of the data taking time.




Figure 93: Effective volumes for different water properties, quantum efficiencies and optical background noise rates for upgoing ν_e CC and $\bar{\nu}_e$ CC events. Different absorption λ_{abs} and scattering length λ_{scat} , and quantum efficiencies (QE) (left) and 10 kHz and 20 kHz single optical noise rates (right).

Effect of undetected parameter variations While the direction and energy resolutions are unaffected, Fig. 94 depicts the ratio of the mean reconstructed energy for nominal and varied water and PMT properties. Variations of the same properties and magnitude as above have been used for this study, but the underlying assumption is now that the variation relative to the nominal values is not known and not accounted for in the reconstruction. An exemplary $\pm 10\%$ variation in scattering length has a negligible effect on the mean reconstructed energy, while the same variation in the absorption length induces a corresponding shift in reconstructed energy of $\pm 8\%$. A decrease in quantum efficiency of 10% results in a corresponding downward shift in energy of 10%.




Figure 94: Ratio of the mean reconstructed energy for nominal and varied water properties (λ_{abs} and λ_{scat}) and quantum efficiencies (QE) for upgoing ν_e CC and $\bar{\nu}_e$ CC events.

¹⁶This is even a factor of 2.5 compared to the measured 8 kHz, cf. Sec. 3.2.

3.5 Flavour identification and muon rejection

The determination of the NMH requires a precise estimate of the neutrino energy and zenith angle and a high-purity event sample. In addition, since neutrino events of all flavours are reconstructed, the discrimination between neutrino flavours is necessary. In this section an event type discrimination algorithm is developed and its performance is outlined. The algorithm is conceived with the distinction between three classes of events in mind. These classes are “atmospheric muons”, “shower-like” and “track-like” neutrino events. In particular the atmospheric muon event class is induced by the passage of downward-going muon bundles coming from a cosmic ray air shower which is misreconstructed as upward-going, i.e. neutrino induced, event. Track-like events are those that are induced by charged current muon neutrino interactions, having the signature of a straight track passing through or nearby the instrumented volume. Finally, shower-like events are those coming from all other neutrino interaction channels and flavours: all neutral current interactions and the charged current interactions of electron and tau neutrinos¹⁷.

3.5.1 Methodology

In order to optimally exploit the information imprinted in the light emission of the events, several machine learning algorithms, so called classifiers, have been evaluated. Finally, a classification algorithm known as *Random Decision Forest* (RDF) [170] has been used in this study.

A RDF consists of many decision trees that individually categorise an event into different classes. Each decision tree consists of several nodes. During classification a number of features, i.e. observables contributing discrimination power, are calculated for an event. At each node a decision in favour of a class is taken and the event is pushed to a child node according to the result of the decision. The individual node decisions in a tree are found by a cut on one of the calculated features. The cuts are chosen so that they maximise performance key figures such as the signal class purity. In this way the event is classified as more likely to be a track, a shower or an atmospheric muon. The decision process is repeated until the event reaches a leaf, a node without children, and the classification into one of the classes is finished.

A decision tree is trained on Monte-Carlo event data. A major disadvantage of single trees, however, is the low ability to generalise the trained tree, i.e. the ability to not only reproduce the features in the training sample. Several methods are proposed in the literature to improve the performance of single decision tree methods and we use the RDF approach. For an RDF many of the above described decision trees are trained simultaneously, a total of 101 in our study. Instead of using all features at once, for each tree a predefined fraction of features and events is selected. Finally, the classification is done by a majority decision of the trees as described above. The purity and efficiency of the classification can be set by defining cuts different from a simple 50% majority decision.

3.5.2 Event preselection

Even if the detector will be located under more than 2000 m of sea water, the number of atmospheric muons arriving at the detector and being triggered (cf. Sec. 3.2.4) is larger than that of atmospheric neutrinos by several orders of magnitude. However, since the atmospheric muon flux is fully shielded by the Earth, looking at upward going events will allow to search for neutrinos. Nonetheless, Cherenkov photons from atmospheric muons can produce a hit pattern in the detector such that reconstruction algorithms still reconstruct the event as upward-going. A pre-selection of events is necessary before training the RDF, since in any case it would not be able to handle such a large contamination of atmospheric muons. Both the reconstruction strategies described in the previous section can produce a proper rejection of the atmospheric background without significantly reducing the amount of good neutrino events.

At first each event is requested to be reconstructed as upward going. This holds both for the track and the shower reconstruction algorithm. Then two different sets of quality criteria are applied, one for the muon and one for the shower reconstruction method. The logical “OR” of the two chosen criteria is used to define the input sample for the RDF.

Concerning the shower reconstruction algorithm, a preliminary event selection is implicitly done in the reconstruction itself, cf. Sec. 3.4.3. Through-going atmospheric muons release a large amount of light in the detector and can easily be separated from low energy neutrino showers. This is done by requiring a proper

¹⁷except for those roughly 18% of τ decays producing a muon.

hit selection in the shower reconstruction itself. This is not the case for the muon track reconstruction algorithm, for which both the signal and the background events show the same hit topology. For bright reconstructed shower events it is required that the hit pattern is compatible with a point-like emission. Here, bright events are defined as events with more than 15 causally connected L2 hits (defined as in Sec. 3.4.2) and the compatibility with a point-like emission is evaluated based on the time residuals of these L2 hits with respect to the reconstructed vertex. If the difference between the 80% and 20% quantiles of the time residuals is smaller than 15 ns, the event is considered as a shower event candidate. This requirement results in a preselection of shower event candidates and efficiently focuses the time-consuming part of the shower reconstruction to neutrino-like events. Additionally, the shower reconstruction algorithm provides many different event by event quality parameters, which provide further rejection power for atmospheric muons and are used as features in the RDF.

As far as the track reconstruction is concerned, the Λ parameter described previously can provide a first rejection of atmospheric muons; however, acting on this parameter alone would also suppress a large part of the neutrino sample at lower energy if high purity is requested. Adding also the reconstructed track starting point information allows an improved rejection of wrongly reconstructed atmospheric muon tracks. Fig. 95 and Fig. 96 show the distribution of the reconstructed track starting point for atmospheric muons and low energy ($E_\nu < 20$ GeV) atmospheric muon neutrinos. A variable R_ν , the radius of a "fiducial cylinder", has been defined and, in combination with the Λ quantities, has been tested in order to achieve a preliminary selection cut. The chosen value for R_ν is equal to the radius of the instrumented volume, i.e. 106 m. The number of wrongly reconstructed atmospheric muons can be reduced by more than 3 orders of magnitude when applying a preliminary selection cut on R_ν and the track quality parameter Λ .




Figure 95: r - z distribution of the reconstructed track starting point for upward going reconstructed tracks: Left: atmospheric muons. Right: atmospheric neutrinos below 20 GeV. Upgoing events with $\Lambda > -6$ are shown. The black lines represent the contour of the instrumented volume.

3.5.3 Classification input

As described above, a decision tree relies on cuts on observables, so called features, that are chosen to discriminate well between the different classes and that are calculated for each event. In the following, the best performing features are ranked according to their discrimination power and very briefly explained.

The ranking is done using the overall classification rate under a majority decision of 50%. For the ranking, the trees of the RDF are trained with each individual feature and the overall RDF performance is evaluated. In the next step the algorithm adds one more feature to the best one and does the training




Figure 96: x-y distribution of the reconstructed track starting point for upward going reconstructed tracks: Left: atmospheric muons. Right: atmospheric neutrinos below 20 GeV. Upgoing events with $\Lambda > -6$ are shown. The black lines represent the contour of the instrumented volume.

once more. This is done for every possible configuration. The best configurations are chosen to do more iterations in the same way and this process iterates as long as the performance increases. In Tab. 13 the ranking of the best features used in the classification is listed.

Rank	Feature Description
1	normalized eigenvalue of tensor of inertia of the hit distribution
2	RMS of time residual distribution with respect to a shower hypothesis
3	χ^2 of linear fit to the cumulative time residual distribution
4	reconstructed neutrino energy from shower reconstruction
5	coverage cov_{20° as defined in Sec. 3.4.3
6	number of hit DOMs within $< 10^\circ$ around the reconstructed shower direction and vertex
7	median of time residual distribution of hits selected under a shower hypothesis
8	ratio between number of selected hits for a track hypothesis and shower hypothesis
9	ordinate intercept of a linear fit to cumulative time residual distribution with respect to a shower hypothesis
10	Bjorken y as reconstructed by the shower reconstruction

Table 13: Ten of the best performing features used in the Random Decision Forest classification algorithm.

3.5.4 Classification performance

In the following the performance of the classification algorithm is evaluated using all events passing the selection criteria shown above.

Definitions It is desirable to maximise the number of correctly classified events for all channels. The following definition is used to evaluate the performance of the classification algorithm. The *fraction of*

correctly classified events R_{corr}^A of class A is defined as the ratio of correctly classified events with succeeded reconstruction $N_{corr,rec}^A$ with respect to the total number of events in this class N_{all}^A :

$$R_{corr}^A = \frac{N_{corr,rec}^A}{N_{all}^A} \quad (34)$$

Classification results Fig. 97 shows the result of the RDF classification. The fraction of correctly classified events per interaction channel is plotted versus the MC neutrino energy. Here the majority vote of the random decision forest was set to 50% as this was the best compromise between all classes. Each color depicts the result for neutrinos and antineutrinos of one flavour in one interaction channel.




Figure 97: Fraction of events classified as tracks (left) or showers (right) for a detector with 6 m vertical DOM spacing.

The left plot in Fig. 97 shows the fraction of events classified as track-like for the different flavours and interaction channels versus the MC neutrino energy. The shown results are obtained for all events used in the last classification step. Classification results for charged current tau neutrino interactions are shown without distinction between track-like and shower-like decay topologies of the resulting tau lepton¹⁸. The high energy range shows an expected increase in identification power for long-track muons from muon neutrinos undergoing a charged current interaction. As can be seen, antineutrinos can be identified more easily than neutrinos. This is expected due to the different reaction inelasticities for neutrinos and antineutrinos. The fraction of interactions with a resulting shower signature wrongly identified as track-like falls below 20% above 10 GeV. Electron neutrinos undergoing a charged current interaction are identified more easily as shower-like than neutral current reactions as they yield more light.

In the right plot the fraction of events recognised as showers is depicted. Most efficiently recognized are electron neutrino charged current interactions. Above a neutrino energy of 15 GeV the fraction of correctly classified events reaches more than 90%. At 6 GeV the fraction reaches 85%. Charged current muon (anti-)neutrino events are falsely classified at a rate of 35% (15%) at 10 GeV.

The results for the configuration with a spacing of 9 m (Fig. 98) shows a drop of around 5% in the identification power for track-like events. The general shape of the distribution remains. The fraction of misclassified shower events stays nearly the same. However, shower events need more energy now to result in a clear signature and successful classification as shower-like events. Therefore, the response curve is shifted by 5 GeV to higher energies.

Fig. 99 shows the particle identification performance for a detector configuration with 12 m vertical spacing. The identification power for the charged current muon neutrinos drops significantly. The fraction of misclassified shower events stays below 20%. Again a shift to higher energies of the response curve for shower-like events is observed.

The contamination of atmospheric muons in the neutrino sample, i.e. downward going atmospheric muons which are reconstructed as upgoing and classified either as neutrino induced tracks or as showers is of the order of a few percent. These wrongly identified muons have equal probability of ending up either

¹⁸Note that tau neutrino interactions have been excluded from the event set used for the RDF training.




Figure 98: Fraction of events classified as tracks (left) or showers (right) for a detector with 9 m vertical DOM spacing.




Figure 99: Fraction of events classified as tracks (left) or showers (right) for a detector with 12 m vertical DOM spacing.

in the “showers” or “tracks” sample. This surviving background is taken into account in the subsequent calculation of the ORCA sensitivity.

3.6 Sensitivity studies for the neutrino mass hierarchy

3.6.1 Global fit

This section describes the main mass hierarchy sensitivity calculation based on pseudo-experiments and log likelihood ratios. It is divided into three parts. First, the modelling of the physics and detector is detailed. This model is used to calculate the expected event rates for given values of the oscillation parameters and systematics. Then, the statistical method for the mass hierarchy sensitivity calculation is described. Finally, an overview is given of the current results using this method. An independent study based on Asimov-sets is described at the end of this section.

Rate calculation KM3NeT/ORCA's data will consist of observed event rates as a function of the reconstructed neutrino energy and zenith angle. By comparing these to the expected rates it will be possible to distinguish between the two mass hierarchy cases. The rate computation is separated into two parts. First the expected neutrino interaction rate at the detector site is calculated as a function of the true neutrino energy and zenith angle. Secondly, the response of the detector itself is modelled, leading to the rates of reconstructed events as a function of the *reconstructed* energy and zenith angle.

As shown in Sec. 3.3 and Sec. 3.4, KM3NeT/ORCA is sensitive to the inelasticity (Bjorken y), potentially




Figure 100: A flowchart showing the different steps in the computation chain. The blue blocks show the intermediate results consisting of sets of histograms as a function of the two variables written below the title. The red blocks describe the steps to go from one intermediate result to the next. The green blocks describe additional inputs that do not use the result from the previous step.

adding a third dimension to the rate histograms. At the moment, the inelasticity is not yet included in the sensitivity study. Doing so will likely improve the mass hierarchy significance, due to its power to discriminate neutrinos and antineutrinos on a statistical basis.

The whole computation chain is summarized in Fig. 100. Each step is described in detail in the following paragraphs.

Detector-independent part

The first half of the simulation chain (leading to intermediate result (c) in Fig. 100) can be summarized as:

$$R_a(E, \theta) = \frac{\rho_{\text{water}}}{m_{\text{nucleon}}} \times \sum_b \sigma_a(E) \times P_{a,b}^{\text{osc}}(E, \theta) \times \Phi_b^{\text{atm}}(E, \theta), \quad (35)$$

where

- R_a is the interaction rate per unit volume at the detector site of (anti)neutrinos of flavour a as a function of the neutrino energy and direction.
- The initial flavour b is summed over $\nu_e, \nu_\mu, \bar{\nu}_e$ and $\bar{\nu}_\mu$.
- Φ_b^{atm} is the atmospheric neutrino flux for neutrinos of flavour b .
- $P_{a,b}^{\text{osc}}$ is the oscillation probability for a neutrino passing through Earth.
- σ_a is the charged-current neutrino-nucleon cross section for a neutrino of flavour a .

A consistent binning is used throughout the calculation. The energy axis is binned linearly in $\log_{10}(E)$ from 2 to 100 GeV in 40 bins. The zenith angle axis is binned linearly in $\cos(\theta)$ from -1 to 0 in 40 bins, where we use the convention that $\cos(\theta) = -1$ corresponds to vertically upgoing neutrinos.

The *atmospheric neutrino fluxes* are modelled by the HKKM2014 simulations [171]. The given flux values are tabulated as a function of energy and zenith angle, averaged over the azimuth angle. In order to deal with the rather coarse binning the values are interpolated. To be more precise, a two-dimensional spline interpolation is made of the *cumulative* tables. The spline's derivatives then yield a *bin-integral conserving interpolation* of the flux tables. The chosen tables are for the Fréjus site (without mountain) at solar minimum, since the Fréjus site is expected to be most similar to the KM3NeT/ORCA detector site.

The *oscillation probabilities* depend on the mixing parameters (including the hierarchy) and the Earth density profile. They are calculated by evaluating the neutrino propagation time evolution operator in a

constant density medium (see [172]) at small steps along the trajectory. The Earth's *density profile* is given by the Preliminary Reference Earth Model (PREM) [143]. To speed up calculations the model is approximated by 42 constant-density shells. The electron density (an ingredient for the oscillation probability calculation) is approximated to be half of the nucleon density.

We use the charged current and neutral current neutrino-nucleon cross sections from the GENIE Monte Carlo generator [161, 162] for an oxygen nucleus and two protons.

The results of the first half of the simulation chain (i.e. at intermediate result (c)) are eight histograms of neutrino interaction rates per unit volume at the detector as a function of the true energy and zenith angle: six for the charged current (CC) interactions (three flavours, neutrinos and antineutrinos) and two for neutral current (NC) interactions of neutrinos and antineutrinos. Throughout the simulation, NC events are approximated as equal for all three flavours.

Detector-dependent part

The second part of the simulation chain models the detector response to neutrino interactions. Each step is based on the results presented in the previous sections of this document.

The energy- and zenith angle-dependent *effective mass* determines how many of the interacting events can be reconstructed. This is step (3) in the flowchart. It is derived from MC simulations as

$$M_{\text{eff}} := V_{\text{gen.}} \times \rho_{\text{water}} \times N_{\text{sel.}}/N_{\text{gen.}}, \quad (36)$$

where $N_{\text{gen.}}$ is the total number of generated events in a large generation volume $V_{\text{gen.}}$. Events that are successfully reconstructed by *either one* of the two reconstruction algorithms are counted in $N_{\text{sel.}}$. The density of sea water ρ_{water} is assumed to be 1025 kg/m^3 . The effective mass is binned as a function of the true neutrino energy and zenith angle, and is evaluated for each of the eight event classes separately.

At this step an additional histogram is created, representing the expected background from *misreconstructed atmospheric muons*. As shown in Sec. 3.5, the contamination of such events can be effectively reduced to a few percent by applying cuts. Due to the high suppression efficiency it is increasingly difficult to generate high statistics samples for this type of background. Therefore, the distribution from a looser cut is used and rescaled to the total number of events found for a stricter cut. This is a conservative estimate as for looser cuts the event distribution turns out to be mostly centered around our signal area (upgoing, around 10 GeV) while the distribution becomes more uniform as we apply stricter cuts.

At the end of this step there are a total of nine histograms.

The next step ((4) in Fig. 100) is *particle identification*. Each input histogram is the basis for two new ones, representing events identified as 'tracks' and 'showers', respectively. The identification probabilities are based on the RDF study described in Sec. 3.5, and depend on the true neutrino energy only. Neutrino events identified as atmospheric muons are discarded. The probability for atmospheric muon background events to be identified as a track/shower has not been determined due to lack of statistics; a simple 50/50 separation is applied. After this step we have eighteen histograms.

In the final step ((5) in Fig. 100) the *energy resolutions* and *angle resolutions* are applied. They are implemented as response matrices filled from simulated data. First the zenith angle is 'smeared out' using a three-dimensional response matrix that provides binned $\cos(\theta_{\text{reco}})$ -distributions as a function of $\cos(\theta_{\text{true}})$ and E_{true} . Then a two-dimensional energy response matrix providing E_{reco} -distributions as a function of E_{true} is used to smear the energy.

The resolutions are evaluated separately for each of the sixteen neutrino event classes. We have, for example:

- ν_{μ} CC interactions identified as tracks
- $\bar{\nu}_e$ CC events identified as showers
- NC ν events identified as showers
- ν_e CC events misidentified as tracks
- ...

Each is smeared using dedicated response matrices. In particular, neutrinos and antineutrinos are smeared differently. So are correctly and wrongly identified events.

The response matrices use a coarser binning than the rate histograms. Depending on the available MC statistics, the number of bins is reduced from 40 to 20 or 10 to avoid artifacts.

<i>parameter</i>	<i>true value distr.</i>	<i>initial value distr.</i>	<i>treatment</i>	<i>prior</i>
θ_{23} [°]	{40, 42, . . . , 50}	uniform over [35, 55] †	fitted	no
θ_{13} [°]	8.42	$\mu = 8.42, \sigma = 0.26$	fitted	yes
θ_{12} [°]	34	$\mu = 34, \sigma = 1$	nuisance	N/A
ΔM^2 [10^{-3} eV ²]	$\mu = 2.4, \sigma = 0.05$	$\mu = 2.4, \sigma = 0.05$	fitted	no
Δm^2 [10^{-5} eV ²]	7.6	$\mu = 7.6, \sigma = 0.2$	nuisance	N/A
δ_{CP} [°]	0	uniform over [0, 360]	fitted	no
overall flux factor	1	$\mu = 1, \sigma = 0.1$	fitted	yes
NC scaling	1	$\mu = 1, \sigma = 0.05$	fitted	yes
$\nu/\bar{\nu}$ skew	0	$\mu = 0, \sigma = 0.03$	fitted	yes
μ/e skew	0	$\mu = 0, \sigma = 0.05$	fitted	yes
energy slope	0	$\mu = 0, \sigma = 0.05$	fitted	yes

Table 14: Default parameter settings used for the LLR analysis. Where μ and σ are given, they refer to a Gaussian distribution. The † indicates that the initial values for θ_{23} are generated in a special way: a total of seven initial values is tried. They are $x + i \times 5^\circ$, where x is the randomly drawn value and $i \in [-3, -2, \dots, 3]$.

After reconstruction all histograms are combined in two final event histograms representing the track channel and the shower channel, respectively.

Sensitivity calculation The sensitivity to the mass hierarchy is calculated using likelihood ratio distributions from *pseudo-experiments* (PEs). The procedure works as follows:

1. Pick a set of true values for the oscillation parameters and other systematics.
2. Calculate the expected number of events for a given period of data taking, using the simulation chain described above.
3. Generate pseudo-data by randomly drawing a detected number of events for each bin based on Poisson statistics. The two histograms thus attained constitute the PE.
4. Find the best-fit likelihoods \mathcal{L}_{NH} and \mathcal{L}_{IH} for the NH and IH assumption, by maximizing the likelihood with respect to the other free parameters in both cases.
5. Calculate the log likelihood ratio $LLR := \log(\mathcal{L}_{NH}/\mathcal{L}_{IH})$. This discriminating variable indicates which hierarchy is favoured by the PE.

The *likelihood* \mathcal{L} of the PE given the model is defined as

$$\mathcal{L} = \prod_{i \in \text{bins}} \mathcal{P}(N_i | \mu_i), \quad (37)$$

where N_i and μ_i are, respectively, the observed and expected number of events in bin i and

$$\mathcal{P}(n | \lambda) = \frac{\lambda^n e^{-\lambda}}{n!} \quad (38)$$

is the Poisson probability to observe n events when the expectation value is λ . The expected event numbers μ_i depend on the parameter values (oscillation parameters and systematics) so that maximizing the likelihood corresponds to finding the parameter values that best fit the PE.

The default parameter settings are summarized in Tab. 14. It shows the *true parameter values* used to generate PEs. Most of these are fixed at some nominal value. The *initial values* are those used as starting values by the minimizer in the fitting procedure that finds the likelihood maximum. These values are chosen randomly for each PE to avoid systematic biases. Most parameters are fitted, meaning they are left free in the minimizer. The likelihood is multiplied by Gaussian priors for some parameters (see Tab. 14). The mean and width of the Gaussian priors correspond to those of the matching initial value distributions. Two parameters (θ_{12} and δm^2) are treated as *nuisance parameters*. This means that, rather than leaving them free in the fit, a random ‘best fit’ value from the initial value distribution is assigned to each PE. It emulates the fact that these parameters will be constrained almost exclusively by external measurements.




Figure 101: Example of log likelihood ratio (LLR) distributions for true NH (red) and true IH (blue) pseudo-experiments. The dashed curves represent Gaussian fits and the dashed vertical lines mark the median of the distributions.

The first six parameters listed in Tab. 14 are the oscillation parameters, where the large mass-squared difference ΔM^2 is defined as

$$\Delta M^2 := \frac{\Delta m_{32}^2 + \Delta m_{31}^2}{2}. \quad (39)$$

The last five entries in the table are *systematics*. The *overall flux factor* and *NC scaling* simply scale the total number of (NC) events by an energy- and zenith-independent factor. The *skew parameters* introduce an additional asymmetry in the ratio of one event type to the other, while conserving the total number of events. They relate to the ratio of neutrinos to antineutrinos and the ratio of μ -flavour events to e -flavour events. Finally, the *energy slope* α introduces an energy-dependent scaling of the number of events of the form E^α .

Because θ_{23} generally has two likelihood maxima, special steps are taken to avoid ending up in the wrong one. The likelihood maximization is repeated several times, starting from a different θ_{23} value each time. Only the best-fit result is considered for either hierarchy.

The distributions for each parameter are uncorrelated and based on the current world uncertainties [89, 91, 173].

The final figure of merit is the *median significance*, computed by comparing LLR distributions for true NH and true IH PEs. An example is shown in Fig. 101. The further these two distributions are apart, the higher the significance. We quote the significance with which the ‘wrong hierarchy’ can be excluded at the median of the ‘true hierarchy’ LLR. In all our simulations the LLR distributions can be excellently approximated by Gaussians. This allows the median significance to be expressed in the following simple form:

$$S_{\text{NH}} := \frac{\mu_{\text{NH}} - \mu_{\text{IH}}}{\sigma_{\text{IH}}}. \quad (40)$$

This gives the median significance in σ ’s to exclude the IH, given true NH, where the μ ’s and σ ’s here refer to the means and widths of the LLR distributions. This number can be interpreted as the minimum significance in σ ’s with which the wrong hierarchy can be excluded in at least half of the pseudo-experiments.

Alternative Hypothesis Initially, the mass hierarchy sensitivity was calculated by comparing LLR distributions generated with identical true oscillation parameter values (other than the hierarchy). However, this approach does not take into account the strong correlation between the measurement of θ_{23} and the hierarchy. From simulations it follows that the best-fit value of θ_{23} depends strongly on the assumed hierarchy. In many cases, the best-fit values for the two hierarchy assumptions are not in the same octant. In the actual measurement we therefore have to distinguish between two cases: NH with some best-fit value θ_{23}^{NH} and IH with a *different* best-fit value θ_{23}^{IH} . Since the two values can be very far apart, the mean and width




Figure 102: Mean and width of the LLR distributions for NH (left) and IH (right) pseudo-experiments as a function of θ_{23} . These values were obtained from a Gaussian fit of the distributions.

of the corresponding LLR distributions can be noticeably different, leading to a different mass hierarchy sensitivity. Note that this effect does not occur for the other parameters, which typically have very similar best-fit values for the two hierarchy assumptions.

To take this effect into account the following procedure was adopted. For each true hypothesis (true hierarchy TH with $\theta_{23,\text{true}}$) the most likely *alternative hypothesis* (other hierarchy OH with $\theta_{23,\text{alt}}$) is determined from the θ_{23} best-fit distribution of PEs generated with the true hypothesis and fitted assuming the OH. The median significance to reject the alternative hypothesis is then calculated:

$$S_{\text{true}} := \frac{|\mu_{\text{true}} - \mu_{\text{alt}}|}{\sigma_{\text{alt}}} \quad (41)$$

A technical issue arises because the LLR distributions were only simulated for certain given values of θ_{23} , while the alternative hypothesis θ_{23} 's can take any value. To overcome this we notice that the LLR distributions' fitted widths and means as a function of θ_{23} look rather smooth, so that they can be reasonably approximated by interpolating between the already calculated values. This is shown in Fig. 102. This method enables us to calculate the mass hierarchy sensitivity for any value of the true and alternative θ_{23} . Fig. 103 illustrates the values of the alternative θ_{23} and the effect on the mass hierarchy sensitivity. All the LLR-method mass hierarchy sensitivity results in this document are produced using this method, unless explicitly stated otherwise.

Fig. 104 shows the effect of the new method on the mass hierarchy sensitivity.

Results Fig. 105 shows the latest mass hierarchy significance plot. The expected significance depends strongly on the true value of θ_{23} and δ_{CP} . Without CP-violation, the neutrino mass hierarchy can be measured with more than 3σ in three years at the current world best fit values of θ_{23}

3.6.2 Spacing studies

Whereas the LLR method (described above) provides the most accurate description of the planned experiment, its application to certain problems might sometimes be prohibitive due to the large number of pseudo-experiments to be generated. Therefore a simplified approach is used to answer dedicated questions.

The starting point is again the set of two histograms (for tracks and showers) in the reconstructed quantities $E_{\text{reco}}, \theta_{\text{reco}}$. In each bin i , the expected number of events (μ_i^{TH}) for a given true hierarchy (TH) hypothesis is calculated. A χ^2 minimisation is performed assuming the wrong hierarchy (WH) marginalizing over the parameters given in Tab. 14. Contrary to the description in Tab. 14, θ_{12}, θ_{13} and Δm^2 are fixed to their true values, whereas all other 8 parameters are fitted unconstrained, *i.e.* without adding any priors. The true value of the CP-phase is fixed to 0. As a result of the minimisation, a χ_{min}^2 is obtained

$$\chi_{\text{min}}^2 = \sum_i \frac{(\mu_i^{\text{TH}} - \mu_i^{\text{WHfit}})^2}{\mu_i^{\text{TH}}} \quad (42)$$




Figure 103: The mass hierarchy sensitivity for the true normal (left) and inverted (right) hierarchy. The horizontal axis indicates the true value of θ_{23} . The vertical axis indicates the ‘alternative’ value of θ_{23} : the value belonging to the hypothesis that is being rejected. The diagonal dashed lines indicate the position where the alternative θ_{23} is the same as the true one. Along these lines the mass hierarchy sensitivity according to the original method can be read off. The solid red and blue lines show the most likely alternative value for each true θ_{23} . They are the most likely value when fitting θ_{23} under the wrong hierarchy assumption. Along these lines the mass hierarchy sensitivity according to the new method can be read off.




Figure 104: Comparison of the mass hierarchy sensitivity calculated using the old method (dashed lines) and the new method (solid lines). In the former, the significance is calculated to reject the other hierarchy at the same θ_{23} , whereas in the latter the alternative hypothesis has a different θ_{23} . The differences are rather small, but there is a noticeable decrease in the second octant IH mass hierarchy sensitivity. This is for the 9 m spacing and three years of operation time, using the default settings (particularly, $\delta_{CP,true} = 0^\circ$).




Figure 105: The mass hierarchy sensitivity for three years of operation time for the 9 m spacing, using the default settings. This includes a fit of θ_{23} , ΔM^2 , δ_{CP} and the five systematics. The dashed lines indicate the mass hierarchy sensitivity when the true value of δ_{CP} is set to 180° .

which determines the probability to refute the WH hypothesis. It can be simply expressed as

$$\sigma = \sqrt{\chi_{min}^2} \quad (43)$$

It has been verified, that the results obtained with this method are generally rather close to those from the full LLR treatment. The simplified method is used to optimize the vertical distance of the DOMs on the DUs. Whereas the horizontal spacing between DUs is determined by deployment constraints (20m distance between DUs is considered a minimum), the vertical distance is a free parameter with little constraints from a technical point of view. Simulations have been performed with DOM distances of 6 m, 9 m and 12 m. The detector performance for these different setups have been shown before. Fig. 106 shows the expected NMH sensitivity after three years of data taking for both hierarchy hypotheses as function of the true mixing angle θ_{23} . An optimal distance is found close to 9 m, as both for 6 m and 12 m the NMH sensitivity degrades, at least in some regions of the parameter space.

3.6.3 Measurement of ΔM^2 and θ_{23}

The derivation of measurement contours for the oscillation parameters is done as well with the simplified procedure, which had been used already for the spacing study. The same set of nuisance parameters is applied. Optionally an energy scale shift is added as additional systematic uncertainty. It is implemented as a free scaling of the neutrino energy in all detector related distributions such as effective mass, particle identification, angular and energy resolution. All nuisance parameters are fitted unconstrained, *i.e.* without priors. Both ΔM^2 and θ_{23} are determined under the assumption that the correct NMH has been already identified. The 1σ measurement contours obtained after three years of data taking for three test points ($\Delta M^2 = 2.45 \cdot 10^{-3} \text{eV}^2$, $\sin^2 \theta_{23} = 0.42, 0.50, 0.58$) are shown on Fig. 107. They are compared to current world best measurements [100, 174] as well as to extrapolations of final results from NOvA and T2K [100, 175], to be expected around 2020. For T2K, the extrapolation is performed by exploiting the published likelihood shape of the present measurement [100] assuming the planned complete beam exposure of $7.8 \cdot 10^{21}$ protons on target. A precision of 3% in ΔM^2 is reached after three years which can be reduced to 2% when suppressing the energy scale uncertainty. The precision in θ_{23} varies between 4% and 10%, depending on its true value and the NMH.




Figure 106: NMH Sensitivity for 3 years of data taking as function of the true mixing angle θ_{23} for both hierarchy hypotheses. Different vertical distances between adjacent DOMs are simulated: red 6 m, blue 9 m, magenta 12 m.




Figure 107: Measurement precision in ΔM^2 and $\sin^2 \theta_{23}$ after three years of data taking with ORCA with (solid red) and without (dashed red) energy scale uncertainty for three test points compared to present results from MINOS (black) [174] and T2K (blue) [100] and predicted performance of NOvA (magenta) [175] and T2K (blue, dashed) [100] in 2020. All contours are at 1σ , left for NH, right IH.

3.6.4 Systematic uncertainties

A substantial list of possible uncertainties is already taken into account while fitting the NMH by marginalizing over the related nuisance parameters, as indicated in Tab. 14. Some of these parameters – such as θ_{23} and ΔM^2 – can be determined together with the NMH with high accuracy, as shown above.

It is crucial to determine reliable priors for the chosen nuisance parameters. The currently used priors are listed as well in Tab. 14. However, it has been verified that loosening the prior conditions or even totally suppressing them has only a small impact on the final NMH sensitivity. Therefore, in future studies some of them might be treated as unconstrained fit parameters, *i.e.* without priors.

Contributions to the uncertainties come from the neutrino flux [171], cross section [155] and from the detector performance. For the latter, a main contribution is expected from the uncertainty in the photon detection efficiency by the PMTs and the related readout electronics. However, as demonstrated in ANTARES and also with the KM3NeT prototype module [5], the measurement of ^{40}K coincidences between adjacent PMTs of the same DOM allows the photon detection efficiency to be monitored in real time with high precision. The variable nature of optical noise due to bioluminescence is controlled by sampling it for each individual PMT with a frequency of 10 Hz. The results of these measurements are directly injected into the simulation, as is done in ANTARES. This excludes bioluminescence as a source of systematic uncertainty of any measurement. Apart from the optical noise due to bioluminescence, sea water is a very stable and homogeneous medium, as monitored over many years by ANTARES. Current knowledge of its light propagation properties are discussed in Sec. 2.4. The residual uncertainties of quantities such as absorption and scattering length have less effect (due to the closer spacing) for ORCA than for ARCA, and are well-covered by the nuisance parameters discussed above, so no separate investigation has been performed.

Additional systematic uncertainties, not yet included in the present study, comprise systematic shifts in the reconstructed energy and zenith angle. These will be considered in the near future. However, it is believed that the energy scale is well-constrained through the knowledge of the absolute PMT efficiency and the water parameters. The angular resolution of neutrino telescopes in sea water is excellent, and it remains better than 10 degrees down to the energies relevant for the NMH determination. Systematic angular offsets are at most in the sub-degree region, as shown by the study of the moon shadow in the cosmic ray signal in ANTARES. A deterioration of the angular resolution due to the movement of the detector elements in the sea current is excluded by permanently re-calibrating them via an acoustic positioning system. Such a system provides a precision of better than about 10 cm for all detector elements, which makes its influence on the angular resolution of reconstructed neutrino events negligible.

The energy and angular resolutions are a crucial input to the sensitivity calculation. Both are estimated from simulations which are subject to uncertainties on their own. These can be parametrised by applying a scaling (*i.e.* broadening or narrowing) to these resolution functions, as is planned for the near future.

Finally, an independent study has been performed to study the variations of the Earth model [143] on the NMH sensitivity. Both the thickness and density of each individual layer have been varied within the tolerance of the model, as well as the sharpness of the layer boundaries. The impact of these variations is found to be negligible for the present study and is therefore ignored as a relevant systematic effect.

3.7 Outlook

The previous sections provide details on the performances of the KM3NeT/ORCA detector in establishing the neutrino mass ordering and improving the precision on the oscillation parameters in the atmospheric sector. While the results obtained so far rely on full Monte Carlo studies and incorporate the leading systematic effects, possible refinements have already been identified and will be scrutinized in the near future. This includes notably the usage of the achieved sensitivity to the interaction inelasticity as a statistical tool to discriminate between neutrino and antineutrinos and to further reject the background from neutral current interactions. As a detailed and ongoing investigation shows, see [156] for first results, event-by-event fluctuations intrinsic to the development of the hadronic system resulting from a neutrino interaction on a nucleon or nucleus limit the achievable resolutions on direction, energy and inelasticity for a given detector geometry. However, some improvements, in particular for the reconstruction of the reaction inelasticity, can still be expected to arise from the development of new reconstruction strategies. Envisaged lines of work comprise e.g. an attempt to identify the leading particle in the hadronic system to constrain its overall

momentum, a combined fit of the hadronic shower and the charged lepton in CC interactions, improved energy estimation techniques, etc.

In addition to the oscillation measurements discussed in the previous section, the size and energy range covered by the KM3NeT/ORCA detector allow for the search of CC interactions of tau neutrinos produced in the oscillation of atmospheric electron and muon neutrinos. While these events can hardly be distinguished on an event-by-event basis, their presence could be revealed by a statistical excess of cascade-like events over the baseline from atmospheric NC interactions and electron neutrino CC interactions. This effect is expected to be seen with high confidence level and statistical power within the first years of operation, but a precise study remains to be carried out. The energy and flavour distributions of observed events in the ORCA detector could in principle also reveal sizeable discrepancies from expectations due to non standard physics interactions (NSI) [176, 177]. While strong deviations from expectations (e.g enhanced CP violation effects) might deteriorate the sensitivity to the NMH, a more likely scenario is that KM3NeT/ORCA will be able to invalidate many NSI processes.

The studies presented here indicate that the current unknown value of the Dirac CP violating phase in the neutrino sector mildly impacts the sensitivity to the neutrino mass ordering. However, the knowledge of the mass ordering could reversely bring sensitivity to the CP phase in the (0.2 – 1) GeV regime [137]. This would imply a denser instrumentation than what is currently envisaged for ORCA, but considering the importance of measuring the CP phase, sensitivity studies could be performed for a further step of the ORCA project. In the same spirit, sensitivity studies for both the NMH and the CP phase have been proposed relying on a putative upgraded neutrino beam to be sent to ORCA from Protvino [178, 179]. Such a strategy would in particular allow for a confirmation of ORCA-only results on the NMH with high statistical power on a short (< 1 yr) timescale [180]. It would require a new beamline to be setup but would offer the advantage to rely on an already built detector.

With its low energy threshold the KM3NeT/ORCA detector offers the possibility to extend searches started with ANTARES (e.g [73, 75]) (and likely to be pursued with KM3NeT/ARCA as well) for extra-terrestrial neutrinos as a signature of the presence of Dark Matter in the centre of the Earth, the Sun and the central region of the Galaxy for which the detector is particularly well located. The low energy threshold of KM3NeT/ORCA is particularly well suited to constrain low-mass WIMP (Weakly Interacting Massive Particle) Dark Matter models. All neutrino flavours could be used for such studies, considering the encouraging first performances in the shower reconstruction channel.

GeV neutrinos are also likely to be emitted by several classes of astrophysical objects like low-energy Gamma-Ray Bursts [181] or Colliding Wind Binaries [182]. Another promising topic is the ability of KM3NeT/ORCA to detect neutrinos from supernovae (SN) explosions. The use of segmented optical modules closely placed to one another indeed offers new detection capabilities: asking for coincidences of many phototubes on individual storeys is expected to strongly reduce the optical background potentially providing high sensitivity to SN up to few tens of kpc. These results will possibly be incorporated in an update of the present document, together with the prospects for several other physics studies that can be undertaken with KM3NeT/ORCA. These span a wide range of scientific fields, including the Earth and Sea Sciences which are not addressed here but are part of the scientific scope of deep-sea neutrino observatories. As an example, a detailed study of the neutrino energy and angular distributions could provide tomographic information on the electron density [183–185], and thus on the composition, of the different Earth layers traversed. Such an approach is complementary to the standard methods used in geophysics, which do not univocally constrain the chemical composition of the Earth, in particular of its innermost layers (mantle and core).

4 Organisation

KM3NeT federates and unifies the various smaller European efforts in the field of Neutrino Astronomy. The process of convergence was supported by an EU funded Design Study (2008–2009) and Preparatory Phase (2008–2012). The KM3NeT consortium has now formed a collaboration with an elected management. The funding agencies (or funding authorities) involved have installed the Resources Review Board (RRB) which oversees the project. The RRB is advised by an international Scientific and Technical Advisory Committee (STAC). A project organisation is setup with the objective to implement the first phase (Phase-1) of the KM3NeT Research Infrastructure. To this end, a Memorandum of Understanding (MoU), covering the total available budget of about 31 M€, has been signed by the members of the RRB. The purpose of this MoU is to define the programme of work to be carried out for this phase and the distribution of charges and responsibilities among the Parties and Institutes for the execution of this work. The MoU sets out i) the organizational, managerial and financial guidelines to be followed by the collaboration, ii) the external scientific and technical review processes and iii) the user access policy. At present, the collaboration consists of more than 240 persons from 52 institutes. The first phase has already started and comprises the final prototyping and preproduction, engineering, construction, calibration, transportation, assembly, installation and commissioning of the elements which form the basis of the KM3NeT neutrino detector and the seafloor and shore station infrastructures as well as the operation of the installed neutrino detectors. The installation is proceeding in two places, off-shore Toulon, France and off-shore Capo Passero, Italy. A third suitable site is available off-shore Pylos, Greece. The construction of the Phase-1 detector has already started with the successful deployment of the first string off-shore Capo Passero and will be completed by 2017.

The Collaboration will offer open access for external users to the KM3NeT Research Infrastructure (Article 15 of the MoU). The KM3NeT Research Infrastructure will also provide user ports for continuous Earth and Sea science measurements in the deep-sea environment. The needs for the Earth and Sea sciences are partly incorporated in the present KM3NeT MoU and other needs will be detailed in designated MoUs between KM3NeT and individual Earth and Sea science groups or more generally with EMSO.

The Phase 1 MoU is a first step towards the intended establishment of a European Research Infrastructure Consortium (ERIC). The collaboration has agreed to host the KM3NeT ERIC in the Netherlands. The neutrino signal recently reported by IceCube has led the KM3NeT Collaboration to propose an intermediate phase (i.e. Phase-2.0). The required actions for the next phase(s) are being taken which include the preparation of requests for additional ERDF funds in France, Italy and Greece as well as requests for national funds. Other support options, e.g. within the framework of Horizon 2020, will also be explored.

Recently, the KM3NeT and ANTARES Collaborations have agreed to organise each general assembly jointly (typically 3–4 times per year). This agreement fosters the scientific progress and the exchange of know-how and limits travel times and expenses. Following a sequence of joint meetings between ANTARES (Mediterranean Sea), IceCube (South Pole), Lake Baikal (Russia) and KM3NeT (Mediterranean Sea), a Memorandum of Understanding for a Global Neutrino Network (GNN) has been signed on 15 October 2013 by the representatives of each project. This step formalises the active collaboration between these projects. Once infrastructures of similar scale are operational on the three continents, the stated aim of the GNN is a worldwide Global Neutrino Observatory.

5 Data policy

The KM3NeT Collaboration has developed a data policy based on the research, educational and outreach goals of the facility. The first exploitation of the data is granted to the collaboration members as they build, maintain and operate the facility and to priority users. Accordingly, each collaboration member has full access rights to all data, software and know-how. Access for non-members is restricted, as long as methods and results have not yet been published. The prompt dissemination of scientific results, new methods and implementations is a central goal of the project, as is education. High-level data (event information enriched with quality information) will be published after an embargo time of two years under an open access policy on a web-based service. Exceptional access rights that correspond to these goals can be granted.

The Collaboration has developed measures to ensure the reproducibility and usability of all scientific results over the full lifetime of the project and in addition 10 years after shutdown. Low-level data (as recorded by the experiment) and high-level data will be stored in parallel at central places. A central software repository, central software builds and operation system images are provided and continuously maintained until the end of the experiment.

The storage and computing needs of the KM3NeT project are highly advanced. The Collaboration has developed a data management plan and a corresponding computing model to answer those needs. The latter is based on the LHC computing models utilizing a hierarchical data processing system with different layers (tiers). Data are stored on two main storage centres (CCIN2P3-Lyon, CNRS and CNAF, INFN); those large data centres are fully interfaced with the major European e-Infrastructures, including GRID-facilities (ReCaS, HellasGRID provide resources to KM3NeT). The main node for processing of the neutrino telescope data is the computer centre in Lyon (CCIN2P3-Lyon). A corresponding long-term and sustainable commitment has already been made by CNRS, which is consistent with the needs for long-term preservation of the data. A specialised service group within the Collaboration will process the data from low-level to high-level and will provide data-related services (including documentation and support on data handling) to the Collaboration and partners. WAN (GRID) access tools (e.g. xrootd, iRODS, and gridFTP) provide the access to high-level data for the Collaboration. The analysis of these data will be pursued at the local e-Infrastructures of the involved institutes (both local and national). The chosen data formats allow for the use of common data analysis tools (e.g. the ROOT data analysis framework) and for integration into e-Infrastructure common services.

The central services are mainly funded through CNRS and INFN that have pledged resources of their main computing centres to the project. Additional storage space and its management are provided by the partner institutes (e.g. INFN has provided 500 TB of disk space for KM3NeT at the ReCaS GRID infrastructure, the Hellenic Open University has pledged 100 TB of disc space and 300 cores to the project).

In addition to the major storage, networking and computing resources provided by the partner institutions and their computing centres, grid resources have been pledged and will be used by KM3NeT (ReCaS, HellasGRID). These will provide significant resources to be used for specialised tasks (as e.g. for special simulation needs). The major resources, however, will be provided by the partners. External services are employed to integrate the KM3NeT e-Infrastructure into the European context of the GRID – in the fields of data management, security and access; services will be implemented in collaboration with EGI.

One of the aims of the KM3NeT data management plan is to play an active role in the development and utilization of e-Infrastructure commons. KM3NeT will therefore contribute to the development of standards and services in the e-Infrastructures both in the specific research field and in general. In the framework of the Global Neutrino Network (GNN), KM3NeT will cooperate with the ANTARES, IceCube and GVD collaborations to contribute to the open science concept by providing access to high-level data and data analysis tools, not only in common data analyses but also for use by citizen scientists.

In the framework of the ASTERICS project, KM3NeT will develop an interface to the Virtual Observatory including training tools and training programmes to enhance the scientific impact of the neutrino telescope and encourage the use of its data by a wide scientific community including interested citizen scientists. Data derived from the operation of the experiment (acoustics, environmental monitoring) will be of interest also outside of the field. Designated documentation and courses for external users will therefore be put in place to facilitate the use of the repositories and tools developed and used by the KM3NeT Collaboration.

6 Cost and Time Schedule

The investment budget for the construction of the first phase (Phase-1) of the KM3NeT research infrastructure, which is fully funded, amounts to about 31 M€. During 2015-2017, 31 strings equipped with 558 optical modules will be assembled and deployed at the French and Italian sites. The overall size of the initial Phase-1 arrays corresponds to about 0.2 building blocks.

The next phase (i.e. KM3NeT 2.0) comprises a complete ARCA and ORCA detector, consisting of 2 and 1 building blocks, respectively. The additional budget for KM3NeT 2.0 is estimated at 95 M€. The cost estimates of KM3NeT 2.0 are based on the actual prices of Phase-1 and thus can be considered accurate.

The breakdown of the cost amongst the major items is illustrated in Fig. 108. They are consistent with the estimations stated in the KM3NeT Technical Design Report published in 2011 and represent a factor of four cost reduction compared to that previously achieved for the ANTARES detector. The cost of a single KM3NeT string is about 230 k€, an additional 90 k€ is needed for the interlink cable, the string deployment and the ROV connection.

Once the funds for Phase-2.0 are available, the array could be constructed within three years. Thus, if funds were forthcoming in 2017, the full array could be completed in 2020. Note that physics studies would already be possible as the array is being constructed, thus reducing the overall time needed to obtain a specified precision. The cost for operation and decommissioning of the infrastructure have been evaluated and amount to about 2 M€ per year and 5 M€, respectively. Hence, the total cost for 10 years of operation and decommissioning of the KM3NeT 2.0 infrastructure adds about 25% to the total budget.




Figure 108: Breakdown of costs amongst the major items.

References

- [1] M. Ageron et al., ANTARES Coll., *ANTARES: the first undersea neutrino telescope*. Nucl. Inst. Meth. **A 656**, 11 (2011), [arXiv:1104.1607 [astro-ph.IM]].
- [2] P. Kooijman for the KM3NeT Coll., in *Proc. 34th Int. Cosmic Ray Conference*, Vol. 001, 1173. PoS, 2015.
- [3] S. Adrián-Martínez et al., KM3NeT Coll., *The prototype detection unit of the KM3NeT detector*, 2015. arXiv:1510.01561 [astro-ph.IM].
- [4] R. Bruijn and D. van Eijk for the KM3NeT Coll., in *Proc. 34th Int. Cosmic Ray Conference*, Vol. 001, 1157. PoS, 2015.
- [5] S. Adrián-Martínez et al., KM3NeT Coll., *Deep sea tests of a prototype of the KM3NeT digital optical module*. Eur. Phys. J. **C 74**, 3056 (2014), [arXiv:1405.0839 [astro-ph.IM]].
- [6] S. Biagi, T. Chiarusi, P. Piattelli and D. Real for the KM3NeT Coll., in *Proc. 34th Int. Cosmic Ray Conference*, Vol. 001, 1172. PoS, 2015.
- [7] B. Bakker, *Trigger studies for the ANTARES and KM3NeT neutrino detector*, 2011. Universiteit van Amsterdam, The Netherlands, M. Sc. Thesis.
- [8] M. de Jong for the KM3NeT Coll., in *Proc. 33rd Int. Cosmic Ray Conference*, Vol. 1, 0891. 2013, available on http://www.cbpf.br/~icrc2013/proc_icrc2013.html.
- [9] S. Adrián-Martínez et al., ANTARES and IceCube Coll., *First combined search for neutrino point-sources in the Southern Hemisphere with the ANTARES and IceCube neutrino telescopes*, 2015. arXiv:1511.02149 [hep-ex].
- [10] M. G. Aartsen et al., IceCube Coll., *Evidence for high-energy extraterrestrial neutrinos at the IceCube detector*. Science **342**, 1242856 (2013), [arXiv:1311.5238 [astro-ph.HE]].
- [11] C. Kopper, W. Giang and N. Kurahashi for the IceCube Coll., in *Proc. 34th Int. Cosmic Ray Conference*, Vol. 001, 1081. PoS, 2015.
- [12] M. G. Aartsen et al., IceCube Coll., *A combined maximum-likelihood analysis of the high-energy astrophysical neutrino flux measured with IceCube*. Astrophys. J. **809**, 98 (2015), [arXiv:1507.03991 [astro-ph.HE]].
- [13] M. G. Aartsen et al., IceCube Coll., *Flavour ratio of astrophysical neutrinos above 35 TeV in IceCube*. Phys. Rev. Lett. **114**, 171102 (2015), [arXiv:1502.03376 [astro-ph.HE]].
- [14] R. Abbasi et al., IceCube Coll., *Search for ultrahigh-energy tau neutrinos with IceCube*. Phys. Rev. **86**, 022005 (2012), [arXiv:1202.4564 [astro-ph.HE]].
- [15] D. Williams for the IceCube Coll., *Search for ultra-high energy tau neutrinos in IceCube*. Nucl. Phys. Proc. Suppl. **B 253**, 155 (2014).
- [16] M. Honda et al., *Calculation of atmospheric neutrino flux using the interaction model calibrated with atmospheric muon data*. Phys. Rev. **D 75**, 043006 (2007), [arXiv:astro-ph/0611418].
- [17] R. Enberg, M. H. Reno and I. Sarcevic, *Prompt neutrino fluxes from atmospheric charm*. Phys. Rev. **D 78**, 043005 (2008), [arXiv:0806.0418 [astro-ph]].
- [18] M. G. Aartsen et al., IceCube Coll., *Search for a diffuse flux of astrophysical muon neutrinos with the IceCube 59-string detector*. Phys. Rev. **D 89**, 062007 (2014), [arXiv:1311.7048 [astro-ph.HE]].
- [19] R. Gauld, J. Rojo, L. Rottoli, S. Sarkar and J. Talbert, *The prompt atmospheric neutrino flux in the light of LHCb*, 2015. arXiv:1511.06346 [astro-ph.HE].
- [20] R. Gandhi, C. Quigg, M. Hall Reno and I. Sarcevic, *Ultrahigh-energy neutrino interactions*. Astropart. Phys. **5**, 81 (1996), [arXiv:hep-ph/9512364].
- [21] G. Ingelman, A. Edin and J. Rathsman, *LEPTO 6.5 – A Monte Carlo generator for deep-inelastic lepton-nucleon scattering*. Comp. Phys. Comm. **101**, 108 (1997), [arXiv:hep-ph/9605286].
- [22] G. D. Barr, *The separation of signals and background in a nucleon decay experiment*. Ph.D. Thesis, University of Oxford, 1987.
- [23] P. Antonioli et al., *A three-dimensional code for muon propagation through the rock: Music*. Astropart. Phys. **7**, 357 (1997), [arXiv:hep-ph/905408].

- [24] S. Jadach, J. H. Kühn and Z. Was, *TAUOLA – a library of Monte Carlo programs to simulate decays of polarized τ leptons*. Comp. Phys. Comm. **64**, 275 (1991).
- [25] C. Carminati et al., *Atmospheric MUons from PArametric formulas: A fast GEnerator for neutrino telescopes (MUPAGE)*. Comp. Phys. Comm. **179**, 915 (2008), [arXiv:0802.0562 [physics.ins-det]].
- [26] D. Heck et al., CORSIKA web page. Available on <http://www.ikp.kit.edu/corsika/>.
- [27] S. Schönert, T. K. Gaisser, E. Resconi and O. Schulz, *Vetoing atmospheric neutrinos in a high energy neutrino telescope*. Phys. Rev. **D 79**, 043009 (2009), [arXiv:0812.4308 [astro-ph]].
- [28] H. S. Fesefeldt, *The simulation of hadronic showers: Physics and applications*, Technical report pitha-85-02, RWTH Aachen, 3. Phys. Inst., 1985.
- [29] N. N. Kalmykov, S. S. Ostapchenko and A. I. Pavlov, *EAS and a quark-gluon string model with jets*. Bull. Russ. Acad. Sci. (Physics) **58**, 1966 (1994).
- [30] T. Heid, C. W. James and K. Pikounis for the KM3NeT Coll., in *Proc. 34th Int. Cosmic Ray Conference*, Vol. 001, 1067. PoS, 2015.
- [31] C. M. F. Hugon, in *Proc. 34th Int. Cosmic Ray Conference*, Vol. 001, 1106. PoS, 2015.
- [32] S. Adrián-Martínez et al., KM3NeT Coll., *Expansion cone for the 3-inch PMTs of the KM3NeT optical modules*. JINST **8**, T 03006 (2013).
- [33] A. Trovato, E. Drakopoulou and P. Sapienza for the KM3NeT Coll., in *Proc. 34th Int. Cosmic Ray Conference*, Vol. 001, 1114. PoS, 2015.
- [34] M. P. Kowalski, *Search for neutrino induced cascades with the AMANDA-II detector*. Ph.D. Thesis, Humbolt Universität Berlin, 2004.
- [35] J. Barrios-Martí, L. A Fusco, T. Heid, A. Heijboer and D. Stransky for the KM3NeT Coll., in *Proc. 34th Int. Cosmic Ray Conference*, Vol. 001, 1186. PoS, 2015.
- [36] T. K. Gaisser, K. Jero, A. Karle and J. van Santen, *Generalized self-veto probability for atmospheric neutrinos*. Phys. Rev. **D 90**, 023009 (2014), [arXiv:1405.0525 [astro-ph.HE]].
- [37] A. Hoecker et al., *TMVA: Toolkit for multivariate data analysis*. PoS **ACAT**, 040 (2007), [arXiv:physics/0703039].
- [38] D. Stransky, R. Coniglione and L. Fusco for the KM3NeT Coll., in *Proc. 34th Int. Cosmic Ray Conference*, Vol. 001, 1107. PoS, 2015.
- [39] S. Adrián-Martínez et al., KM3NeT Coll., *Detection potential of the KM3NeT detector for high-energy neutrinos from the Fermi bubbles*. Astropart. Phys. **42**, 7 (2013), [arXiv:1208.1226 [astro-ph.HE]].
- [40] M. G. Aartsen et al., IceCube Coll., *Observation of high-energy astrophysical neutrinos in three years of IceCube data*. Phys. Rev. Lett. **113**, 101101 (2014), [arXiv:1405.5303 [astro-ph.HE]].
- [41] M. G. Aartsen et al., IceCube Coll., *Atmospheric and astrophysical neutrinos above 1 TeV interacting in IceCube*. Phys. Rev. **D 91**, 022001 (2015), [arXiv:1410.1749 [astro-ph.HE]].
- [42] A. A. Abdo et al., Milagro Coll., *A measurement of the spatial distribution of diffuse TeV gamma-ray emission from the Galactic Plane with Milagro*. Astrophys. J. **688**, 1078 (2008), [arXiv:0805.0417 [astro-ph]].
- [43] F. Aharonian et al., H.E.S.S. Coll., *Discovery of very-high-energy γ -rays from the Galactic Centre ridge*. Nature **439**, 695 (2006), [arXiv:astro-ph/0603021].
- [44] M. Ackermann et al., Fermi-LAT Coll., *Fermi-LAT observations of the diffuse γ -ray emission: Implications for cosmic rays and the interstellar medium*. Astrophys. J. **750**, 3 (2012), [arXiv:1202.4039 [astro-ph.HE]].
- [45] M. Ahlers and K. Murase, *Probing the Galactic origin of the IceCube excess with gamma rays*. Phys. Rev. **D 90**, 023010 (2014), [arXiv:1309.4077 [astro-ph.HE]].
- [46] A. Neronov and D. V. Semikoz, *Neutrinos from Extra-Large Hadron Collider in the Milky Way*. Astropart. Phys. **72**, 32 (2016), [arXiv:1412.1690 [astro-ph.HE]].
- [47] D. Gaggero et al., *The Gamma-Ray and neutrino sky: A consistent picture of Fermi-LAT, milagro, and icecube results*. Astrophys. J. **815**, L 25 (2015), [arXiv:1504.00227 [astro-ph.HE]].

REFERENCES

- [48] M. Ahlers, Y. Bai, V. Barger and R. Lu, *Galactic TeV–PeV neutrinos*, 2015. arXiv:1505.03156 [hep-ph].
- [49] A. Neronov and D. V. Semikoz, *Evidence the Galactic contribution to the IceCube astrophysical neutrino flux*. *Astropart. Phys.* **75**, 60 (2016), [arXiv:1509.03522 [astro-ph.HE]].
- [50] D. Gaggero et al., *A hadronic scenario for the Galactic Ridge*, 2015. arXiv:1508.03681 [astro-ph.HE].
- [51] M. G. Aartsen et al., IceCube Coll., *Evidence for astrophysical muon neutrinos from the northern sky with icecube*. *Phys. Rev. Lett.* **115**, 081102 (2015), [arXiv:1507.04005 [astro-ph.HE]].
- [52] A. Kappes, J. Hinton, C. Stegmann and F. A. Aharonian, *Potential neutrino signals from galactic gamma-ray sources*. *Astrophys. J.* **656**, 870 (2007), [arXiv:astro-ph/0607286]. Erratum in *Astrophys. J.* 661 (2007) 1348.
- [53] F. Aharonian, J. Buckley, T. Kifune and G. Sinnis, *High energy astrophysics with ground-based gamma ray detectors*. *Rep. Prog. Phys.* **71**, 096901 (2008).
- [54] A. A. Abdo et al., Milagro Coll., *Milagro observations of multi-TeV emission from Galactic sources in the Fermi bright source list*. *Astrophys. J. Lett.* **700**, L127 (2009), [arXiv:0904:1018 [astro-ph.HE]].
- [55] K. Mannheim, *Neutrino signatures of the origins of cosmic rays*. *Nuc. Phys. B Proc. Supp.* **256**, 264 (2014).
- [56] F. Aharonian et al., H.E.S.S. Coll., *Primary particle acceleration above 100 TeV in the shell-type supernova remnant RX J1713.7-3946 with deep H.E.S.S. observations*. *Astron. Astrophys.* **464**, 235 (2007), [arXiv:astro-ph/0611813].
- [57] F. Aharonian et al., H.E.S.S. Coll., *Primary particle acceleration above 100 TeV in the shell-type supernova remnant RX J1713.7-3946 with deep H.E.S.S. observations (Corrigendum)*. *Astron. Astrophys.* **531**, C1 (2011).
- [58] S. R. Kelner, F. A. Aharonian and V. V. Bugayov, *Energy spectra of gamma rays, electrons, and neutrinos produced at proton–proton interactions in the very high energy regime*. *Phys. Rev.* **D 74**, 034018 (2006), [arXiv:0802.0562 [physics.ins-det]]. Erratum in *Phys. Rev.* **D 78** 039901 (2009).
- [59] J. Braun et al., *Methods for point source analysis in high energy neutrino telescopes*. *Astropart. Phys.* **29**, 299 (2008), [arXiv:0801.1604 [astro-ph]].
- [60] F. Aharonian et al., H.E.S.S. Coll., *First detection of a VHE gamma-ray spectral maximum from a cosmic source: HESS discovery of the Vela X nebula*. *Astron. Astrophys.* **448**, L43 (2006), [arXiv:astro-ph/0601575].
- [61] A. Abramowski et al., H.E.S.S. Coll., *Probing the extent of the non-thermal emission from the Vela X region at TeV energies with H.E.S.S.* *Astron. Astrophys.* **548**, A38 (2012), [arXiv:1210.1359 [astro-ph.HE]].
- [62] F. L. Villante and F. Vissani, *How precisely can neutrino emission from supernova remnants be constrained by gamma ray observations?* *Phys. Rev.* **D 78**, 103007 (2008).
- [63] F. Vissani and F. L. Villante, *Cosmic rays and neutrinos from supernova remnants (or: the time when H.E.S.S. met Ginzburg and Syrovatskii)*. *Nucl. Inst. Meth.* **A 588**, 123 (2008). Proc. First Int. Conf. Astroparticle Physics, RICAP 07.
- [64] F. Vissani, *Neutrinos from galactic sources of cosmic rays with known γ -ray spectra*. *Astropart. Phys.* **26**, 310 (2006).
- [65] M. G. Aartsen et al., IceCube Coll., *Searches for extended and point-like neutrino sources with four years of IceCube data*. *Astrophys. J.* **796**, 109 (2014).
- [66] S. Adrián-Martínez et al., ANTARES Coll., *Searches for point-like and extended neutrino sources close to the Galactic Center using the ANTARES neutrino telescope*. *Astrophys. J. Lett.* **786**, L5 (2014).
- [67] T. Michael for the ANTARES Coll., in *Proc. 34th Int. Cosmic Ray Conference*, Vol. 001, 1078. PoS, 2015.
- [68] P. Meszaros and M. J. Rees, *Relativistic fireballs and their impact on external matter - Models for cosmological gamma-ray bursts*. *Astrophys. J.* **405**, 278 (1993).

- [69] E. Waxman and J. Bahcall, *High energy neutrinos from cosmological gamma-ray burst fireballs*. Phys. Rev. Lett. **78**, 2292 (1997), [arXiv:astro-ph/9701231].
- [70] M. G. Aartsen et al., IceCube Coll., *Search for prompt neutrino emission from Gamma-Ray Bursts with IceCube*. Astrophys. J. **805**, L 5 (2015), [arXiv:1412.6510 [astro-ph.HE]].
- [71] S. Adrián-Martínez et al., ANTARES Coll., *Search for muon neutrinos from gamma-ray bursts with the ANTARES neutrino telescope using 2008 to 2011 data*. Astron. Astrophys. **559**, A 9 (2013), [arXiv:1307.0304 [astro-ph.HE]].
- [72] S. Adrián-Martínez et al., ANTARES Coll., *Optical and X-ray early follow-up of ANTARES neutrino alerts*, 2015. arXiv:1508.01180 [astro-ph.HE].
- [73] S. Adrián-Martínez et al., ANTARES Coll., *First results on dark matter annihilation in the Sun using the ANTARES neutrino telescope*. JCAP **1311**, 032 (2013), [arXiv:1302.6516 [astro-ph.HE]].
- [74] M. G. Aartsen et al., IceCube Coll., *Search for dark matter annihilations in the Sun with the 79-string IceCube detector*. Phys. Rev. Lett. **110**, 131302 (2013), [arXiv:1212.4097 [astro-ph.HE]].
- [75] S. Adrián-Martínez et al., ANTARES Coll., *Search of dark matter annihilation in the Galactic Centre using the ANTARES neutrino telescope*. JCAP **10**, 068 (2015), [arXiv:1505.04866 [astro-ph.HE]].
- [76] M. G. Aartsen et al., IceCube Coll., *Search for dark matter annihilation in the Galactic Center with IceCube-79*. Eur. Phys. J. **C 75**, 492 (2015), [arXiv:1505.07259 [astro-ph.HE]].
- [77] M. G. Aartsen et al., IceCube Coll., *Multipole analysis of IceCube data to search for dark matter accumulated in the Galactic halo*. Eur. Phys. J. **C 75**, 20 (2015), [arXiv:1408.0634 [astro-ph.HE]].
- [78] M. G. Aartsen et al., IceCube Coll., *IceCube search for dark matter annihilation in nearby galaxies and galaxy clusters*. Phys. Rev. **D 88**, 122001 (2013), [arXiv:1307.3473 [astro-ph.HE]].
- [79] B. Feldstein, A. Kusenko, S. Matsumoto and T. Yanagida, *Neutrinos at IceCube from Heavy Decaying Dark Matter*. Phys. Rev. **D88**, 015004 (2013), [arXiv:1303.7320 [hep-ph]].
- [80] A. Esmaili and P. D. Serpico, *Are IceCube neutrinos unveiling PeV-scale decaying dark matter?* JCAP **1311**, 054 (2013), [arXiv:1308.1105 [hep-ph]].
- [81] Zavala, Jesús, *Galactic PeV neutrinos from dark matter annihilation*. Phys. Rev. **D89**, 123516 (2014), [arXiv:1404.2932 [astro-ph.HE]].
- [82] K. Murase, R. Laha, S. Ando and M. Ahlers, *Testing the Dark Matter Scenario for PeV Neutrinos Observed in IceCube*. Phys. Rev. Lett. **115**, 071301 (2015), [arXiv:1503.04663 [hep-ph]].
- [83] S. Adrián-Martínez et al., ANTARES Coll., *Search for relativistic magnetic monopoles with the ANTARES neutrino telescope*. Astropart. Phys. **35**, 634 (2012), [arXiv:1110.2656 [astro-ph.HE]].
- [84] M. G. Aartsen et al., IceCube Coll., *Searches for Relativistic Magnetic Monopoles in IceCube*, 2015. arXiv:1511.01350 [astro-ph].
- [85] G. Riccobene et al., *Deep seawater inherent optical properties in the Southern Ionian Sea*. Astropart. Phys. **27**, 1 (2007), [arXiv:astro-ph/0603701].
- [86] V. Van Elewycck, P. Keller and M. Lindsey Clark for the KM3NeT Coll., in *Proc. 34th Int. Cosmic Ray Conference*, Vol. 001, 1160. PoS, 2015.
- [87] A. de Gouvea et al., Intensity Frontier Neutrino Working Group, *Working group report: Neutrinos*, 2013. arXiv:1310.4340 [hep-ex]. Community Summer Study 2013: Snowmass on the Mississippi (CSS2013) Minneapolis, MN, USA.
- [88] K. A. Olive et al., Particle Data Group, *Review of particle physics*. Chin. Phys. **C 38**, 090001 (2014).
- [89] F. Capozzi et al., *Status of three-neutrino oscillation parameters, circa 2013*. Phys. Rev. **D 89**, 093018 (2014), [arXiv:1312.2878 [hep-ph]].
- [90] M. C. Gonzalez-Garcia, M. Maltoni and T. Schwetz, *Updated fit to three neutrino mixing: status of leptonic CP violation*. JHEP **11**, 052 (2014), [arXiv:1409.5439 [hep-ph]].
- [91] D. V. Forero, M. Tortola and J. W. F. Valle, *Neutrino oscillations refitted*. Phys. Rev. **D 90**, 093006 (2014), [arXiv:1405.7540 [hep-ph]].
- [92] Y. Abe et al., Double Chooz Coll., *Indication for the disappearance of reactor electron antineutrinos in the Double Chooz experiment*. Phys. Rev. Lett. **108**, 131801 (2012), [arXiv:1112.6353 [hep-ex]].

REFERENCES

- [93] J. K. Ahn et al., RENO Coll., *Observation of reactor electron antineutrino disappearance in the RENO experiment*. Phys. Rev. Lett. **108**, 191802 (2012), [arXiv:1204.0626 [hep-ex]].
- [94] F. P. An et al., Daya Bay Coll., *Observation of electron-antineutrino disappearance at Daya Bay*. Phys. Rev. Lett. **108**, 171803 (2012), [arXiv:1203.1669 [hep-ex]].
- [95] W. Winter, *Neutrino mass hierarchy: Theory and phenomenology*. AIP Conf. Proc. **1666**, 120001 (2015). Proc. 26th International Conference on Neutrino Physics and Astrophysics (Neutrino 2014).
- [96] C. Lunardini, Cecilia and A. Yu. Smirnov, *Probing the neutrino mass hierarchy and the 13 mixing with supernovae*. JCAP **0306**, 009 (2003), [arXiv:hep-ph/0302033].
- [97] V. Barger, P. Huber and D. Marfatia, *Supernova neutrinos can tell us the neutrino mass hierarchy independently of flux models*. Phys. Lett. **B617**, 167 (2005), [arXiv:hep-ph/0501184].
- [98] Y.-F. Li, J. Cao, Y. Wang and L. Zhan, *Unambiguous determination of the neutrino mass hierarchy using reactor neutrinos*. Phys. Rev. **D88**, 013008 (2013), [arXiv:1303.6733 [hep-ex]].
- [99] R. B. Patterson for the NOvA Coll., *The NOvA experiment: Status and outlook*. Nucl. Phys. Proc. Suppl. **236-263**, 151 (2013), [arXiv:1209.0716 [hep-ex]]. Proc. 25th International Conference on Neutrino Physics and Astrophysics (Neutrino 2012).
- [100] K. Abe et al., T2K Coll., *Measurements of neutrino oscillation in appearance and disappearance channels by the T2K experiment with 6.6×10^{20} protons on target*. Phys. Rev. **D91**, 072010 (2015), [arXiv:1502.01550 [hep-ex]].
- [101] P. Adamson et al., *CHerenkov detectors In mine PitS (CHIPS) Letter of Intent to FNAL*, 2013. arXiv:1307.5918 [physics.ins-det]. Community Summer Study 2013: Snowmass on the Mississippi (CSS2013) Minneapolis, MN, USA.
- [102] C. Adams et al., LBNE Coll., *The long-baseline neutrino experiment: Exploring fundamental symmetries of the Universe*, 2013. arXiv:1307.7335 [hep-ex].
- [103] A. Stahl et al., *Expression of interest for a very long baseline neutrino oscillation experiment (LBNO)*, 2012. CERN-SPSC-2012-021, SPSC-EOI-007.
- [104] DUNE Coll. Available on <http://docs.dunescience.org/cgi-bin/ShowDocument?docid=180>.
- [105] A. Samanta, *The mass hierarchy with atmospheric neutrinos at INO*. Phys. Lett. **B673**, 37 (2009), [arXiv:hep-ph/0610196].
- [106] K. Abe et al., *Letter of Intent: The Hyper-Kamiokande experiment – detector design and physics potential –*, 2011. arXiv:1109.3262 [hep-ex].
- [107] M. G. Aartsen et al., IceCube PINGU Coll., *Letter of Intent: The Precision IceCube Next Generation Upgrade (PINGU)*, 2014. arXiv:1401.2046 [physics.ins-det].
- [108] J. Bernabeu, S. Palomares-Ruiz, A. Perez and S. T. Petcov, *The Earth mantle core effect in matter induced asymmetries for atmospheric neutrino oscillations*. Phys. Lett. **B531**, 90 (2002), [arXiv:hep-ph/0110071].
- [109] S. Palomares-Ruiz and S. T. Petcov, *Three-neutrino oscillations of atmospheric neutrinos, $\theta(13)$, neutrino mass hierarchy and iron magnetized detectors*. Nucl. Phys. **B712**, 392 (2005), [arXiv:hep-ph/0406096].
- [110] D. Indumathi and M. V. N. Murthy, *A Question of hierarchy: Matter effects with atmospheric neutrinos and anti-neutrinos*. Phys. Rev. **D71**, 013001 (2005), [arXiv:hep-ph/0407336].
- [111] R. Gandhi, P. Ghoshal, S. Goswami, P. Mehta and S. U. Sankar, *Earth matter effects at very long baselines and the neutrino mass hierarchy*. Phys. Rev. **D73**, 053001 (2006), [arXiv:hep-ph/0411252].
- [112] S. T. Petcov and T. Schwetz, *Determining the neutrino mass hierarchy with atmospheric neutrinos*. Nucl. Phys. **B740**, 1 (2006), [arXiv:hep-ph/0511277].
- [113] D. Indumathi, M. V. N. Murthy, G. Rajasekaran and N. Sinha, *Neutrino oscillation probabilities: Sensitivity to parameters*. Phys. Rev. **D74**, 053004 (2006), [arXiv:hep-ph/0603264].
- [114] R. Gandhi, P. Ghoshal, S. Goswami, P. Mehta, S. Sankar and S. Shalgar, *Mass Hierarchy Determination via future Atmospheric Neutrino Detectors*. Phys. Rev. **D76**, 073012 (2007), [arXiv:0707.1723 [hep-ph]].

- [115] R. Gandhi, Raj, P. Ghoshal, S. Goswami and S. U. Sankar, *Resolving the Mass Hierarchy with Atmospheric Neutrinos using a Liquid Argon Detector*. Phys. Rev. **D78**, 073001 (2008), [arXiv:0807.2759 [hep-ph]].
- [116] Samanta, Abhijit, *Discrimination of mass hierarchy with atmospheric neutrinos at a magnetized muon detector*. Phys. Rev. **D81**, 037302 (2010), [arXiv:0907.3540 [hep-ph]].
- [117] A. Samanta and A. Yu. Smirnov, *The 2-3 mixing and mass split: atmospheric neutrinos and magnetized spectrometers*. JHEP **07**, 048 (2011), [arXiv:1012.0360 [hep-ph]].
- [118] M. Blennow and T. Schwetz, *Identifying the Neutrino mass Ordering with INO and NOvA*. JHEP **08**, 058 (2012), [arXiv:1203.3388 [hep-ph]]. [Erratum: JHEP11,098(2012)].
- [119] A. Ghosh, T. Thakore and S. Choubey, *Determining the Neutrino Mass Hierarchy with INO, T2K, NOvA and Reactor Experiments*. JHEP **04**, 009 (2013), [arXiv:1212.1305 [hep-ph]].
- [120] A. Ghosh and S. Choubey, *Measuring the Mass Hierarchy with Muon and Hadron Events in Atmospheric Neutrino Experiments*. JHEP **10**, 174 (2013), [arXiv:1306.1423 [hep-ph]].
- [121] S. T. Petcov, *Diffraction-like (or parametric resonance-like?) enhancement of the Earth (day-night) effect for solar neutrinos crossing the Earth core*. Phys. Lett. **B 434**, 321 (1998), [arXiv:hep-ph/9805262].
- [122] E. K. Akhmedov, A. Dighe, P. Lipari and A. Yu. Smirnov, *Atmospheric neutrinos at Super-Kamiokande and parametric resonance in neutrino oscillations*. Nucl. Phys. **B542**, 3 (1999), [arXiv:hep-ph/9808270].
- [123] M. Chizhov, M. Maris and S. T. Petcov, *On the oscillation length resonance in the transitions of solar and atmospheric neutrinos crossing the earth core*, 1998. arXiv:hep-ph/9810501.
- [124] J. Bernabeu, S. Palomares Ruiz and S. T. Petcov, *Atmospheric neutrino oscillations, θ_{13} and neutrino mass hierarchy*. Nucl. Phys. **B669**, 255 (2003), [arXiv:hep-ph/0305152].
- [125] S. T. Petcov and S. Palomares-Ruiz, *On the atmospheric neutrino oscillations, θ_{13} and neutrino mass hierarchy*, 2004. arXiv:hep-ph/0406106.
- [126] P. Huber, M. Maltoni and T. Schwetz, *Resolving parameter degeneracies in long-baseline experiments by atmospheric neutrino data*. Phys. Rev. **D71**, 053006 (2005), [arXiv:hep-ph/0501037].
- [127] O. Mena, I. Mocioiu and S. Razzaque, *Neutrino mass hierarchy extraction using atmospheric neutrinos in ice*. Phys. Rev. **D 78**, 093003 (2008), [arXiv:0803.3044 [hep-ph]].
- [128] E. Kh. Akhmedov, S. Razzaque and A. Yu. Smirnov, *Mass hierarchy, 2-3 mixing and CP-phase with huge atmospheric neutrino detectors*. JHEP **02**, 082 (2013), [arXiv:1205.7071 [hep-ph]]. Erratum in JHEP 07 (2013) 026.
- [129] S. K. Agarwalla, T. Li, O. Mena and S. Palomares-Ruiz, *Exploring the Earth matter effect with atmospheric neutrinos in ice*, 2012. arXiv:1212.2238 [hep-ph].
- [130] D. Franco et al., *Mass hierarchy discrimination with atmospheric neutrinos in large volume ice/water Cherenkov detectors*. JHEP **04**, 008 (2013), [arXiv:1301.4332 [hep-ex]].
- [131] M. Ribordy and A. Y. Smirnov, *Improving the neutrino mass hierarchy identification with inelasticity measurement in PINGU and ORCA*. Phys. Rev. **D 87**, 113007 (2013), [arXiv:1303.0758 [hep-ph]].
- [132] W. Winter, *Neutrino mass hierarchy determination with IceCube-PINGU*. Phys. Rev. **D 88**, 013013 (2013), [arXiv:1305.5539 [hep-ph]].
- [133] M. Blennow and T. Schwetz, *Determination of the neutrino mass ordering by combining PINGU and Daya Bay II*. JHEP **09**, 089 (2013), [arXiv:1306.3988 [hep-ph]].
- [134] S.-F. Ge, K. Hagiwara and C. Rott, *A Novel Approach to Study Atmospheric Neutrino Oscillation*. JHEP **06**, 150 (2014), [arXiv:1309.3176 [hep-ph]].
- [135] S. Choubey and A. Ghosh, *Determining the Octant of θ_{23} with PINGU, T2K, NOvA and Reactor Data*. JHEP **11**, 166 (2013), [arXiv:1309.5760 [hep-ph]].
- [136] M. Blennow, P. Coloma, P. Huber and T. Schwetz, *Quantifying the sensitivity of oscillation experiments to the neutrino mass ordering*. JHEP **03**, 028 (2014), [arXiv:1311.1822 [hep-ph]].
- [137] S. Razzaque and A. Yu. Smirnov, *Super-PINGU for measurement of the leptonic CP-phase with atmospheric neutrinos*. JHEP **05**, 139 (2015), [arXiv:1406.1407 [hep-ph]].

REFERENCES

- [138] Capozzi, F. and Lisi, E. and Marrone, A., *PINGU and the neutrino mass hierarchy: Statistical and systematic aspects*. Phys. Rev. **D91**, 073011 (2015), [arXiv:1503.01999 [hep-ph]].
- [139] J. P. Yañez and A. Kouchner, *Measurement of atmospheric neutrino oscillations with very large volume neutrino telescopes*, 2015. arXiv:1509.08404 [hep-ex].
- [140] L. Wolfenstein, *Neutrino oscillations in matter*. Phys. Rev. **D 17**, 2369 (1978).
- [141] S. P. Mikheev and A. Yu. Smirnov, *Resonance amplification of oscillations in matter and spectroscopy of Solar neutrinos*. Sov. J. Nucl. Phys. **42**, 913 (1985).
- [142] Choubey, Sandhya and Roy, Probir, *Probing the deviation from maximal mixing of atmospheric neutrinos*. Phys. Rev. D **73**, 013006 (2006), [arXiv:hep-ph/0509197].
- [143] A. M. Dziewonski and D. L. Anderson, *Preliminary reference Earth model*. Phys. Earth Planet. Inter. **25**, 297 (1981).
- [144] E. K. Akhmedov, *Neutrino oscillations in inhomogeneous matter. (In Russian)*. Sov. J. Nucl. Phys. **47**, 301 (1988). [Yad. Fiz.47,475(1988)].
- [145] V. K. Ermilova, V. A. Tsarev, and V. A. Chechin, *Restoration of the Density Distribution of Material Based on Neutrino Oscillations*. Bull. Lebedev Phys. Inst. **1988N3**, 51 (1988). [Bull. Lebedev Phys. Inst.3,51(1988)].
- [146] P. I. Krastev and A. Yu. Smirnov, *Parametric Effects in Neutrino Oscillations*. Phys. Lett. **B226**, 341 (1989).
- [147] Q. Y. Liu and A. Yu. Smirnov, *Neutrino mass spectrum with muon-neutrino \rightarrow sterile-neutrino oscillations of atmospheric neutrinos*. Nucl. Phys. **B524**, 505 (1998), [arXiv:hep-ph/9712493].
- [148] E. K. Akhmedov, *Parametric resonance of neutrino oscillations and passage of solar and atmospheric neutrinos through the earth*. Nucl. Phys. **B538**, 25 (1999), [arXiv:hep-ph/9805272].
- [149] M. V. Chizhov and S. T. Petcov, *New conditions for a total neutrino conversion in a medium*. Phys. Rev. Lett. **83**, 1096 (1999), [arXiv:hep-ph/9903399].
- [150] M. V. Chizhov, and S. T. Petcov, *Enhancing mechanisms of neutrino transitions in a medium of nonperiodic constant density layers and in the earth*. Phys. Rev. **D63**, 073003 (2001), [arXiv:hep-ph/9903424].
- [151] E. K. Akhmedov, M. Maltoni and A. Yu. Smirnov, *Oscillations of high energy neutrinos in matter: Precise formalism and parametric resonance*. Phys. Rev. Lett. **95**, 211801 (2005), [arXiv:hep-ph/0506064].
- [152] E. K. Akhmedov, M. Maltoni and A. Yu. Smirnov, *1-3 leptonic mixing and the neutrino oscillograms of the Earth*. JHEP **05**, 077 (2007), [arXiv:hep-ph/0612285].
- [153] S. Adrián-Martínez et al., ANTARES Coll., *Measurement of atmospheric neutrino oscillations with the ANTARES neutrino telescope*. Phys. Lett. **B 714**, 224 (2012), [arXiv:1206.0745 [hep-ex]].
- [154] M. G. Aartsen et al., IceCube Coll., *Measurement of atmospheric neutrino oscillations with IceCube*. Phys. Rev. Lett. **111**, 081801 (2013), [arXiv:1305.3909 [hep-ex]].
- [155] J. A. Formaggio and G. P. Zeller, *From eV to EeV: Neutrino cross sections across energy scales*. Rev. Mod. Phys. **84**, 1307 (2012), [arXiv:1305.7513 [hep-ex]].
- [156] J. Hofestädt and C. W. James for the KM3NeT Coll., in *Proc. 34th Int. Cosmic Ray Conference*, Vol. 001, 1084. PoS, 2015.
- [157] X. Qian et al., *Statistical evaluation of experimental determinations of neutrino mass hierarchy*. Phys. Rev. **D 86**, 113011 (2012), [arXiv:1210.3651 [hep-ph]].
- [158] E. Ciuffoli, J. Evslin and X. Zhang, *Confidence in a neutrino mass hierarchy determination*. JHEP **01**, 095 (2014), [arXiv:1305.5150 [hep-ph]].
- [159] M. Blennow, *On the Bayesian approach to neutrino mass ordering*. JHEP **01**, 139 (2014), [arXiv:1311.3183 [hep-ph]].
- [160] A. Himmel for the Super-Kamiokande Coll., *Recent results from Super-Kamiokande*. AIP Conf. Proc. **1604**, 345 (2014), [arXiv:1310.6677 [hep-ex]]. Proc. 7th Int. Conf. Interconnection between Particle Physics and Cosmology (PPC 2013), Lead/Deadwood, SD, USA.
- [161] C. Andreopoulos et al., *The GENIE neutrino Monte Carlo generator*. Nucl. Inst. Meth. **A 614**, 87 (2010), [arXiv:0905.2517 [hep-ph]].

- [162] C. Andreopoulos et al., *The GENIE neutrino Monte Carlo generator: Physics and user manual*, 2015. arXiv:1510.05494 [hep-ph].
- [163] G. D. Barr et al., *A three-dimensional calculation of atmospheric neutrinos*. Phys. Rev. **D 70**, 023006 (2004), [arXiv:astro-ph/0403630].
- [164] A. G. Tsirigotis, A. Leisos and S. E. Tzamarias, *HOU reconstruction & simulation (HOURS): A complete simulation and reconstruction package for very large volume underwater neutrino telescopes*. Nucl. Inst. Meth. **A 626-627**, S 185 (2011).
- [165] Y. Becherini, A. Margiotta, M. Sioli and M. Spurio, *A parameterisation of single and multiple muons in the deep water or ice*. Astropart. Phys. **25**, 1 (2006), [arXiv:hep-ph/0507228].
- [166] C. Forti et al., *Simulation of atmospheric cascades and deep underground muons*. Phys. Rev. **D 42**, 3668 (1990).
- [167] J. Hofestädt, *Performance studies for the ORCA neutrino detector*. Ph.D. Thesis, Universität Erlangen-Nürnberg, Germany, 2016. To be published.
- [168] A. Heijboer, *Track reconstruction and point source searches with ANTARES*. Ph.D. Thesis, Universiteit van Amsterdam, The Netherlands, 2004.
- [169] C. Kopper, *Performance studies for the KM3NeT neutrino telescope*. Ph.D. Thesis, University of Erlangen-Nürnberg, Germany, 2010.
- [170] L. Breiman, *Random forests*. Machine Learning **45**, 5 (2001).
- [171] M. Honda et al., *Atmospheric neutrino flux calculation using the NRLMSISE-00 atmospheric model*. Phys. Rev. **D 92**, 023004 (2015), [arXiv:1502.03916 [astro-ph.HE]].
- [172] T. Ohlsson and H. Snellman, *Three flavor neutrino oscillations in matter*. J. Math. Phys. **41**, 2768 (2000), [arXiv:hep-ph/9910546]. Erratum in J. Math. Phys. **42** (2001) 2345.
- [173] F. P. An et al., Daya Bay Coll., *New measurement of antineutrino oscillation with the full detector configuration at Daya Bay*. Phys. Rev. Lett. **115**, 111802 (2015), [arXiv:1505.03456 [hep-ex]].
- [174] P. Adamson et al., MINOS Coll., *Combined analysis of ν_μ disappearance and $\nu_\mu \rightarrow \nu_e$ appearance in MINOS using accelerator and atmospheric neutrinos*. Phys. Rev. Lett. **112**, 191801 (2014), [arXiv:1403.0867 [hep-ex]].
- [175] S. Lein, Ph.D. Thesis, University of Minnesota, *Muon Neutrino Contained Disappearance in NOvA*, 2015. Available on <http://ccd.fnal.gov/techpubs/fermilab-reports-thesis.html>.
- [176] T. Ohlsson, H. Zhang and S. Zhou, *Effects of nonstandard neutrino interactions at PINGU*. Phys. Rev. **D 88**, 013001 (2013), [arXiv:1303.6130 [hep-ph]].
- [177] M. C. Gonzalez-Garcia, F. Halzen and M. Maltoni, *Physics reach of high-energy and high-statistics IceCube atmospheric neutrino data*. Phys. Rev. **D 71**, 093010 (2005), [arXiv:hep-ph/0502223].
- [178] J. Brunner, *Counting electrons to probe the neutrino mass hierarchy*, 2013. arXiv:1304.6230 [hep-ex].
- [179] J. Brunner, *Measurement of neutrino oscillations with neutrino telescopes*. Adv. High Energy Phys. **2013**, 782538 (2013).
- [180] S. Bourret, *Sensitivity to the neutrino mass hierarchy and discovery of CP violation with the Protvino-to-ORCA proposal*, 2015. M. Sc. Thesis, ETH Zürich (Switzerland) and Ecole Polytechnique (France).
- [181] K. Asano and K. Murase, *Gamma-Ray Bursts as Multienergy Neutrino Sources*. Adv. Astron. **2015**, 568516 (2015).
- [182] J. Becker Tjus, *Neutrinos from Colliding Wind Binaries: Future Prospects for PINGU and ORCA*, 2014. arXiv:1405.0471 [astro-ph.HE].
- [183] W. Winter, *Neutrino tomography: Learning about the Earth's interior using the propagation of neutrinos*. Earth Moon Planets **99**, 285 (2006), [arXiv:physics/0602049]. Proc. Neutrino Sciences 2005, Honolulu, USA.
- [184] C. Rott, A. Taketa and D. Bose, *Spectrometry of the Earth using neutrino oscillations*. Sci. Rep. **5**, 15225 (2015), [arXiv:1502.04930 [physics.geo-ph]].
- [185] W. Winter, *Atmospheric neutrino oscillations for Earth tomography*, 2015. arXiv:1511.05154 [hep-ph].